

See discussions, stats, and author profiles for this publication at:
<https://www.researchgate.net/publication/265089771>

Microbial Acceleration of Olivine Dissolution via Siderophore Production

ARTICLE · DECEMBER 2014

DOI: 10.1016/j.proeps.2014.08.041

CITATION

1

READS

23

3 AUTHORS, INCLUDING:



A. Joshua West

University of Southern California

60 PUBLICATIONS 1,140 CITATIONS

SEE PROFILE



Kenneth H Neelson

University of Southern California

398 PUBLICATIONS 22,157 CITATIONS

SEE PROFILE

Geochemistry of the Earth's Surface meeting, GES-10

Microbial acceleration of olivine dissolution via siderophore production

Mark A. Torres^{a*}, A. Joshua West^a, Kenneth Nealson^a^aUniversity of Southern California - Department of Earth Sciences, 3651 Trousdale Parkway, Los Angeles 90089, United States of America

Abstract

Silicate minerals represent an important reservoir of essential nutrients at Earth's surface. Due to the slow kinetics of primary silicate mineral dissolution and the potential for nutrient sequestration by secondary mineral precipitation, the bioavailability of many silicate-bound nutrients may be limited by the ability of microorganisms to actively scavenge these nutrients via organic ligand production. In this study, the effectiveness of ligand production as a means to scavenge Fe from Fe-silicates is addressed through targeted laboratory experiments using olivine as a model mineral.

Preliminary results show that microbial Fe-binding ligands (i.e. siderophores) can accelerate olivine dissolution rates stoichiometrically by almost an order of magnitude in experiments buffered at circumneutral pH. In addition to higher reaction rates, organic Fe-binding ligands fostered the accumulation of dissolved Fe in solution, which was below detection in the abiotic experiments due to the precipitation of secondary Fe minerals in the presence of O₂. Accelerated olivine dissolution rates in the presence of microbial Fe-binding ligands is somewhat unexpected because these ligands are known to be highly selective towards Fe³⁺ whereas olivine dominantly contains Fe²⁺. Spectrophotometric analysis of the ligand complexes produced during reaction with olivine reveals the dominance for Fe³⁺-ligand complexes in solution.

© 2014 Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

Peer-review under responsibility of the Scientific Committee of GES-10

*marktorr@usc.edu

Keywords: Mineral dissolution kinetics; siderophores; microbial weathering

1. Introduction

The biological demand for Fe and other trace metals despite their low solubility in many Earth surface environments is thought to be met via the production of strong and highly selective ligands known as siderophores¹. These molecules bind to trace metals, facilitating their biological uptake. The hypothesis that siderophores play a

key role in mobilizing lithogenic nutrients is corroborated by much experimental work showing that the production of these ligands by microorganisms is induced in response to metal deficiencies and that the metal-ligand complexes are actively taken up into living cells²⁻³. Furthermore, siderophores have been measured in natural soils and soil porewaters⁴⁻⁵.

Understanding the role of siderophores in biogeochemical cycles requires information about how they interact with mineral substrates. To date, most experimental work has focused on the interactions between siderophores and aqueous species⁶⁻⁷ as well as secondary Fe³⁺-bearing phases⁸. Comparatively, less is known about the interactions between siderophores and primary Fe-bearing phases (e.g. silicates), especially those that do not contain Al³⁺ (e.g. ref. 9), which is also known to be strongly complexed by siderophores¹⁰. In this work, olivine ((Mg,Fe)₂SiO₄) is used as a model mineral to investigate the interaction between siderophores and Fe-bearing primary silicates in laboratory dissolution experiments.

2. Methods

Large (~1-3 mm) olivine crystals were hand picked underneath a binocular microscope in order to avoid grains with weathered surfaces or mineral inclusions. The un-weathered grains were crushed in an agate mortar and pestle, dry sieved into different grain size fractions, and cleaned ultrasonically with ethanol in order to remove fine-grained particles produced during grinding. Dissolution experiments were performed in both batch and mixed flow systems with a defined grain size of olivine (75-175 µm) and varying concentrations (i.e. 0 – 280 µM) of the commercially available hydroxamate siderophore deferoxamine. Batch experiments were buffered to a pH of ~7.5 with a TRIS/HCl buffer (10 mM) and reacted at 30°C in an incubated shaking table that kept the solutions well mixed by rotating at 200 rpm for the duration of the experiments. The batch experiments were sampled regularly for up to 40 hours and analyzed for Mg, Fe, and Si concentrations by MP-OES after acidification with concentrated HCl.

Experiments in mixed-flow reactors were performed using a flow rate of either 0.1 or 0.2 mL/min in order to allow for fluid residence times of 4 or 8 hours respectively. Constant solution volume in the reactors was maintained by pressurizing the reactors with laboratory air so that fluid exited the reactor upon reaching the height of the outlet tubing. The solution pH was held constant at a value of ~7.5 with either a TRIS/HCl buffer (10 mM) or a phosphate/NaOH buffer (500 µM). The mixed flow experiments were performed in an incubated shaking table set at the same temperature and rotation speed as in the batch experiments. The reactors were sampled regularly until a steady state was reached and analyzed for Mg, Fe, and Si concentrations by MP-OES after acidification with concentrated HCl. In addition to concentration measurements, ligand speciation was monitored in selected samples by measuring the solution absorbance at wavelengths between 200 and 600 nm with a spectrophotometer.

3. Results

In batch experiments, measured olivine dissolution rates increase non-linearly with increasing deferoxamine concentrations in a similar manner to that seen with the mineral hornblende (Fig. 1; ref. 9). However, a greater enhancement in dissolution rate at lower deferoxamine concentrations is seen for olivine relative to hornblende. Additionally, olivine Mg, Si, and Fe release rates are all increased stoichiometrically by deferoxamine, which is not seen for the constituent ions of hornblende⁹. At deferoxamine concentrations of ≥200 µM, dissolution rates appear to plateau at a value that is almost an order of magnitude greater than the dissolution rate measured in the absence of any ligands (Fig. 1).

Generally, the rates measured in the mixed-flow experiments are consistent with those derived from the batch experiments (Fig. 1). Ligand speciation analysis of the outlet fluids from the mixed flow reactors reveals the dominance of Fe^{3+} complexes due to the presence of an absorbance peak at around 430 nm that can be re-produced when Fe^{3+} is added to the initial solution, but not when either Mg^{2+} or Si^{4+} are added (Fig. 2).

4. Discussion

The ability of the siderophore deferoxamine to appreciably accelerate olivine dissolution rates at circumneutral pH is interesting in that olivine dominantly contains Fe^{2+} whereas deferoxamine is highly selective towards binding Fe^{3+} . Three mechanisms could potentially explain this observation. If secondary Fe-oxide phases inhibit olivine dissolution by armoring the mineral surface, then the presence of a strong Fe^{3+} binding ligand could remove these phases and act to increase the reactive surface area of olivine. In the ligand-free batch experiments, secondary Fe-oxides are likely precipitating as the concentration of Fe is below the analytical detection limit. However, the removal of secondary Fe-oxide phases is unlikely to be the dominant mechanism contributing to the accelerated olivine dissolution rates measured in our system as ligand-free dissolution rates do not decrease with time after stoichiometric dissolution with respect to Mg and Si is reached.

Another possibility is that adsorbed deferoxamine accelerates the oxidation of Fe^{2+} at the mineral surface and then the strong capacity for deferoxamine to bind Fe^{3+} leads to the accelerated dissolution by facilitating bond breakage. This is similar to the mechanism proposed by ref. 11 to describe the effect of Mg^{2+} complexation by oxalate on olivine dissolution rates. This mechanism would explain the observed production of Fe^{3+} - ligand complexes (Fig. 2). Alternatively, the lower affinity (i.e. 10^{10} v.s. 10^{30} ; ref. 7) of deferoxamine for Fe^{2+} may still be high enough to accelerate olivine dissolution rates by binding Fe^{2+} and facilitating bond breakage. If this is the case, then the oxidation of Fe^{2+} to Fe^{3+} must occur in solution in order to produce the observed Fe^{3+} complexes. While subtly different, these mechanisms have implications for how siderophores react with ferrous silicates in natural environments, because variations in pO_2 strongly affect Fe oxidation kinetics¹². Further work will seek to identify which mechanism dominates the interaction between olivine and deferoxamine.

References

1. Sandy, M. & Butler, A. Microbial iron acquisition: marine and terrestrial siderophores. *Chem. Rev.* 109, 4580–95 (2009).
2. Müller, G. & Raymond, K. N. Specificity and mechanism of ferrioxamine-mediated iron transport in *Streptomyces pilosus*. *J. Bacteriol.* 160, 304–12 (1984).
3. Bellenger, J. P., Wichard, T., Kustka, a. B. & Kraepiel, a. M. L. Uptake of molybdenum and vanadium by a nitrogen-fixing soil bacterium using siderophores. *Nat. Geosci.* 1, 243–246 (2008).
4. Holmström, S. & Lundström, U. Siderophores in forest soil solution. *Biogeochemistry* 71, 247–258 (2004).
5. Essén, S. a, Bylund, D., Holmström, S. J. M., Moberg, M. & Lundström, U. S. Quantification of hydroxamate siderophores in soil solutions of podzolic soil profiles in Sweden. *Biometals* 19, 269–82 (2006).
6. Enyedy, A., Csoka, H. & Farkas, E. A comparison between the chelating properties of some dihydroxamic acids, desferrioxamine B and acetohydroxamic acid. 18, 2391–2398 (1999).
7. Spasojevic, I., Armstrong, S. K., Brickman, T. J. & Crumbliss, A. L. Electrochemical Behavior of the Fe(III) Complexes of the Cyclic Hydroxamate Siderophores Alcaligin and Desferrioxamine E. *Inorg. Chem.* 38, 449–454 (1999).
8. Reichard, P. U., Kretzschmar, R. & Kraemer, S. M. Dissolution mechanisms of goethite in the presence of siderophores and organic acids. *Geochim. Cosmochim. Acta* 71, 5635–5650 (2007).

9. Liermann, L. J., Kalinowski, B. E., Brantley, S. L. & Ferry, J. G. Role of bacterial siderophores in dissolution of hornblende. *Geochim. Cosmochim. Acta* 64, 587–602 (2000).
10. Evers, A., Hancock, R. D., Martell, A. E. & Motekaitis, R. J. Metal Ion Recognition in Ligands with Negatively Charged Oxygen Donor Groups. Complexation of Fe(III), Ga(III), In(III), Al(III), and Other Highly Charged Metal Ions. *Inorg. Chem.* 28, 2189–2195 (1989).
11. Olsen, A. A. & Rimstidt, J. D. Oxalate-promoted forsterite dissolution at low pH. *Geochim. Cosmochim. Acta* 72, 1758–1766 (2008).
12. Sugimori, H., Kanzaki, Y. & Murakami, T. Relationships between Fe redistribution and pO_2 during mineral dissolution under low O_2 conditions. *Geochim. Cosmochim. Acta* 84, 29–46 (2012).




Figure 1. Silicon release rates from olivine in the presence of deferoxamine at pH ~7.5.




Figure 2. Absorbance spectra of Si^{4+} -DFOB, Mg^{2+} -DFOB, Fe^{3+} -DFOB, and DFOB reacted with olivine