

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/255764624>

ChemInform Abstract: Recent Advances in New Multicomponent Synthesis of Structurally Diversified 1,4-Dihydropyridines

ARTICLE *in* RSC ADVANCES · OCTOBER 2012

Impact Factor: 3.84 · DOI: 10.1039/C2RA21406G

CITATIONS

55

READS

272

2 AUTHORS, INCLUDING:



Jieping Wan

Jiangxi Normal University

82 PUBLICATIONS 927 CITATIONS

SEE PROFILE

Cite this: *RSC Advances*, 2012, **2**, 9763–9777

www.rsc.org/advances

REVIEW

Recent advances in new multicomponent synthesis of structurally diversified 1,4-dihydropyridines

Jie-Ping Wan* and Yunyun Liu

Received 6th May 2012, Accepted 8th August 2012

DOI: 10.1039/c2ra21406g

1,4-Dihydropyridines (DHPs) are one of the most important classes of heterocyclic compounds as privileged pharmacophores. They are therefore attractive synthetic targets of organic chemistry. Multicomponent reactions (MCRs) are the most efficient strategies for the synthesis of 1,4-DHPs in terms of providing both sufficient structural diversity and numbers for compound libraries. Following the classical multicomponent synthesis of 1,4-DHPs by the Hantzsch reaction, chemists have developed a large number of new MCRs to access 1,4-DHPs with significantly extended structural diversity, which is pivotal for the process of discovering new lead compounds and drugs of 1,4-DHPs. The advances on the synthesis of structurally diversified 1,4-DHPs through new MCRs beyond the classical Hantzsch reaction is reviewed.

1. Introduction

Among the known strategies of drug discovery, high throughput screening is obviously the one that is serving most efficiently to the pharmaceutical industry. To guarantee the success on acquiring lead compounds *via* this kind of screening tactic, sources of large amount of molecular libraries are important preconditions.¹ To satisfy such requirements, new synthetic strategies with robust efficiency in providing a massive amount

of structurally diverse organic products are the only solution. MCRs which employ three or more reactants to furnish products containing structure or substructure of all starting materials in one-pot and one set of fixed reaction conditions are such examples with powerful productivity.² The Hantzsch reaction (Scheme 1) which provides 1,4-DHPs as products is one of the earliest and most well-known MCRs owing to the excellent pharmaceutical profile of 1,4-DHPs.³ The most typical examples of 1,4-DHPs pharmaceuticals are those broadly prescribed calcium channel blockers, for example, Nifedipine (**1**), Felodipine (**2**), Nicardipine (**3**) and Nimodipine (**4**), to name but a few (Scheme 2). In addition, other versatile biological profiles of 1,4-DHPs such as anticonvulsant activity,⁴ selective

*Key Laboratory of Functional Small Organic Molecules, Ministry of Education and College of Chemistry and Chemical Engineering, Jiangxi Normal University, Nanchang, 330022, P. R. China.
E-mail: wanjieping@gmail.com; Fax: +86 791 88120380*




Jie-Ping Wan

Dr Jie-Ping Wan obtained his Bachelor's degree by studying chemistry in Nanchang University in 2000–2004. He continued his graduate study in Zhejiang University in 2005–2010 in Professor Yuanjiang Pan's group and obtained his Ph.D degree in 2010. In July, 2010, he joined the College of Chemistry and Chemical Engineering in Jiangxi Normal University as an assistant professor. In September 2011, under the support of Chinese-Deutsche Zentrum für Wissenschaftsförderung Dr Wan joined Prof. Dieter Enders group in RWTH Aachen University as a postdoctoral fellow working in organocatalysis. His present research interest contains multicomponent reactions, organocatalysis and metal-catalyzed synthesis.



Yunyun Liu


Dr Yunyun Liu was born in 1983 in Shandong Province, China. She obtained her Bachelor Degree in Qufu Normal University in 2005. She then moved to Zhejiang University to continue her graduate study in the Department of Chemistry. Under the supervision of Professor Weiliang Bao, she worked in the field of copper-catalyzed Ullmann coupling reaction and related tandem reactions for her doctorate study. She obtained her doctorate degree in 2010 and presently is an assistant professor in Jiangxi Normal University. She is currently interested in the research of metal-catalyzed organic synthesis and the development of new cascade organic reactions.



Scheme 1 Classical Hantzsch reaction.

adenosine-A3 receptor antagonism,⁵ radioprotective activity,⁶ sirtuin activation and inhibition,⁷ *etc.* have also been reported. Therefore, it is not surprising that 1,4-DHPs receive daily increasing interests as synthetic targets.

Conventionally, 1,4-DHPs could be accessed *via* the Hantzsch reaction, reduction of pyridines, addition to pyridines or cycloadditions *etc.*⁸ As a facile and broadly tolerable protocol, the Hantzsch reaction remains as a frequently employed tactic for the synthesis of 1,4-DHPs in a large number of areas such as stereoselective synthesis and green chemistry.⁹ On the other hand, following the rapid progress of chemical and pharmaceutical industries, some limits have also been noticed on this classical multicomponent synthetic route. A main limit of the Hantzsch reaction is that the 1,4-DHPs provided by the reaction are all symmetrical in the heterocyclic fragment since this unit was constructed by employing two molecules of 1,3-dicarbonyl compounds. Although Hantzsch reactions using two different dicarbonyl compounds had been later achieved, the symmetrical 1,4-DHPs formed by the homo-condensation of same dicarbonyl compound remained more or less as side products in these entries. In terms of biological screening, acquiring diversified derivatives of a known pharmacophore is a main and efficient approach for discovering new biologically active lead compounds.¹⁰ Even though many reported biologically active 1,4-DHPs are symmetrical, this was not the requirement of biological receptors, but the results of employing the Hantzsch reaction for 1,4-DHPs synthesis in most cases. For the sake of finding more 1,4-DHPs possessing new and improved biological functions, the synthesis of structurally diverse 1,4-DHPs such as unsymmetrical 1,4-DHPs is pivotal work. Therefore, new MCRs




Scheme 2 1,4-DHPs used as clinical drugs.


that are capable of generating novel 1,4-DHPs have attracted significant attention in recent years. Considering the merits of 1,4-DHPs for the development of chemical biology and medicinal chemistry as well as the impressive progress being made in this field, it is therefore necessary to summarize the research advances on the multicomponent synthesis of 1,4-DHPs with different new MCRs. We present herein the first review on the advances of 1,4-DHPs synthesis with new MCRs beyond the Hantzsch reaction. Based on the analysis of all known literature on this topic, we classified these reactions into different sections according to the amino sources employed for the construction of the dihydropyridine ring. The main content of this review covered the period from 2004 to 2012.

2. Multicomponent synthesis of 1,4-DHPs using amine as amino source

Amines are obviously the most easily available and widely utilized amino sources in organic synthesis. They have been frequently employed as the amine component in the classical Hantzsch reaction. Similarly, amines are also favorable amino sources in the devisal of new multicomponent synthesis of 1,4-DHPs. For example, employing α,β -unsaturated aldehydes, 1,3-dicarbonyl compounds and amines has been found as a very practical approach for the synthesis of unsymmetrical 1,4-DHPs. During their work of synthesizing 1,4-DHPs **6** by two-component condensation of β -enamino esters **5** and α,β -unsaturated aldehydes, Renaud *et al.* conducted tentative investigations on direct three-component reaction of 1,3-dicarbonyl compounds, amines and α,β -unsaturated aldehydes to access 1,4-DHPs with Lewis acid catalysis. Moderate to excellent yields have been obtained by performing the reactions in a stepwise manner (Scheme 3).¹¹ Later on, the one step, three-component reaction of a 1,3-dicarbonyl compound, primary amine and α,β -unsaturated aldehyde has been realized by Menéndez and coworkers. Their method was established by employing cerium ammonium nitrate (CAN) as a Lewis acid catalyst. This protocol has been demonstrated with excellent substrate tolerance by providing a sound number of 1,4-DHPs of type **6** with moderate to good yields at room temperature. And a noteworthy feature of this methodology was that β -ketothioesters could be efficiently employed as the dicarbonyl component to correspondingly give 1,4-DHPs containing a reactive thioester group as shown by typical results outlined in Scheme 4.¹² In addition, as a widely employed catalysis tactic, organocatalysis had also turned out to be practical for this kind of three-component 1,4-DHPs synthesis



Scheme 3 Three-component stepwise synthesis of 1,4-DHPs.




Scheme 4 Typical results on enal-based synthesis of unsymmetrical 1,4-DHPs.

according to the results presented by Kumar *et al.* While a series of different organocatalysts such as (–)-cinchonidine, (–)-ephedrine, L-lysine, L-histidine, L-glutamic acid, L-aspartic acid, L-pipecolic acid and L-proline have been screened, L-proline displayed the best activity as an organocatalyst. By performing the reactions of different α,β -unsaturated aldehydes, amines and dicarbonyl compounds at room temperature and solvent free conditions, various 1,4-DHPs of type **6** have been obtained with generally excellent yields. For example, the reaction of cinnamaldehyde, aniline and 2-acetyl ethyl acetate provided the corresponding 1,4-DHP in 90% yield.¹³ Following these results, some other different catalysts such as nano CuO ,¹⁴ silica-supported perchloric acid ($HClO_4 \cdot SiO_2$)¹⁵ and $InCl_3$ ¹⁶ were also discovered as practical catalysts for this three-component 1,4-DHPs synthesis.

One of the most important properties of unsymmetrical 1,4-DHPs is the chiral center embedded in the central heterocycle. Therefore, performing this kind of multicomponent reactions in an enantioselective manner is of great significance. Few methods of asymmetric catalysis on 1,4-DHPs synthesis are currently available. The representative results on this area was reported by Gong and coworkers, in which a chiral phosphoric acid was used as a Brønsted acid to catalyze the three-component reaction of α,β -unsaturated aldehydes, amines and dicarbonyl compounds giving (*S*)-**6** with good to excellent enantioselectivity in general (66–97% ee) and fair to good yields (31–93%). Some typical

results on the synthesized chiral 1,4-DHPs of type **6** are shown in Scheme 5.¹⁷ Notably, elegant results on the enantioselective synthesis of 1,4-DHPs by the Hantzsch reaction has also recently been accomplished by Gestwicki and co-workers employing BINOL-phosphoric acid catalysis.¹⁸

Alkynes were another type of frequently employed building block in the multicomponent synthesis of 1,4-DHPs of both symmetrical and unsymmetrical structures. For example, Wang, *et al.* developed an interesting multicomponent synthesis of 1,4-DHPs of type **7** via TFA-catalyzed assembly of 3 molecules of *E*-3-(2-formylphenoxy)propenoates **8** and amines (Scheme 6). Corresponding salicaldehydes were generated as side products from **8** after incorporating amines and two ethyl propiolate building blocks. The salicaldehydes were able to be regenerated to **8** by incorporating ethyl propiolate. This transformation has been found to be specifically applicable for the starting materials of type **8** since no corresponding reactions were observed when using either the *para*-substituted isomers of **8** to incorporate amines or direct three-component reactions of salicaldehyde, aniline and ethyl propiolate/ethyl 3-phenoxyacrylate under standard reaction conditions.¹⁹ However, it was later discovered that the direct MCRs of ethyl propiolate, benzaldehyde and a primary amine were capable of providing corresponding 1,4-DHPs **9** by heating the starting materials in acetic acid. The application scope of this protocol, however, was not defined, as only 4 examples of 1,4-DHPs **9** were prepared for the sake of biological research (Scheme 7).⁷ Also employing propiolates as building blocks for the dihydropyridine ring construction, Yan *et al.* established a four-component synthesis of 1,4-DHPs **10** from methyl propiolate, aryl amines, aromatic aldehydes and acetonitriles. With the catalysis of triethyl amine, the Knoevenagel condensed intermediates **11** incorporated intermediates **12** from an aza-Michael addition to provide intermediates **13** which underwent subsequent intramolecular cyclization and tautomerization to 1,4-DHPs **10** as final products. On the other hand, when ethyl 2-cyanoacetate was used as the starting material in the place of malonitrile/2-cyanoacetamide, 2-pyridinones **15** were afforded as main or single products in the corresponding entries (Scheme 8).²⁰ As alkyne substrates with similar reactivity as alkyl propiolates, dialkyl acetylenedicarboxylates were also found to be versatile materials for the synthesis of novel functionalized 1,4-DHPs. For example, under the catalysis of organocatalyst L-proline, acetylene-dicarboxylates **16** efficiently reacted with amines to generate β -enamino esters of type **17**. Based on this transformation, Jiang and coworkers designed an L-proline catalyzed multicomponent synthetic method starting from acetylenedicarboxylates (alkyl propiolates as well), amines, aldehydes and 1,3-dicarbonyl compounds to synthesize 1,4-DHPs of type **21**. As shown in Scheme 9, active intermediates **17** generated from the addition of amines to electron deficient alkynes were believed to directly couple with Knoevenagel condensed intermediates **18** to form **19/20**, which were efficiently transformed to 1,4-DHPs **21** via further intramolecular condensation. Mild reaction conditions and wide substrate tolerance on all starting materials were advantages in this method.²¹ Similar to acyclic 1,3-dicarbonyl compounds, cyclic 1,3-dicarbonyl compounds also reacted with aromatic amines, aromatic aldehydes and dimethyl acetylenedicarboxylate to give corresponding 1,4-DHPs **22** in a fused




R ¹	R ²	R ³	Ar	Yield (%)	ee (%)
4-NO ₂ C ₆ H ₄	Me	OEt	4-MeOC ₆ H ₄	82	92
4-NO ₂ C ₆ H ₄	Me	OEt	3-MeOC ₆ H ₄	53	97
4-NO ₂ C ₆ H ₄	Me	OEt	4-ClC ₆ H ₄	60	89
Ph	Me	OPr- <i>i</i>	4-MeOC ₆ H ₄	40	81
2,3-Cl ₂ C ₆ H ₃	Me	OEt	4-MeOC ₆ H ₄	72	87
2,3-Cl ₂ C ₆ H ₃	Me	OEt	3-MeOC ₆ H ₄	72	96
3-NO ₂ C ₆ H ₄	Me	OEt	4-MeOC ₆ H ₄	70	91
2-NO ₂ C ₆ H ₄	Me	OEt	4-MeOC ₆ H ₄	72	94
2-BrC ₆ H ₄	Me	Me	4-MeOC ₆ H ₄	63	73
4-NO ₂ C ₆ H ₄	Et	OMe	4-MeOC ₆ H ₄	38	91

Scheme 5 Enal-based enantioselective synthesis of 1,4-DHPs.


structure by employing HOAc as solvent as well as acid catalyst (Scheme 10).²²

Instead of enone intermediates **18**, α,β -unsaturated aldehydes could be directly employed as building blocks to prepare 1,4-DHPs by reacting with acetylenedicarboxylates and primary amines. In the presence of TFA or TfOH, 1,4-DHPs **23** were easily afforded by the three-component reactions of dimethyl acetylenedicarboxylate, amines and α,β -unsaturated aldehydes at room temperature in THF. 1,2-DHPs **24** were formed as minor side products in these reactions. Based on the proposed reaction




Scheme 6 Three-component synthesis of 1,4-DHPs initiated by *E*-3-(2-formylphenoxy)propenoates.


mechanism, the initial N-nucleophilic site in β -enamino esters attack the different electrophilic sites in α,β -unsaturated aldehydes was believed to determine the regioselectivity. N-attacking led to the formation of 1,4-DHPs, while minor 1,2-DHPs products was proposed to be provided by the aza-Michael addition of the N-nucleophilic site to enals (Scheme 11).²³ Interestingly, the homo-condensation of acetylenedicarboxylates with amines and aldehydes were also capable of delivering 1,4-DHPs. Yan *et al.* discovered the synthesis of fully substituted 1,4-DHPs **25** by directly reacting acetylenedicarboxylates with aromatic amines and aromatic aldehydes in absolute ethanol. When 95% EtOH was employed, 2-hydroxyhydropyridines **26** were obtained diastereoselectively probably by the addition of H₂O to **25** (Scheme 12).²⁴



Scheme 7 Direct synthesis of symmetrical 1,4-DHPs from propiolate.



Scheme 8 Four-component synthesis of 1,4-DHPs with aldehyde, amine, propiolate and nitrile.




Scheme 9 Synthesis of fused 1,4-DHPs using cyclic diketones.


Recently, Kumar *et al.* discovered that malonitrile or ethyl cyanoacetate was able to efficiently react with diethyl acetylenedicarboxylate, aromatic amines and aromatic aldehydes to provide 1,4-DHPs of type **27** by a solvent- and catalyst-free

grinding protocol. This clean synthetic method proceeded also *via* enamino esters and Knoevenagel condensed intermediates (Scheme 13).²⁵


Besides those aforementioned alkynes, alkenes have also been found as applicable substrates in 1,4-DHPs synthesis. In 2011, Jiang *et al.* reported an interesting multicomponent synthesis of 1,4-DHPs directly employing aldehydes, amines and electron deficient alkenes **28** with palladium(II) catalysis in O₂ atmosphere. The 1,4-DHPs of type **29** were given in moderate to good yields. Compared with benzaldehyde derivatives, heteroaromatic aldehydes were not good reaction partners as lower yields of the corresponding products were afforded (Scheme 14).²⁶ Another multicomponent synthesis of 2,6-unsubstituted 1,4-DHPs of type **29** employing electron rich alkenes **30** as precursors of enamino intermediates **31** was reported recently. In the presence of the Lewis acid Sc(OTf)₃, corresponding 1,4-DHPs were afforded in fair to good yields. However, 2–19 days of




Scheme 10 Synthesis of 1,4-DHPs using aldehyde, amine, a 1,3-dicarbonyl compound and electron deficient alkynes.




Scheme 11 Regioselective three-component synthesis of 1,4-DHPs using dimethylacetylenedicarboxylate, enal and amine.



Scheme 12 Regioselective three-component synthesis of 1,4-DHPs and tetrahydropyridines.



Scheme 13 Solvent- and catalyst-free synthesis of 1,4-DHPs.




Scheme 15 Alkoxy activated alkene initiated synthesis of 1,4-DHPs.

reaction time and the presence of side products **31** as well as other types of side products in the reactions were the main restrictions of this method (Scheme 15).²⁷


As reactants possessing analogous reactivity with **30**, the β-ester acetals **32** could also be employed as C-nucleophile sources for 1,4-DHP synthesis by serving as the precursors of β-enamino esters. The *in situ* generated β-enamino esters from starting acetals and aromatic amines efficiently reacted with aldehydes in the manner of homo-condensation to provide 1,4-DHPs **33** by heating at 90 °C in dioxane in the presence of Lewis acid Yb(OTf)₃ (Scheme 16).²⁸

In addition to the utility of new C-nucleophiles for devising new MCRs assembling 1,4-DHPs, the employment of a new

electrophilic species to replace traditional aldehyde electrophiles has also been recently developed as practical tactic. And some well defined electrophilic species such as isatin and analogous 1,2-dicarbonyl compounds have been found particularly interesting in this regard by affording spirocyclic 1,4-DHPs derivatives through MCRs. For example, isatin **34** as well as analogous dicarbonyl substrate **36** have been both successfully utilized to react with cyclic 1,3-dicarbonyl compounds and aromatic primary amines respectively to construct polycyclic 1,4-DHPs **35**²⁹ and **37**³⁰ which containing both spiro- and fused architecture (Scheme 17). It should be noted that the essence of these reactions is identical as that of the Hantzsch reaction.




Scheme 14 Electron deficient alkene-based synthesis of 1,4-DHPs.



Scheme 16 β-Ester acetal initiated synthesis of 1,4-DHPs.

Reasonably, following the consideration on designing new MCRs for the synthesis of structurally diverse 1,4-DHPs employing aldehydes as C-electrophiles, new types of MCRs employing electrophilic isatins had also been developed for 1,4-DHP library synthesis. Through an analogous route described in Scheme 13, Perumal *et al.* performed MCRs for synthesis of spiro-1,4-DHPs **40** wherein isatin took the place of the aldehyde component in Scheme 13. With the catalysis of triethylamine, key Knoevenagel condensed intermediates **38** and enamino ethers **39** assembled to provide corresponding products in generally good yields (Scheme 18).³¹ On the other hand, Alizadeh *et al.* employed diamines **41** and 1,1-bis(methylthio)-2-nitroethylene **42** as precursors of enaminone intermediates **43**, which incorporated **38** to provide spiro- and fused 1,4-DHPs **44** under the promotion of piperidine (Scheme 18).³² After their synthesis of 1,4-DHPs **44**, Alizadeh *et al.* later developed another four-component reaction for 1,4-DHP synthesis. The protocol employed Wittig reagent **45** to react with isatin to form an electron deficient Wittig intermediate which was efficiently captured by the enamino ester intermediates *in situ* generated from ketoesters and amines to give 1,4-DHPs products **46** in good to excellent yields (Scheme 19).³³ On the other hand,




Scheme 17 Synthesis of spiro-1,4-DHPs with 1,2-dicarbonyl substrates.


Sambri and coworkers developed an unconventional multi-component protocol for the synthesis of 1,4-DHPs **47** from ethyl propiolate, two 1,3-dicarbonyl compounds and primary amines wherein ethyl propiolate served as electrophile (Scheme 20).³⁴

3. Multicomponent synthesis of 1,4-DHPs using ammonia as amino source


Ammonia was used as the amino source in the earliest version of the Hantzsch 1,4-DHPs synthesis, the reactivity of ammonia was identical as primary amines in most 1,4-DHPs synthesis, the difference was that 1,4-DHPs containing a free NH-group in the DHP ring were the products when ammonia was utilized as the amino source. Similar to the cases of amine-based reactions, novel MCRs have been known for the synthesis of new 1,4-DHPs libraries bearing a free NH group in the central ring by using ammonia as the amino source. For instance, starting from aldehyde, alcohols, diketene **48** and ammonia acetate, an alternative four-component protocol for synthesizing Hantzsch-type 1,4-DHPs **50** in the presence of 10 mol% H₂SO₄ had been



Scheme 19 Four-component synthesis of spiro-1,4-DHPs involving Wittig reagent and isatin.




Scheme 18 Multicomponent synthesis of spiro-1,4-DHPs using isatins as carbonyl electrophiles.




Scheme 20 Propiolate as electrophile in 1,4-DHPs synthesis.

achieved. The reactions proceeded *via* key intermediates **49** which were generated from 4-methyleneoxetan-2-one **48**, aldehydes and alcohols (Scheme 21).³⁵ In another example, the (2-aryl)methylene malononitriles **50** which were prepared from aldehydes and malonitrile have been successfully employed to react with ethyl acetoacetate and ammonia acetate to provide 1,4-DHPs **51** by grinding under solvent-free conditions (Scheme 22).³⁶ Safari *et al.* developed a new three-component synthesis of NH-free 1,4-DHPs of type **53** using chalcones **52**, ethyl acetoacetate and ammonia. Excellent yields of products were obtained in refluxing water with the catalysis of cellulose sulfuric acid (Scheme 23).³⁷


Ammonia acetate has also been used successfully as the reaction partner of cyclic 1,3-diketone **54**. Acyclic 1,3-dicarbonyl compounds and isatin for multicomponent synthesis of 1,4-DHPs, unsymmetrical sipro- and fused cyclic scaffolds **55** were provided in good to excellent yield by subjecting the four-component reactions to refluxing toluene in the presence of 15 mol% pyridine (Scheme 24).³⁸



Scheme 21 Synthesis of 1,4-DHPs involving 4-methyleneoxetan-2-one.



Scheme 22 Three-component synthesis of 1,4-DHPs using dienemalononitriles.




Scheme 23 Chalcone-based three-component 1,4-DHPs synthesis.


4. Multicomponent synthesis of 1,4-DHPs using enaminone, β -enamino esters or enamines as amino source

Enaminones, β -enamino esters and enamines are dinucleophilic species which possess broad reactivity through employing both the amino group and β -nucleophilic sp^2 carbon. These compounds have been proved particularly useful for the synthesis of various heterocyclic products. As introduced in the above contents, β -enamino esters were key intermediates in many amine-based MCRs synthesis of 1,4-DHPs, it is therefore not surprising that β -enamino esters as well as analogous substrates were also frequently employed as starting materials for 1,4-DHPs synthesis directly. Actually, during chemists' earliest efforts in devising new synthesis of structurally unsymmetrical 1,4-DHPs, β -enamino esters were selected as ideal candidates as starting building blocks. Usually, β -enamino esters are first prepared by subjecting an amine to incorporate a 1,3-dicarbonyl compound. The obtained β -enamino ester was then employed with an aldehyde and a different 1,3-dicarbonyl compound to provide unsymmetrical 1,4-DHPs in the manner of a three-component reaction. This kind of strategy had been applied for the synthesis of pyrazolyl functionalized 1,4-DHPs **57** by Perumal and co-workers. As a result, acyclic 1,3-dicarbonyl compounds, β -enamino ester **56** and pyrazolyl aldehydes were successfully assembled to provide **57** in ionic liquid [bmim]Cl in the presence of 3,4,5-trifluorobenzene boronic acid (Scheme 25).³⁹ Wang *et al.* conducted a systematic investigation on this kind of synthesis using cyclic diketone based β -enamino esters **58**, aromatic aldehydes and a different cyclic diketone. These reactions were found practical in refluxing water or solvent-free heating conditions to yield 1,4-DHPs **59** (Scheme 26).⁴⁰


An important feature of β -enamino ester/enaminone based MCRs was that they are not only useful in the synthesis of 1,4-DHPs with unsymmetrical structure, but also capable of serving as building blocks in the synthesis of symmetrical 1,4-DHPs. Dondoni *et al.* synthesized highly functionalized 1,4-DHPs **61**



Scheme 24 Ammonia-based synthesis of fused spiro-1,4-DHPs.




Scheme 25 Enaminone-based synthesis of pyrazolyl 1,4-DHPs.



Scheme 26 Enaminone-based synthesis of fused 1,4-DHPs.

using β -keto esters and β -enamino esters prepared from identical β -keto esters and ammonia to incorporate functionalized aldehyde **60**, the products were obtained as enantioenriched compounds owing to the use of functional chiral aldehyde as starting materials (Eq 1, Scheme 27).⁴¹ Later on, the same group developed an organocatalyzed three-component reaction of sugar aldehydes **62**, enaminones **63** and 1,3-dicarbonyl compounds in the presence of L-proline catalyst, 1,4-DHPs **64** were furnished in excellent diastereoselectivity with the inducement of chiral sugar aldehydes (Eq 2, Scheme 27).⁴² Mirza-Aghayan *et al.* developed a three-component synthesis of symmetrical 1,4-DHPs **65** by directly employing two molecules of identical β -enamino esters and aldehydes (Eq 3, Scheme 27).⁴³ In addition, the free NH group functionalized enaminones/ β -enamino esters of **66** have also been discovered as practical materials for the synthesis 1,4-DHPs. Li and coworkers successfully performed the assembly of 2 molecules of **66** and aldehydes to provide 2,6-unsubstituted 1,4-DHPs **67** with the catalysis of TsOH·H₂O or NaAuCl₄·2H₂O. Excellent tolerance had been demonstrated with this protocol as R² and R³ could both be alkyl or aryl groups when incorporating aromatic aldehydes (Eq 4, Scheme 27).⁴⁴


Interestingly, besides serving as dinucleophiles, β -enamino esters could also act as electrophilic acceptors for multicomponent 1,4-DHPs synthesis under proper conditions. Ajavakom *et al.* discovered that the Lewis acid TiCl₄ catalyzes the assembly of three molecules of β -enamino ester **68** to yield symmetrical 1,4-DHPs **72**. This three-component process was initiated by a



Scheme 27 Various multicomponent synthesis of symmetrical 1,4-DHPs.

Michael addition between 2 molecules of **66** affording intermediates **69**. Upon the elimination of one molecule of amine on **69**, intermediates **70** were generated as new Michael acceptors. The addition of one more **69** to **70** led to formation of intermediates **71**, and the intramolecular elimination cyclization of **71** provided products **72** (Scheme 28).⁴⁵

Owing to the excellent reactivity and diverse availability of enaminones, many diversity oriented, enaminone-based synthesis of 1,4-DHPs have been reported by using either new 1,3-dicarbonyl substrates or enaminones/ β -enamino esters. For example, enaminone **73** has been found as an ideal reaction partner of isatin and cyclic 1,3-diketones to yield spiro-fused 1,4-DHPs **74** in refluxing water with the catalysis of Brønsted acid *p*-TSA (Eq 1 in Scheme 29),⁴⁶ while enaminone **75** with the




Scheme 28 Assembly of 1,4-DHPs from three enaminones.

coumarin backbone has been reported as practical precursor for the synthesis of highly functionalized 1,4-DHPs **76** employing (\pm) lactic acid and ethyl lactate as catalyst and green solvent, respectively (Eq 2 in Scheme 29).⁴⁷ Recently, the easily available 6-amino-1,3-dimethyl uracil **77**, which was also formally an enaminone building block, has been demonstrated as a useful starting material in the three-component synthesis of 1,4-DHPs **78** by incorporating aromatic aldehydes and β -ketoesters in water (50 °C) in the presence of thiourea dioxide (Eq 3 in Scheme 29).⁴⁸


In the process of exploring new MCRs for heterocyclic scaffold synthesis, Li *et al.* established the synthesis of fused multicyclic 1,4-DHPs **81** from β -ketoester, aldehydes and ketene aminal compounds (cyclic enaminones) **79**. It was the 1,4-DHPs

80 that were first generated from a triethyl amine promoted three-component assembly, as intermediate products, **80** underwent efficient intramolecular coupling cyclization with K_2CO_3 catalysis to produce the final products. On the other hand, fused pyridine derivatives **83** were yielded when Meldrum's acid **82** was used as the dicarbonyl compound for the reaction under standard reaction conditions (Scheme 30).⁴⁹

Similar to the cases of amine-based multicomponent synthesis of 1,4-DHPs, substituted acetonitriles and enals were also frequently employed reaction partners in enaminone/ β -enamino ester-based multicomponent synthesis of 1,4-DHPs, particularly for the synthesis of unsymmetrical 1,4-DHPs. For example, starting from enaminones **79**, aldehydes and malonitrile, Li *et al.* attained 1,4-DHPs **84** in refluxing MeCN in the presence of



Scheme 29 Various synthesis of spiro- or fused 1,4-DHPs with different enaminones.




Scheme 30 1,4-DHPs-based polycycles synthesis *via* enaminone-based three-component reaction.


triethylamine, and further treatment with K_2CO_3 in DMF on **84** led to the production of **85** (Scheme 31).⁵⁰ On the other hand, interesting chemo-selective formation of 1,4-DHPs were discovered in the three-component reactions of enaminones **86**, aldehydes and β -cyanoketones, wherein the selective condensation of the carbonyl instead of cyano group in the β -cyanoketones allowed the production of 3-cyano functionalized 1,4-DHPs **87**,⁵¹ and a similar transformation was also applied for the synthesis of indolyl substituted 1,4-DHPs **88** by using corresponding indolyl functionalized cyanoketones (Scheme 31).⁵² More recently, malonitrile and ethyl 2-cyanoacetates were also successfully subjected to incorporate aldehydes and nitro-functionalized cyclic enaminones **89** for 1,4-DHP synthesis. Unlike the reaction process involved in the MCRs using β -cyanoketones, 2-amino functionalized fused 1,4-DHPs **90** were afforded in the reactions *via* a nucleophilic addition transformation on the cyano group of the β -cyanoester (Scheme 31).⁵³ In the area of multicomponent synthesis of 1,4-DHPs employing enaminone as amino source, Pan and coworkers developed the three-component regioselective synthesis of 1,4-DHPs of type **92** and isomeric 1,2-DHPs **93** using N,N' -dimethyl amino functionalized enaminones **91**, primary amines and enals. The designed systematic experiments using primary amines of different

electronic and steric properties suggested that a strong electron withdrawing group or bulky *ortho*-group tend to induce the formation of 1,2-DHPs **93**, combined with the observation of compounds **94** in some entries, it was proposed that **94** were the key intermediates of the reaction. The normal transformation of the N-nucleophilic condensation with the formyl group in the enal component led to the production of 1,4-DHPs **92**, when a strong electron withdrawing or bulky *ortho*-group was included in the amine component, the nucleophilicity of the amino group in **94** was significantly hindered, which allowed the nucleophilic carbon atom in **94** to attack the enal formyl first and led to the regioselective formation of 1,2-DHPs **93**. This work was the first example on the tunable selective synthesis of 1,4-DHPs and 1,2-DHPs with three-component reactions (Scheme 32).⁵⁴

Although not frequently available, simple enamines have also been reported to be capable of serving as dinucleophilic building blocks for multicomponent synthesis of 1,4-DHPs by acting similarly as enaminones and β -enamino esters. Quiroga *et al.* reported the synthesis of elaborated polycyclic products **96** with heteroaryl amines **95**, isatin and various 1,3-dicarbonyl substrates. Although **96** are formally fused aromatic heterocycles, the central 1,4-DHP backbone was formed during the transformation *via* cross-condensed intermediates **97** (Scheme 33).⁵⁵




Scheme 31 Various synthesis of 1,4-DHPs-based heterocycle synthesis with different enaminones.




Scheme 32 Enaminone initiated regioselective synthesis of 1,4- and 1,2-DHPs.

Rather recently, a similar three-component transformation using enamine **98**, cyclic diketones and aromatic aldehydes have been reported to provide multicyclic elaborated 1,4-DHPs **99** (Scheme 34).⁵⁶



Scheme 33 Cyclic enamine-based three-component synthesis of highly functionalized 1,4-DHPs.




Scheme 34 Multicyclic 1,4-DHPs synthesis using cyclic enamines.

5. Multicomponent synthesis of 1,4-DHPs using other amino sources

According to related results in the known literature, amines, ammonia and enaminone analogs were obviously the most frequently and conventional amino sources used for the synthesis of 1,4-DHPs, however, some other types of N-containing species such as cyanides, imines *etc.* were also able to serve as the amino sources for the synthesis of novel 1,4-DHPs *via* MCRs as occasional cases. Depending on the properties of these amino resources, both 1,4-DHPs with novel elaboration and 1,4-DHPs with similar structure as those equivalents obtained from conventional amino resources were possible.

5.1. Using cyanide as amino source

As reactive starting materials, cyanide derivatives such as malonitrile were frequently utilized in the synthesis of 1,4-DHPs in amine, ammonia or enaminone participated MCRs. In these kinds of reactions, as introduced above, it was amine, enaminone or ammonia that served as the amino source to construct the heterocyclic backbone while cyano groups were mainly involved as the donor of electrophilic carbons in those MCRs. Interestingly, in some reactions, it was also possible to directly employ the N atom in the cyano group as the amino source of 1,4-DHPs. Fustero and coworkers developed the three-component protocol involving fluorinated nitriles (cyanides) **100**, alkyl propiolates and chiral allyl *p*-tolyl sulfoxide **101** for the synthesis of 1,4-DHPs **104**. The reactions proceeded under the promotion of $\text{KN}(\text{SiMe}_3)_2$ *via* two steps of cascade transformations, the first nucleophilic addition of **101** to **100** under catalysis of base gave intermediates **102**, the second step addition of **102** to propiolates led to the formation of intermediates **103**, and intramolecular Michael addition on **103** provided enantioenriched 1,4-DHPs **104** in generally excellent diastereoselectivity, which was induced by the chiral *p*-tolyl sulfoxide (Scheme 35).⁵⁷




Scheme 35 Diastereoselective synthesis of 1,4-DHPs using cyanide amino source.

5.2. Using imine as amino source

Imines were usually the resulted products from the condensation of amines and aldehydes. Direct employment of aldehydes, amines and propiolates have been a well established protocol for the synthesis of 1,4-DHPs. Interestingly, Fukuzawa *et al.* developed a three-component reaction of imines **105** and two molecules of propiolates to provide 1,4-DHPs of type **9** (see also Scheme 7) in the presence of 10 mol% Lewis acid catalyst $\text{Sc}(\text{OTf})_3$. In these reactions, one key step was the decomposition of imines **105** to aldehydes and amines. Only low to moderate yields of products were obtained by this methodology with imines prepared from the condensation of aromatic aldehydes with aromatic/aliphatic amines (Scheme 36).⁵⁸

5.3. Using thioamide as amino source

In devising the MCRs synthesis of structurally diverse 1,4-DHPs employing new C-nucleophilic building blocks, β -arylthioamides **106** have been found as useful starting materials in the synthesis of 1,4-DHPs **108** containing a thioester functional group. During the reaction process, **106** tautomerized as **107** to incorporate aldehydes in the presence of electron withdrawing group




Scheme 36 Imine in three-component synthesis of 1,4-DHPs.


functionalized acetonitriles and alkyl halide. Microwave irradiation had been found to give superior results over traditional heating conditions to this Et_3N -catalyzed 1,4-DHPs synthetic method (Scheme 37).⁵⁹ On the basis of this transformation, a further devised cascade three-component protocol for the synthesis of fused heterocyclic 1,4-DHPs **112** was reported by employing 2-chlorophenyl functionalized thioamides **109**, aldehydes and malonitrile. In the presence of KF /neutral Al_2O_3 catalysts with PEG6000 and microwave irradiation, **112** were efficiently provided through the intramolecular cyclization of intermediates **110/111** (Scheme 38).⁶⁰

6. Conclusions

As a class of typical heterocyclic scaffolds with broad spectrum of biological and pharmaceutical profiles, 1,4-DHPs have been regarded as significant targets of organic synthesis. Following the daily increasing requirement on molecular libraries during the discovering process of new lead compounds and drugs, urgent promotion was simultaneously brought to the chemistry of 1,4-DHPs synthesis. Particularly, the synthesis of 1,4-DHPs possessing novel and diverse elaboration consists of the main effort in this field. With those various types of newly designed MCRs, it is now possible to access highly functionalized 1,4-DHPs with enormously expanded diversity on the basis of the classical Hantzsch multicomponent synthesis. The breakthrough in the field should be ascribed to those highly diversified and reactive starting materials including amine resources, various C-electrophiles such as aldehydes and isatins, bifunctional building blocks such as enals, enones, 1,3-dicarbonyl compounds



Scheme 37 Three-component synthesis of thioester functionalized 1,4-DHPs.



Scheme 38 Cascade three-component synthesis of sulphur-containing heterocycle fused 1,4-DHPs.

as well as different alkynes/alkenes *etc.* In this regard, devising more new MCRs for the synthesis of structurally novel 1,4-DHPs relies on further endeavours in exploring more unconventional building blocks.

In addition, in terms of sustainable synthesis, to develop 1,4-DHPs synthesis with MCRs under milder, greener conditions is also highly desirable in this field. More importantly, as most new MCRs for 1,4-DHPs assembly provide products containing chiral centers, methods of directly accessing enantiomerically pure 1,4-DHPs with MCRs is in urgent requirement. Establishment of efficient methodologies providing chiral 1,4-DHPs is highly expectable to significantly enhance the progress of discovering more medicinally valuable heterocyclic compounds.

Acknowledgements

The authors gratefully acknowledge the financial support from the NSFC of China (21102059), the NSF of Jiangxi Province (20114BAB213005) and a Sponsored Program for Cultivating Youths of Outstanding Ability from Jiangxi Normal University.

References

- (a) S. L. Schreiber, *Science*, 2000, **287**, 1964–1969; (b) D. S. Tan, *Nat. Chem. Biol.*, 2005, **1**, 74–84; (c) T. E. Nielsen and S. L. Schreiber, *Angew. Chem., Int. Ed.*, 2008, **47**, 48–56; (d) S. Shang and D. S. Tan, *Curr. Opin. Chem. Biol.*, 2005, **9**, 248–258; (e) G. L. Thomas, E. E. Wyatt and D. R. Spring, *Curr. Opin. Drug Discov. Dev.*, 2006, **9**, 700–712.
- (a) A. Dömling, *Chem. Rev.*, 2006, **106**, 17; (b) A. Dömling and I. Ugi, *Angew. Chem., Int. Ed.*, 2000, **39**, 3168; (c) V. Nair, C. Rajesh, A. U. Vinod, S. Bindu, A. R. Sreekanth, J. S. Mathen and L. Balagopal, *Acc. Chem. Res.*, 2003, **36**, 899–907; (d) J.-P. Wan and Y. Liu, *Curr. Org. Chem.*, 2011, **15**, 2758–2773.
- A. Hantzsch, *Justus Liebigs Ann. Chem.*, 1882, **215**, 1–82.
- K. K. Borowicz, M. Gasior, Z. Kleinrok and S. J. Czuczwar, *Eur. J. Pharmacol.*, 1997, **323**, 45–51.
- A. H. Li, S. Moro, N. Forsyth, N. Melman, X.-D. Ji and K. A. Jacobson, *J. Med. Chem.*, 1999, **42**, 706–721.
- I. O. Donkor, X. Zhou, J. Schmidt, K. C. Agrawal and V. Kishore, *Bioorg. Med. Chem.*, 1998, **6**, 563–568.
- A. Mai, S. Valente, S. Meade, V. Carafa, M. Tardugno, A. Nebbioso, A. Galmozzi, N. Mitro, E. D. Fabiani, L. Altucci and A. Kazantsev, *J. Med. Chem.*, 2009, **52**, 5496–5504.
- (a) For known reviews on 1,4-dihydropyridine chemistry, see: U. Eisner and J. Kuthan, *Chem. Rev.*, 1972, **72**, 1–42; (b) D. M. Stout and A. I. Meyers, *Chem. Rev.*, 1982, **82**, 223–243; (c) A. Sausins and G. Duburs, *Heterocycles*, 1988, **27**, 269–289.
- (a) For some recent reviews wherein the Hantzsch synthesis of 1,4-DHPs was mentioned, see: C. de Graaff, E. Ruijter and R. V. A. Orru, *Chem. Soc. Rev.*, 2012, **41**, 3969–4009; (b) A. Dömling, W. Wang and K. Wang, *Chem. Rev.*, 2012, **112**, 3083–3135; (c) A. Moyano and R. Rios, *Chem. Rev.*, 2011, **111**, 4703–4832; (d) A. Dondoni and A. Massi, *Acc. Chem. Res.*, 2006, **39**, 451–463; (e) N. Isambert, M. M. Sanchez Duque, J.-C. Plaquevent, Y. Génisson, J. Rodriguez and T. Constantieux, *Chem. Soc. Rev.*, 2011, **40**, 1347–1357; (f) D. Bonne, Y. Coquerel, T. Constantieux and J. Rodriguez, *Tetrahedron: Asymmetry*, 2010, **21**, 1085–1109; (g) J. Yu, F. Shi and L.-Z. Gong, *Acc. Chem. Res.*, 2011, **44**, 1156–1171; (h) H. Pellissier, *Adv. Synth. Catal.*, 2012, **354**, 237–294.
- J.-P. Wan and Y. Liu, *Synthesis*, 2010, 3943–3953.
- R. K. Vohra, C. Bruneau and J.-L. Renaud, *Adv. Synth. Catal.*, 2006, **348**, 2571–2574.
- V. Sridharan, P. T. Perumal, C. Avendaño and J. C. Menéndez, *Tetrahedron*, 2007, **63**, 4407–4413.
- A. Kumar and R. A. Maurya, *Tetrahedron*, 2008, **64**, 3477–3482.
- M. L. Kantam, T. Tamani, L. Chakrapani and B. M. Choudary, *Catal. Commun.*, 2009, **10**, 370–372.
- D. Ramesh, S. Rajaram, M. Narasimhulu, T. S. Reddy, K. C. Mahesh, G. Manasa and Y. Venkateswarlu, *Chin. J. Chem.*, 2011, **29**, 2471–2475.
- S. Maiti, V. Sridharan and J. C. Menéndez, *J. Comb. Chem.*, 2010, **12**, 713–722.
- J. Jiang, J. Yu, X.-X. Sun, Q.-Q. Rao and L.-Z. Gong, *Angew. Chem., Int. Ed.*, 2008, **47**, 2458–2462.
- C. G. Evans and J. E. Gestwicki, *Org. Lett.*, 2009, **11**, 2957–2959.
- S.-L. Cui, J. Wang, X.-F. Lin and Y.-G. Wang, *J. Org. Chem.*, 2007, **72**, 7779–7782.
- J. Sun, Y. Sun, E.-Y. Xia and C.-G. Yan, *ACS Comb. Sci.*, 2011, **13**, 436–441.
- H. Jiang, R. Mai, H. Cao, Q. Zhu and X. Liu, *Org. Biomol. Chem.*, 2009, **7**, 4943–4953.
- J. Sun, Y. Sun, H. Gao and C.-G. Yan, *Eur. J. Org. Chem.*, 2011, 6952–6956.
- A. T. Khan and M. M. Khan, *Tetrahedron Lett.*, 2011, **52**, 3455–3459.
- J. Sun, Q. Wu, E.-Y. Xia and C.-G. Yan, *Eur. J. Org. Chem.*, 2011, 2981–2986.
- A. Kumar and S. Sharma, *Green Chem.*, 2011, **13**, 2017–2020.
- H. Jiang, X. Ji, Y. Li, Z. Chen and A. Wang, *Org. Biomol. Chem.*, 2011, **9**, 5358–5361.
- P. R. Girling, A. S. Batsanov, H. C. Shen and A. Whiting, *Chem. Commun.*, 2012, **48**, 4893–4895.
- S. Sueki, R. Takei, J. Abe and I. Shimizu, *Tetrahedron Lett.*, 2011, **52**, 4473–4477.
- R. Ghahremanzadeh, G. I. Shakibaei, S. Ahadi and A. Bazgir, *J. Comb. Chem.*, 2010, **12**, 191–194.
- R. Ghahremanzadeh, S. Ahadi, G. I. Shakibaei and A. Bazgir, *Tetrahedron Lett.*, 2010, **51**, 499–502.
- S. E. Kiruthika, N. V. Lakshmi, B. R. Banu and P. T. Perumal, *Tetrahedron Lett.*, 2011, **52**, 6508–6511.
- A. Alizadeh, T. Firuziyar and A. Mikaeili, *Synthesis*, 2010, 3913–3917.
- A. Alizadeh and J. Mokhtari, *Tetrahedron*, 2011, **67**, 3519–3523.
- G. Bartoli, M. Bosco, P. Galzerano, R. Giri, A. Mazzanti, P. Melchiorre and L. Sambri, *Eur. J. Org. Chem.*, 2008, 3970–3975.
- A. Alizadeh and S. Rostamnia, *Synthesis*, 2010, 4057–4060.
- A. M. Zonouz and D. Moghani, *Synth. Commun.*, 2011, **41**, 2152–2160.
- J. Safari, S. H. Banitaba and S. D. Khalili, *J. Mol. Catal. A: Chem.*, 2011, **335**, 46–50.
- A. Feiz, G. I. Shakibaei, Z. Yasaei, H. R. Khavasi and A. Bazgir, *Helv. Chim. Acta*, 2011, **94**, 1628–1637.
- R. Sridhar and P. T. Perumal, *Tetrahedron*, 2005, **61**, 2465–2470.
- G.-W. Wang and C.-B. Miao, *Green Chem.*, 2006, **8**, 1080–1085.
- A. Dondoni, A. Massi, E. Minghini and V. Bertolasi, *Tetrahedron*, 2004, **60**, 2311–2326.
- D. R. B. Ducatti, A. Massi, M. D. Nosedá, M. E. R. Duarte and A. Dondoni, *Org. Biomol. Chem.*, 2009, **7**, 1980–1986.
- M. Mirza-Aghayan, M. K. Langrodi, M. Rahimifard and R. Boukherroub, *Appl. Organomet. Chem.*, 2009, **23**, 267–271.
- J. Yang, C. Wang, X. Xie, H. Li and Y. Li, *Eur. J. Org. Chem.*, 2010, 4189–4193.
- T. Sirijindalert, K. Hansuthirakul, P. Rashatasakhon, M. Sukwattanasinitt and A. Ajavakom, *Tetrahedron*, 2010, **66**, 5161–5167.
- K. Jadidi, R. Ghahremanzadeh, P. Mirzaei and A. Bazgir, *J. Heterocycl. Chem.*, 2011, **48**, 1014–1018.
- S. Paul and A. R. Das, *Tetrahedron Lett.*, 2012, **53**, 2206–2210.
- S. Verma and S. L. Jain, *Tetrahedron Lett.*, 2012, **53**, 2595–2600.
- L.-R. Wen, C. Liu, M. Li and L.-J. Wang, *J. Org. Chem.*, 2010, **75**, 7605–7614.
- L.-R. Wen, C.-Y. Jiang, M. Li and L.-J. Wang, *Tetrahedron*, 2011, **67**, 293–302.
- J.-X. Wang, G.-L. Shen and D.-Q. Shi, *J. Heterocycl. Chem.*, 2011, **48**, 1145–1148.
- T. Chen, X.-P. Xu, H.-F. Liu and S.-J. Ji, *Tetrahedron*, 2011, **67**, 5469–5476.
- C. Altug, A. K. Burnett, E. Caner, Y. Dürüst, M. C. Elliott, R. P. J. Glanville, C. Guy and A. D. Westwell, *Tetrahedron*, 2011, **67**, 9522–9528.
- J.-P. Wan, S.-F. Gan, G.-L. Sun and Y.-J. Pan, *J. Org. Chem.*, 2009, **74**, 2862–2865.

-
- 55 J. Quiroga, S. Portillo, A. Péres, J. Gálves, R. Abonia and B. Insuasty, *Tetrahedron Lett.*, 2011, **52**, 2664–2666.
- 56 A. T. Khan and D. K. Das, *Tetrahedron Lett.*, 2012, **53**, 2345–2351.
- 57 S. Fustero, S. Catalán, M. Sánchez-Roselló, A. Simón-Fuentes and C. del Pozo, *Org. Lett.*, 2010, **12**, 3484–3487.
- 58 S. Kikuchi, M. Iwai, H. Murayama and S.-I. Fukuzawa, *Tetrahedron Lett.*, 2008, **49**, 114–116.
- 59 M. Li, Z. Zuo, L. Wen and S. Wang, *J. Comb. Chem.*, 2008, **10**, 436–441.
- 60 L. Wen, C. Ji, Y. Li and M. Li, *J. Comb. Chem.*, 2009, **11**, 799–805.