

A New Approach to the Stereospecific Total Synthesis of Racemic *Cecropia* Juvenile Hormone†

By KIYOSI KONDO,* AKIRA NEGISHI, KIYOHIDE MATSUI, and DAIEI TUNEMOTO
(Sagami Chemical Research Center, 3100 Onuma, Sagamihara-shi 229, Japan)

and SATORU MASAMUNE
(Department of Chemistry, University of Alberta, Edmonton, Alberta, Canada)

Summary 4-Methyl-5,6-dihydro-2*H*-thiopyran and its dimer (**4**) are used as structural units for the synthesis of C₁₈-juvenile hormone.

WE describe herein a new and economically feasible method for the stereospecific synthesis¹ of a racemic *Cecropia* juvenile hormone (C₁₈-J.H.), based on condensation of dihydrothiopyrans. The structure of C₁₈-J.H. can be divided into three structural units (A)—(C). The configuration of two ethyl groups in units (A) and (B) might be retained if they were blocked with two sulphur atoms [see


(A') and (B')]. These blocking atoms could easily be removed by reductive desulphurization.

A key intermediate, the thiopyran (**4**), was synthesised by two methods. Condensation of tetrahydrothiopyran-4-one² with dimethyloxosulphonium methylide afforded the epoxide (**2**)‡ (65%), m.p. 52°. The carbanion prepared from the thiopyran (**1**)³ and BuⁿLi in the presence of 2,3-diazabicyclo[2,2,2]octane (DABCO) was treated with the epoxide (**2**) in THF at -20° and the resulting adduct was further dehydrated with SOCl₂-pyridine to give the desired dimeric dihydrothiopyran (**4**) [73% based on (**2**)],

† Presented at the 26th Annual Meeting of Japan Chemical Society, April 1st, 1972, Hiratsuka, Japan.

‡ All new compounds gave satisfactory elemental analyses and i.r. and n.m.r. spectra consistent with the assigned structures.


b.p. 85° at 0.2 mmHg. The other method was based on the condensation of (1) with (3), which was prepared from tetrahydrothiopyran-4-one by treatment with HCN, followed by solvolysis with EtOH, dehydration, reduction, and bromination with PBr₃.



Attachment of the final unit (C) and the removal of the blocking sulphur atoms were carried out by the following routes.

Route A: The dimer (4) was converted into a carbanion by treatment with BuⁿLi-DABCO and was then treated with propylene oxide to afford the alcohol (5) (84%). Reductive desulphurization of (5) by treatment with metallic Li in ethylamine⁴ at -20° produced the dodecadienol (6) (60%), b.p. 97–98° at 0.45 mmHg. Oxidation of the alcohol (6) with Jones reagent yielded the ketone (7) (85%), b.p. 98–99° at 1.5 mmHg, which was further transformed into deoxy C₁₈-J.H. by condensation with diethyl methoxycarbonylmethylphosphonate.⁵ Epoxidation⁶ of the ester gave racemic J.H., which was identical spectroscopically with natural J.H.⁷

Route B: The carbanion of dimer (4) was condensed with *trans*-4-chloro-3-methylbut-2-enyl tetrahydropyranyl ether, prepared from isoprene,⁸ to yield the triene (8; R = THP) (60%). The trienol (8; R = H) was treated with Li in ethylamine at -70° and the resulting dithio-alcohol (9; R = H) was converted into the corresponding acetate (9; R = Ac) (80%). Final desulphurization of (9; R = Ac) was achieved by treatment with excess of deactivated Raney nickel (W-2/acetone) to give the acetate (10) as a colourless oil (55%), b.p. 115° at 0.15 mmHg. Deoxy-C₁₈-J.H. obtained from (10) by the method of Corey *et al.*⁹ was identical spectroscopically with that prepared from the ketone (7).



C₁₈-J.H. can thus be obtained with 100% stereospecificity at C-6 and C-10 and 95% or more at C-2¹⁰ by route A, and with complete stereospecificity by route B.

This investigation was supported by Kawakami Memorial Foundation. S.M. thanks the Defence Research Board of Canada and Hoffmann-La Roche, Inc. for financial support.

(Received, 30th August 1972; Com. 1514.)

¹ E. J. Corey, J. A. Katzenellenbogen, N. W. Gilman, S. A. Roman, and B. W. Erickson, *J. Amer. Chem. Soc.*, 1966, **90**, 5618; R. Zurflüh, E. N. Wall, J. B. Siddall, and J. A. Edwards, *ibid.*, 6224; W. S. Johnson, T. Li, D. J. Faulkner, and S. F. Campbell, *ibid.*, p. 6225; E. J. Corey and H. Yamamoto, *ibid.*, 1970, **92**, 6636; E. E. van Tamelen, P. McCurry, and U. Huber, *Proc. Nat. Acad. Sci. U.S.A.*, 1971, **68**, 1294.

² E. A. Fehnel and M. Carmack, *J. Amer. Chem. Soc.*, 1948, **70**, 1813.

³ R. F. Naylor, *J. Chem. Soc.*, 1949, 2749.

⁴ A. S. Hallsworth, H. B. Henbest, and T. I. Wrigley, *J. Chem. Soc.*, 1957, 1969.

⁵ K. H. Dahm, B. M. Trost, and H. Röller, *J. Amer. Chem. Soc.*, 1967, **89**, 5292.

⁶ E. E. van Tamelen and T. J. Curphey, *Tetrahedron Letters*, 1962, 121.

⁷ B. M. Trost, *Accounts Chem. Res.*, 1970, **3**, 120.

⁸ W. Oroshnik and R. A. Mallory, *J. Amer. Chem. Soc.*, 1950, **72**, 4608.

⁹ E. J. Corey, N. W. Gilman, and B. E. Ganem, *J. Amer. Chem. Soc.*, 1968, **90**, 5616.

¹⁰ Y. Butsugan, M. Murai, and T. Matsuura, *Nippon Kagaku Zasshi*, 1969, **89**, 431.