

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/233893086>

Anion-driven self-assembly: From discrete cages to infinite polycatenanes step by step

ARTICLE in CHEMICAL COMMUNICATIONS · DECEMBER 2012

Impact Factor: 6.83 · DOI: 10.1039/c2cc36921d · Source: PubMed

CITATIONS

14

READS

20

6 AUTHORS, INCLUDING:



Feilong Jiang Prof

Chinese Academy of Sciences

184 PUBLICATIONS 2,742 CITATIONS

SEE PROFILE



Daqiang Yuan

Chinese Academy of Sciences

279 PUBLICATIONS 10,113 CITATIONS

SEE PROFILE



Guangxun Lyu

Chinese Academy of Sciences

5 PUBLICATIONS 30 CITATIONS

SEE PROFILE



Maochun Hong

Chinese Academy of Sciences

402 PUBLICATIONS 9,661 CITATIONS

SEE PROFILE

Anion-driven self-assembly: from discrete cages to infinite polycatenanes step by step†

Cite this: *Chem. Commun.*, 2013, **49**, 719Received 24th September 2012,
Accepted 27th November 2012

DOI: 10.1039/c2cc36921d

www.rsc.org/chemcomm

Qihui Chen, Feilong Jiang, Daqiang Yuan, Lian Chen, Guangxun Lyu and
Maochun Hong*


Induced by different anions, two different polycatenanes constructed from Ag_6L_4 units with different conformations have been prepared. Intermediates Ag_6L_4 and Ag_5L_4 are observed to participate in interlocking, and the related possible self-assembly routes are also presented.

Nature has the ability to assemble simple molecule into extremely complex biomolecules by using a variety of weak interactions such as hydrogen bonding and charge–charge interactions.¹ Inspired by this deliberate approach, many elegant assemblies driven by coordination have been prepared.^{2–4} Catenanes and rotaxanes have attracted considerable attention not only for their interesting topological structures, but also for their astonishing applications such as in molecular machines and motors.⁵ In 1999, the Fujita group firstly synthesized a [2]catenane based on discrete three-dimensional (3D) interlocking metallacages.⁶ Since then, there have been several examples of [2]catenanes constructed from discrete 3D metallacages,^{6–9} however, an infinite polycatenane assembled from discrete metallacages seems more intriguing and challenging. The first infinite polycatenane assembled from discrete cages was reported by the Lu group in 2010.¹⁰ Recently, another type of polycatenane was prepared by Dehnen and coworkers.¹¹ To the best of our knowledge, only two infinite polycatenanes constructed from discrete metallacages have been reported, and the assembly process seems still obscure.^{10,11} Thus, construction of such highly organized architectures from discrete metallacages and an understanding of these processes have therefore remained a challenge for chemists. Herein, we report an interesting self-assembly system: a new polycatenane is constructed from Ag_6L_4 cages through introducing ClO_4^- anions. If larger PF_6^- anions are introduced, a different polycatenane with larger Ag_{12}L_8 cages composed of two Ag_6L_4 units is produced. The key intermediate which participates in interlocking directly is observed firstly, and the related possible self-assembly routes are also presented.

In order to achieve a metallacage with a suitable window for interlocking, a novel tripodal ligand **L** (**L** = tris(isonicotinoxy-methyl)ethane) was designed and synthesized. The ligand **L** is so flexible that it can adopt various conformations to satisfy the construction of a metallacage. For example, a M_6L_4 cage can be constructed from four molecular **L** and six molecular 2-coordinated metal ions. (Scheme 1)

Layering a CHCl_3 solution of ligand **L** with a MeOH solution of AgClO_4 provided colorless rectangular crystals of compound **1**. Single crystal X-ray diffraction analysis indicated that compound **1**, $[\text{Ag}_6\text{L}_4(\text{ClO}_4)_6 \cdot \text{CHCl}_3]_n$, has a polycatenated chain consisting of interlocking discrete metallacages. As shown in Fig. 1, the cage has a similar hypothetical structure to that illustrated in Scheme 1, which is comprised of six $\text{Ag}(\text{i})$ ions and four **L** ligands. The angles between the three coordinated pyridines in **L** are about 30, 81 and 102° , respectively. The Ag_6L_4 cage has a $(20 \times 16 \times 4 \text{ \AA})$ cavity with the ClO_4^- ions near the 2-coordinated $\text{Ag}(\text{i})$ to stabilize it.

As shown in Fig. 2, the polycatenane in **1** is constructed by the interlocking of neighboring Ag_6L_4 cages. The neighboring pyridine groups of **L** ligands twist in the same direction with the distance of *ca.* 4 Å between them, which indicates that the



Scheme 1 Ligand **L** and a possible cage constructed from *anti* conformations of **L** with **M** (**M** = 2-coordinated metal ions, such as Ag).




Fig. 1 The Ag_6L_4 cage in compound **1** (H atoms are omitted for clarity, grey: C; red: O; blue: N; green: Cl; teal: Ag).

State Key Laboratory of Structure Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou, 350002, China.
E-mail: hmc@fjirm.ac.cn; Fax: +86 591-83794946; Tel: +86 591-83712460

† Electronic supplementary information (ESI) available: Experimental section and figures. CCDC 888500 and 888501. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c2cc36921d




Fig. 2 Left, structural fragment containing three interlaced metallacages; right, the interlocking knot is further stabilized by strong charge-charge interactions.

interlocking of the Ag_6L_4 cages is driven by $\pi \cdots \pi$ interactions. The interlocking knot is further stabilized by the strong charge-charge interactions between two ClO_4^- anions and two Ag^+ ions, in which the distances between Ag and O atoms are 2.797 Å.

The formation of infinite architectures from discrete cages is very difficult and sensitive;^{10,11} even if using similar anions such as I^- and Br^- anions to replace Cl^- anions, the results are entirely different.¹¹ The 2-coordinated $\text{Ag}(\text{I})$ in compound **1** is unsaturated, and we wanted to know whether the Ag_6L_4 cage could be extended into a larger unit induced by a larger counter anion.

Employing the PF_6^- anion as the larger counter anion led to a new compound **2**.¹² Single crystal X-ray diffraction analysis indicated that compound **2**, $[\text{Ag}_{12}\text{L}_8(\text{PF}_6)_{12} \cdot x\text{CHCl}_3]_n$, is another new polycatenane, however, its basic units here for interlocking are Ag_{12}L_8 cages, composed of two Ag_6L_4 units with different conformation from compound **1** (the angles between the three coordinated pyridines are about 34, 101 and 111°, respectively) stabilized by the larger PF_6^- anions. The cavity of the Ag_{12}L_8 cage is up to $(22 \times 17 \times 10 \text{ Å})$ (Fig. 3).

The interlocking knot is stabilized only by the strong charge-charge interactions from one PF_6^- anion and four $\text{Ag}(\text{I})$ ions, with a distance of 2.872 Å between the Ag and F atoms. Driven by these interactions, the Ag_{12}L_8 cages interlock with each other to generate an elegant polycatenane (Fig. 4).

Induced by different anions, two different polycatenanes constructed from Ag_6L_4 units with different conformations have been prepared. The two novel structures represent the third example of the extension of discrete metallacages into an infinite polycatenane.^{10,11}

There are several examples in which discrete cages are extended into complicated topological motifs,^{6–11} however, only three examples demonstrate that the discrete cages can assemble into an infinite polycatenane rather than a less complex [2]catenane.^{10,11} There is a question as to why some discrete cages can generate infinite frameworks instead of limited [2]catenanes.^{6–9} Herein, we give our opinion on what induces discrete cages into infinite polycatenated frameworks.




Fig. 3 The Ag_{12}L_8 cage in **2**; which is comprised of two Ag_6L_4 units linked by $\text{Ag} \cdots \text{Ag}$ interactions (H atoms are omitted for clarity, grey: C; red: O; blue: N; green: F; teal: Ag; pink: P).





Fig. 4 Left, two interlocked metallacages stabilized by the strong charge-charge interactions from one PF_6^- anion and four Ag^+ ions; right, structural fragment containing three interlaced metallacages.

Firstly, a suitable window and large cavity; to the best of our knowledge, all of the discrete cages that can generate complicated motifs possess suitable windows and large cavities.^{6–11} Secondly, the interlocking is an entropically unfavourable process and needs suitable interactions to stabilize it. In Fujita, Clever, Kuroda and Hardie's examples,^{6–9} the cavities in newly-formed [2]catenanes are not large enough for the next interlock. In our case, the large windows and cavities from the cages sustain interlocking and, even after interlocking twice, the cavities in the polycatenanes still seem large (Fig. 2 and 4). Moreover, strong charge-charge interactions between counter anions and positively charged cages make the interlocking stable. There are also discrete cages with large windows and cavities and suitable interactions in Lu and Dehnen's systems.^{10,11}

The reason why discrete cages can further extend into infinite polycatenated architectures rather than limited [2]catenanes seems clear; nevertheless, the concrete process is still inexplicit.^{6–11} There may be two possible paths for the formation of [2]catenanes: path I, two open precursors are firstly locked to give an interlocking knot and closing this intermediate finally leads to the [2]catenane;¹³ path II, the closed unit is firstly formed, the unit has reversible 'on/off' ability, and the interlocking of two units directly leads to the [2]catenane (Scheme 2).

After many attempts, we found that the Ag_6L_4 unit could be assembled quickly in $\text{MeOH}-\text{CHCl}_3$ solution induced by ClO_4^- anions (see ESI†). More importantly, an unexpected spectral peak (2677) corresponding to Ag_5L_4 is firstly observed, which results from the loss of one $\text{Ag}(\text{I})$ ion from an Ag_6L_4 unit. The peak intensities of two different units (Ag_6L_4 and Ag_5L_4) are comparable, indicating a fast dynamic equilibrium between the Ag_6L_4 unit and Ag_5L_4 unit¹⁴ (Fig. 5). These observed intermediates demonstrate that the assembly process of compound **1** should follow path II rather than path I.

A possible assembly process of compound **1** is presented in detail (Scheme 3): the ligand **L** and $\text{Ag}(\text{I})$ quickly assemble into a Ag_6L_4 cage induced by ClO_4^- ions in CHCl_3 - MeOH solution (within 5 min); the Ag_6L_4 cage can open and lose a $\text{Ag}(\text{I})$ ion to form Ag_5L_4 and the process is reversible and fast; interlocking of one Ag_6L_4 cage with the newly formed Ag_5L_4 unit leads to a [2]catenane; the newly-formed [2]catenane continues to interlock, finally resulting in



Scheme 2 The two possible paths for the formation of [2]catenanes: (1) locking, (2) closing, (3) interlocking directly.





Fig. 5 ESI-MS spectrum of Ag_6L_4 cage and Ag_5L_4 unit generated *in situ*, in which the 1391 peak corresponds to $\text{Ag}_6\text{L}_4(\text{ClO}_4)_4^{2+}$, the 2885 peak to $\text{Ag}_6\text{L}_4(\text{ClO}_4)_5^+$, and the 2677 peak to $\text{Ag}_5\text{L}_4(\text{ClO}_4)_4^+$.



Scheme 3 Possible process of formation of compound **1**, in which the reactions (1), (2), (3), (4), (5) are fast, and the reaction (6) is slow.

compound **1**. The interlocking process is slow and can self-correct (only the interlocking knots with lowest energy are stable), even if the basic cage unit can be assembled within 5 min, the infinite interlocking products are obtained after two weeks. Induced by larger PF_6^- ions, Ag_6L_4 units rather than Ag_{12}L_8 are assembled immediately (see ESI†). Thus, the formation of compound **2** is presented as follows: stabilized by the larger counter anions, the *in situ* formed Ag_6L_4 units can spontaneously dimerize into larger Ag_{12}L_8 cages, and finally lead to compound **2** after infinite interlocking and self-correcting.

As we know, Ag_5L_4 is a key unit for interlocking in our system, while no such similar intermediates for interlocking have been reported.^{10,11} So we believe that suitable metal-donor joints such as Pd-N ,^{6,8,9} Ag-N ,^{7b,10} Zn-N ,^{7a,11} which can quickly associate and disassociate should be another important factor for interlocking. So far, two different polycatenanes based on discrete cages have been synthesized successfully, which may be ascribed to the following factors: firstly, suitable windows and large cavities of the cages can satisfy the interlocking; secondly, suitable metal-donor joints can provide the key intermediates for interlocking; thirdly, the strong charge-charge interactions from anions and positive frameworks make the interlocking knots stable.

In summary, two different polycatenanes constructed from different discrete cages have been prepared based on a rationally designed ligand **L** and Ag(I) . Further research indicates that the basic units for interlocking, Ag_6L_4 and Ag_5L_4 , can be generated immediately, induced by both ClO_4^- and PF_6^- anions. Possible routes of our assembly system and new design principles to prepare novel polycatenated assemblies based on discrete metallacages have been presented. Further research on preparing more complicated assemblies is ongoing.

We are thankful for financial support from the 973 Program (2011CB932504, 2011CBA00507), the National Nature Science

Foundation of China (21131006, 21141006) and the Nature Science Foundation of Fujian Province (2011J05038).

Notes and references

- (a) J. W. Steed, D. R. Turner and K. J. Wallace, *Core Concepts in Supramolecular Chemistry and Nanochemistry*, Wiley, Chichester, UK, 2007; (b) M. Groll, L. Dizel, J. Lowe, D. Stock, M. Bochter, H. D. Bartunik and R. Huber, *Nature*, 1997, **386**, 463–471.
- (a) P. J. Stang and B. Olenyuk, *Acc. Chem. Res.*, 1997, **30**, 502–518; (b) S. R. Seidel and P. J. Stang, *Acc. Chem. Res.*, 2002, **35**, 972–983; (c) M. Fujita, *Chem. Soc. Rev.*, 1998, **27**, 417–425; (d) M. Fujita, M. Tominaga, A. Hori and B. Therrien, *Acc. Chem. Res.*, 2005, **38**, 369–378; (e) D. L. Caulder and K. N. Raymond, *Acc. Chem. Res.*, 1999, **32**, 975–982; (f) N. C. Gianneschi, M. S. Masar III and C. A. Mirkin, *Acc. Chem. Res.*, 2005, **38**, 825–837; (g) C. G. Oliver, P. A. Ulman, M. J. Wiester and C. A. Mirkin, *Acc. Chem. Res.*, 2008, **41**, 1618–1629; (h) J. R. Nitschke, *Acc. Chem. Res.*, 2007, **40**, 103–112; (i) S. De, K. Mahata and M. Schmittel, *Chem. Soc. Rev.*, 2010, **39**, 1555–1575; (j) M. J. Hardie, *Chem. Soc. Rev.*, 2010, **39**, 516–527.
- (a) Y. Liu, C. Hu, A. Comotti and M. D. Ward, *Science*, 2011, **333**, 436–440; (b) Q.-F. Sun, S. Sato and M. Fujita, *Nat. Chem.*, 2012, **4**, 330–333; (c) J.-R. Li and H.-C. Zhou, *Nat. Chem.*, 2010, **2**, 893–898; (d) B. Kilbas, S. Mirtschin, R. Scopelliti and K. Severin, *Chem. Sci.*, 2012, **3**, 701–704; (e) E. Alessio, M. Casanova, E. Zangrando and E. Iengo, *Chem. Commun.*, 2012, **48**, 5112–5114; (f) N. Ousaka, J. K. Clegg and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2012, **51**, 1464–1468; (g) W. Meng, J. K. Clegg and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2012, **51**, 1881–1884; (h) Q.-Q. Wang, V. W. Day and K. Bowman-James, *Angew. Chem., Int. Ed.*, 2012, **51**, 2119–2123; (i) S. H. Lim, Y. Su and S. M. Cohen, *Angew. Chem., Int. Ed.*, 2012, **51**, 5106–5109; (j) K. Xiong, F. Jiang, Y. Gai, D. Yuan, L. Chen, M. Wu, K. Su and M. Hong, *Chem. Sci.*, 2012, **3**, 2321–2325; (k) O. Chepelin, J. Ujma, P. E. Barran and P. J. Lusby, *Angew. Chem., Int. Ed.*, 2012, **51**, 4194–4197; (l) X.-J. Li, F.-L. Jiang, M.-Y. Wu, S.-Q. Zhang, Y.-F. Zhou and M.-C. Hong, *Inorg. Chem.*, 2012, **51**, 4116–4122.
- (a) V. Vajpayee, Y. H. Song, T. R. Cook, H. Kim, Y. Lee, P. J. Stang and K.-W. Chi, *J. Am. Chem. Soc.*, 2011, **133**, 19646–19649; (b) A. Granzhan, C. Schouwey, T. Riis-Johannessen, R. Scopelliti and K. Severin, *J. Am. Chem. Soc.*, 2011, **133**, 7106–7115; (c) A. Stephenson, S. P. Argent, T. R. Johannessen, I. S. Tidmarsh and M. D. Ward, *J. Am. Chem. Soc.*, 2011, **133**, 858–870; (d) N. Kishi, Z. Li, K. Yoza, M. Akita and M. Yoshizawa, *J. Am. Chem. Soc.*, 2011, **133**, 11438–11441; (e) S. C. Serra, E. Coronado, P. Gaviña, J. Ponce and S. Tatay, *Chem. Commun.*, 2011, **47**, 8235–8237; (f) I. D. Giles, H. T. Chifotides, M. Shatruk and K. R. Dunbar, *Chem. Commun.*, 2011, **47**, 12604–12606; (g) O. Zava, J. Mattsson, B. Therrien and P. Dyson, *Chem.-Eur. J.*, 2010, **16**, 1428–1431; (h) M.-C. Hong, Y.-J. Zhao, W.-P. Su, R. Cao, M. Fujita, Z.-Y. Zhou and A. S. C. Chan, *J. Am. Chem. Soc.*, 2000, **122**, 4819–4820; (i) N. Li, F.-L. Jiang, L. Chen, X.-J. Li, Q.-H. Chen and M.-C. Hong, *Chem. Commun.*, 2011, **47**, 2327–2329.
- (a) V. Balzani, A. Credì, F. M. Raymo and J. F. Stoddart, *Angew. Chem., Int. Ed.*, 2000, **39**, 3348–3391; (b) J. V. Hernandez, E. R. Kay and D. A. Leigh, *Science*, 2004, **306**, 1532–1537.
- M. Fujita, N. Fujita, K. Ogura and K. Yamaguchi, *Nature*, 1999, **400**, 52–55.
- (a) A. Westcott, J. Fisher, L. P. Harding, P. Rizkallah and M. J. Hardie, *J. Am. Chem. Soc.*, 2008, **130**, 2950–2951; (b) J. J. Henkels, T. R. Ronson, L. P. Harding and M. J. Hardie, *Chem. Commun.*, 2011, **47**, 6560–6562.
- (a) M. Fukuda, R. Sekiya and R. Kuroda, *Angew. Chem., Int. Ed.*, 2008, **47**, 706–710; (b) R. Sekiya, M. Fukuda and R. Kuroda, *J. Am. Chem. Soc.*, 2012, **134**, 10987–10997.
- S. Freye, J. Hey, A. Torras-Galan, D. Stalke, R. Herbst-Irmer, M. John and G. H. Clever, *Angew. Chem., Int. Ed.*, 2012, **51**, 2191–2194.
- X. Kuang, X. Wu, R. Yu, J. P. Donahue, J. Huang and C.-Z. Lu, *Nat. Chem.*, 2010, **2**, 461–465.
- J. Heine, J. S. Gunne and S. Dehnen, *J. Am. Chem. Soc.*, 2011, **133**, 10018–10021.
- Using other anions such as F^- , NO_3^- , CF_3CO_2^- , and CF_3SO_3^- cannot obtain similar compounds.
- (a) D. A. Leigh, P. J. Lusby, R. T. McBurney, A. Morelli, A. M. Z. Slawin, A. R. Thomson and D. B. Walker, *J. Am. Chem. Soc.*, 2009, **131**, 3762–3771; (b) M. R. Sambrook, P. D. Beer, J. A. Wisner, R. L. Paul and A. R. Cowley, *J. Am. Chem. Soc.*, 2004, **126**, 15364–15365.
- The intermediates Ag_6L_4 and Ag_5L_4 recorded by ESI-MS may be mixtures with different conformations; attempts to find Ag_6L_4 and Ag_5L_4 by MALDI-TOF failed. Larger species such as the [2]catenane were not observed using ESI-MS.