

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/50865063>

Structural Analysis of a Ni–Methyl Species in Methyl–Coenzyme M Reductase from *Methanothermobacter marburgensis*

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MARCH 2011

Impact Factor: 12.11 · DOI: 10.1021/ja110492p · Source: PubMed

CITATIONS

13

READS

22

7 AUTHORS, INCLUDING:



Mishtu Dey

University of Iowa

22 PUBLICATIONS 309 CITATIONS

SEE PROFILE



Xianghui li

New England Biolabs

12 PUBLICATIONS 292 CITATIONS

SEE PROFILE



Ritimukta Sarangi

Stanford University

55 PUBLICATIONS 1,376 CITATIONS

SEE PROFILE



Stephen W Ragsdale

University of Michigan

201 PUBLICATIONS 7,465 CITATIONS

SEE PROFILE


Structural Analysis of a Ni-Methyl Species in Methyl-Coenzyme M Reductase from *Methanothermobacter marburgensis*

Peder E. Cedervall,[†] Mishtu Dey,^{‡,§,#} Xianghui Li,^{‡,§,||} Ritimukta Sarangi,[⊥] Britt Hedman,[⊥] Stephen W. Ragsdale,[‡] and Carrie M. Wilmot^{*,†}

[†]Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, Minnesota 55455, United States

[‡]Department of Biological Chemistry, University of Michigan, Ann Arbor, Michigan 48109, United States

[⊥]SLAC National Accelerator Laboratory, Stanford Synchrotron Radiation Lightsource, Menlo Park, California 94025, United States

 Supporting Information

ABSTRACT: We present the 1.2 Å resolution X-ray crystal structure of a Ni-methyl species that is a proposed catalytic intermediate in methyl-coenzyme M reductase (MCR), the enzyme that catalyzes the biological formation of methane. The methyl group is situated 2.1 Å proximal of the Ni atom of the MCR coenzyme F₄₃₀. A rearrangement of the substrate channel has been posited to bring together substrate species, but Ni(III)-methyl formation alone does not lead to any observable structural changes in the channel.

Methane has potential as an alternative fuel but is also a potent greenhouse gas, with surplus methanogenesis currently leading to a slow increase in methane concentrations within the atmosphere. The major source of methane produced in nature is the end product of the archaeal decomposition of organic matter in strict anaerobic environments such as lake sediments and the intestinal tract of animals.¹ In the methanogenic archaea, methyl-coenzyme M reductase (MCR) catalyzes the final and rate-limiting step in methane biogenesis: the reduction of methyl-coenzyme M (methyl-SCoM, 2-(methylthio)ethanesulfonate) by coenzyme B (CoBSH, *N*-7-mercaptoheptanoylthreonine phosphate) to methane and a heterodisulfide (Supporting Information (SI), Scheme S1). The 273 kDa MCR enzyme contains two deeply buried active sites, each containing a highly reduced Ni-tetrapyrrole cofactor, named coenzyme F₄₃₀ (F₄₃₀).² The Ni atom in F₄₃₀ is redox active and has been proposed to shuttle between oxidation states +1, +2, and +3 during turnover^{1,3,4} (Scheme 1, path 1), while other studies have suggested the involvement of only Ni oxidation states +1 and +2 (SI, Scheme S2).^{5–7} The actual mechanism remains elusive since no true catalytic intermediate of MCR has ever been observed. However, a stable MCR-Ni(III)-methyl (MCR_{Me}) state can be artificially produced by treating the resting active Ni(I) form of the enzyme (MCR_{red1}) with methyl halide (Scheme 1, path 2).^{8,9} Treatment of MCR_{Me} with coenzyme M (HSCoM, demethylated methyl-SCoM) generates MCR_{red1} along with the substrate methyl-SCoM, supporting this species as a possible catalytic intermediate.⁸

The geometric and electronic structure of MCR_{Me} was recently characterized in solution by X-ray absorption spectroscopy (XAS)

and density functional theory (DFT).¹⁰ Here we describe the X-ray crystal structure of the protein in this mechanistically relevant state. To confirm that MCR_{Me} could form in crystals, a Ni K-edge XAS spectrum was obtained from a MCR_{red1} crystal treated anaerobically with methyl iodide, which demonstrated MCR_{Me} formation in the crystal (SI Methods, Results, and Figure S1).


X-ray diffraction data were collected on a MCR_{Me} crystal, and modeling and refinement led to final *R*-values of 13.2% and 15.6% for *R*_{work} and *R*_{free}, respectively (SI Methods, Results, and Table S1). The electron density map at 1.2 Å resolution contained two distinct species bound proximal to the Ni of F₄₃₀. Besides a background of bound HSCoM (added during purification to stabilize MCR), there was clear *F*_o – *F*_c difference density, consistent with a Ni-coordinated methyl group (Figure 1A). To deconvolute the multiple species from within the active site, anomalous data generated from the native data set (wavelength 0.98 Å) and at two additional wavelengths, 1.51 and 2.29 Å, were used (Figure 1B; SI Methods, Results, Table S2, and Figure S2). The Ni–S distance refines to ~2.4 Å, which is in line with the previous crystal structures containing HSCoM^{2,11–13} and the distance reported by XAS.¹⁰ The Ni–C distance refines to 2.07 and 2.10 Å in the two crystallographically independent active sites in the asymmetric unit (the distance determined by XAS for MCR_{Me} in solution is 2.04 Å¹⁰). The substrate channel contains a mixture of coenzyme B analogue (CoB₆SH (SI, Figure S3), a slow substrate added in crystallization) and CoBSH (copurifies with the enzyme) (Figure 2). The respective thiols of CoB₆SH and the substrate CoBSH are coincident and situated ~8.7 Å away from the Ni in F₄₃₀.¹¹ As CoBSH is a substrate, the long distance between the thiol and Ni(II) in the previous inactive MCR crystal structures has been puzzling.^{2,11–13} Thus, a conformational change has been proposed that would bring the CoBSH closer to the Ni,^{11,15} which is consistent with recent transient kinetic studies revealing substrate-triggered conformational changes in the catalytic resting state MCR_{red1}.¹⁶

Our data suggest that if Ni(III)-methyl is an intermediate in methane formation, its presence alone does not affect the architecture of the MCR substrate channel, which appears

Received: November 22, 2010

Published: March 25, 2011

Scheme 1. (1) Proposed MCR Mechanism Involving a Ni(III)-Methyl Intermediate and (2) Generation of a Ni(III)-Methyl Species Using Methyl Iodide^a



^a The resting Ni(I) state is circled with a dashed line. For clarity the axial glutamine residue has been excluded.




Figure 1. Ni-methyl MCR electron density. (A) Stereo figure showing positive $F_o - F_c$ electron density (green mesh, contoured at 5σ) when methyl is not included in the structural model. $2F_o - F_c$ electron density (contoured at 1σ) is shown as blue mesh. (B) Anomalous electron density map from X-ray diffraction data collected at wavelength 2.29 Å (orange mesh, contoured at 5σ) (f'' (electrons): sulfur, 1.14; nickel, 1.03). Coenzymes are shown in stick colored by atom (carbons from CoB₆SH, pale yellow; CoBSH, pale green; HSCoM, light gray); coenzyme F₄₃₀, dark gray with the Ni atom shown as a green sphere, iodide as a purple sphere, and a water molecule as a red sphere. Protein is shown as cartoon.

unchanged compared to the structure of the channel in the Ni(II) structures. The release of methane involves a protonation step, which has been postulated to be provided by CoBSH either via a CoM species (Scheme 1, path 1) or possibly via a water molecule (Figure 2), two Tyr residues² (Tyr α 333 and Tyr β 367, whose hydroxyls point toward the Ni), or the C-ring nitrogen of F₄₃₀.³ Since no catalytic intermediates of MCR have been observed thus far by fast spectroscopic techniques, any Ni(III)-methyl catalytic intermediate formed from the native substrates must be fleeting during turnover. Recently transient kinetic methods using the natural substrate, methyl-SCoM, and the slow substrate CoB₆SH enabled observation of a short-lived Ni(III)-alkyl intermediate and its subsequent decay at catalytically competent rates.¹⁶ In contrast, when Ni(III)-methyl is artificially generated using methyl iodide, as in this study, it is relatively stable, enabling spectroscopic and biochemical characterization.^{8,9}




Figure 2. Final structural model and electron density. The final model contains a 60:40 mixture of coenzyme B₆ (CoB₆SH) and coenzyme B (CoBSH), 50% coenzyme M (HSCoM), 50% methyl, and 35% iodide. Coenzymes are shown in stick colored by atom (carbon from CoB₆SH, pale yellow; CoBSH, pale green; HSCoM, light gray); coenzyme F₄₃₀, dark gray with the Ni atom shown as a green sphere, methyl displayed as a gray sphere, iodide as a purple sphere, and a water molecule as a red sphere. Protein is shown as cartoon. $2F_o - F_c$ electron density (contoured at 1σ) is shown as blue mesh.

Interestingly, MCR is also implicated in “reverse” methanogenesis in the anaerobic oxidation of methane within microbial mats found at methane seeps in the deep ocean.^{17,18} The dissociation energy of the C–H bond (+439 kJ/mol) is larger than in any other organic compound except benzene. Thauer and Shima have proposed that in “reverse” methanogenesis only the strong electrophile F₄₃₀-Ni(III) of MCR is capable of breaking the C–H bond in a modified “reversed” version of the Ni(III)-methyl mechanism proposed for methanogenesis (Scheme 1, path 1).¹⁹ The recent evidence of a Ni-(σ -alkane) intermediate in the reaction of MCR is consistent with this proposal.²⁰

The crystal structure of MCR_{Me} is the first structure resembling a postulated intermediate in one of the proposed mechanisms of MCR. The high resolution of the X-ray diffraction data, coupled with anomalous diffraction at several wavelengths, has enabled us to extract the relevant Ni-methyl structure from confounding species. The active Ni(I) enzyme, its complex with methyl-SCoM, substrate channel conformational changes, and true catalytic intermediates trapped during turnover still wait to be structurally defined.

■ ASSOCIATED CONTENT

S Supporting Information. Methods, supporting results, X-ray crystallographic data table, anomalous signal table, single-crystal X-ray absorption spectra, anomalous electron density figure, drawing of CoBSH and analogue CoB₆SH, EPR spectra, catalytic reaction scheme, and enzymatic mechanism scheme. This material is available free of charge via the Internet at <http://pubs.acs.org>. The structure is deposited in the Protein Data Bank with PDB entry 3POT.

■ AUTHOR INFORMATION

Corresponding Author

wilmo004@umn.edu

Present Addresses

[#]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, United States^{||}Shanghai R&D Center, New England Biolabs, Inc., Shanghai, China 201204

Author Contributions

^SThese authors contributed equally to this work

■ ACKNOWLEDGMENT

This work was supported by U.S. Department of Energy grant DE-FG02-08ER15931 to S.W.R and supplement to C.M.W. and by Minnesota Partnership for Biotechnology and Medical Genomics grant SPAP-05-0013-P-FY06. We thank Stephan L. Ginell and the staff at the SBC-CAT beamline 19-ID-D at the Advanced Photon Source, Argonne National Laboratory, for valuable assistance during X-ray data collection. Argonne is operated by UChicago Argonne, LLC, for the U.S. Department of Energy, Office of Biological and Environmental Research, under contract DE-AC02-06CH11357. SSRL operations are funded by the U.S. Department of Energy, Office of Basic Energy Sciences. The SSRL Structural Molecular Biology program is supported by the National Institutes of Health, National Center for Research Resources, Biomedical Technology Program, and the U.S. Department of Energy, Office of Biological and Environmental Research. This publication was supported by Grant Number 5 P41 RR001209 from the National Center for Research Resources (NCRR), a component of the National Institutes of Health (NIH). Its contents are solely the responsibility of the authors and do not necessarily represent the official view of NCRR or NIH. Computer resources were provided by the Basic Sciences Computing Laboratory of the University of Minnesota Supercomputing Institute, and we thank Can Ergenekan for his support. We also thank Ed Hoeffner at the Kahlert Structural Biology Laboratory (KSBL) at the University of Minnesota.

■ REFERENCES

- (1) Thauer, R. K. *Microbiology* **1998**, *144*, 2377–2406.
- (2) Ermler, U.; Grabarse, W.; Shima, S.; Goubeaud, M.; Thauer, R. K. *Science* **1997**, *278*, 1457–1462.
- (3) Duin, E. C.; McKee, M. L. *J. Phys. Chem. B* **2008**, *112*, 2466–2482.
- (4) Horng, Y. C.; Becker, D. F.; Ragsdale, S. W. *Biochemistry* **2001**, *40*, 12875–12885.
- (5) Chen, S. L.; Pelmenschikov, V.; Blomberg, M. R.; Siegbahn, P. E. *J. Am. Chem. Soc.* **2009**, *131*, 9912–9913.
- (6) Pelmenschikov, V.; Blomberg, M. R. A.; Siegbahn, P. E. M.; Crabtree, R. H. *J. Am. Chem. Soc.* **2002**, *124*, 4039–4049.
- (7) Pelmenschikov, V.; Siegbahn, P. E. *J. Biol. Inorg. Chem.* **2003**, *8*, 653–662.
- (8) Dey, M.; Telser, J.; Kunz, R. C.; Lees, N. S.; Ragsdale, S. W.; Hoffman, B. M. *J. Am. Chem. Soc.* **2007**, *129*, 11030–11032.
- (9) Yang, N.; Reiher, M.; Wang, M.; Harmer, J.; Duin, E. C. *J. Am. Chem. Soc.* **2007**, *129*, 11028–11029.
- (10) Sarangi, R.; Dey, M.; Ragsdale, S. W. *Biochemistry* **2009**, *48*, 3146–3156.
- (11) Cedervall, P. E.; Dey, M.; Pearson, A. R.; Ragsdale, S. W.; Wilmot, C. M. *Biochemistry* **2010**, *49*, 7683–7693.
- (12) Grabarse, W.; Mahlert, F.; Duin, E. C.; Goubeaud, M.; Shima, S.; Thauer, R. K.; Lamzin, V.; Ermler, U. *J. Mol. Biol.* **2001**, *309*, 315–330.
- (13) Grabarse, W.; Mahlert, F.; Shima, S.; Thauer, R. K.; Ermler, U. *J. Mol. Biol.* **2000**, *303*, 329–344.
- (14) Dey, M.; Kunz, R. C.; Lyons, D. M.; Ragsdale, S. W. *Biochemistry* **2007**, *46*, 11969–11978.
- (15) Ebner, S.; Jaun, B.; Goenrich, M.; Thauer, R. K.; Harmer, J. *J. Am. Chem. Soc.* **2010**, *132*, 567–575.
- (16) Dey, M.; Li, X.; Kunz, R. C.; Ragsdale, S. W. *Biochemistry* **2010**, *49*, 10902–10911.
- (17) Hallam, S. J.; Girguis, P. R.; Preston, C. M.; Richardson, P. M.; DeLong, E. F. *Appl. Environ. Microbiol.* **2003**, *69*, 5483–5491.
- (18) Scheller, S.; Goenrich, M.; Boecher, R.; Thauer, R. K.; Jaun, B. *Nature* **2010**, *465*, 606–608.
- (19) Thauer, R. K.; Shima, S. *Ann. N.Y. Acad. Sci.* **2008**, *1125*, 158–170.
- (20) Scheller, S.; Goenrich, M.; Mayr, S.; Thauer, R. K.; Jaun, B. *Angew. Chem., Int. Ed.* **2010**, *49*, 8112–8115.