

# Synthesis enables a structural revision of the *Mycobacterium tuberculosis*-produced diterpene, edaxadiene†

Jillian E. Spangler, Cheryl A. Carson and Erik J. Sorensen\*

Received 27th April 2010, Accepted 29th April 2010

DOI: 10.1039/c0sc00284d

A stereodivergent synthesis of the [3.3.1] bicyclic core of edaxadiene was completed utilizing a key intramolecular oxidative ketone allylation. Significant discrepancies between the spectroscopic data obtained for the synthetic construct and the natural isolate raised questions about the structural assignment of edaxadiene. A subsequent structural reassignment was validated by completion of a total synthesis of the correct structure of the natural product.

Tuberculosis is a pulmonary disease caused by the pathogen *Mycobacterium tuberculosis*. Roughly two billion people, or one-third of the world's population, are believed to be infected by the bacterium, resulting in over 1.5 million deaths each year.<sup>1</sup> Despite these grave statistics, details about the infectivity and virulence of *M. tuberculosis* are only partially understood.<sup>2</sup> Peters and co-workers recently disclosed their isolation and structural elucidation of a halimane-type diterpenoid, edaxadiene (**1**), a compound produced by *M. tuberculosis*.<sup>3</sup> The genetic operon responsible for the production of edaxadiene is present only in pathogenic strains of mycobacteria.<sup>4</sup> In addition, edaxadiene was demonstrated to be an *in vitro* inhibitor of macrophage maturation.<sup>3</sup> These experimental observations suggest that production of edaxadiene is crucial for the infectivity and virulence of *M. tuberculosis*.

Our group has a growing interest in molecules related to the biology and treatment of tuberculosis.<sup>5</sup> The biological importance as well as the scarcity of edaxadiene (**1**) led us to actively pursue a total synthesis of this intriguing natural product. Our goal was to produce sufficient quantities of this molecule to confirm its structural assignment and enable a search for its biomolecular target.

We recognized that at the heart of this challenge is the synthesis of a [3.3.1] bicyclic core bearing five contiguous stereogenic centers, including two all-carbon quaternary stereogenic centers. In addition, the relative configuration of the C<sub>13</sub> stereogenic center was not assigned during structural elucidation. Although edaxadiene (**1**) has conservation of the halimane skeleton (Fig. 1), formation of a unique C<sub>7</sub>–C<sub>13</sub> bond locks the conformation of the B-ring and produces a highly unfavorable 1,3-diaxial interaction between the C<sub>17</sub> methyl group and C<sub>1</sub> methylene that is absent in related diterpenes.<sup>6</sup>

We envisioned that edaxadiene could arise from enone **3** by annulation of the leftmost ring onto the convex face of the [3.3.1] bicycle with subsequent deoxygenation of the C<sub>6</sub> ketone (**4**) to form the C<sub>5</sub>–C<sub>6</sub> olefin (Scheme 1). We reasoned that this enone (**3**) could be fashioned through an intramolecular ketone





Fig. 1 Proposed structure of edaxadiene. Stereochemistry is undefined at the starred (\*) carbon.




Scheme 1 Proposed synthesis of **1** via an intramolecular ketone allylation. Stereochemistry is undefined at the starred (\*) carbon.

allylation of an enone of type **2** to forge the C<sub>7</sub>–C<sub>13</sub> bond. Importantly, we anticipated that variation of olefin geometry and reaction conditions in this cyclization would provide access to both epimers at the unassigned C<sub>13</sub> stereogenic center.

Our efforts thus began with the development of a synthesis of enone **2**. The thermal Diels–Alder cycloaddition of tigraldehyde and a modified Rawal diene<sup>7</sup> (**5**) provided silyl enol ether **6** (Scheme 2). Subsequent methylenation of the aldehyde provided alkene **7** in good yield on a large scale (>15 g). Regioselective hydroboration of the terminal olefin with 9-BBN furnished a *B*-alkyl borane that was subjected to a Suzuki–Miyaura cross-coupling reaction<sup>8</sup> with (*Z*)-2-bromo-2-butene (**10a**) or (*E*)-2-bromo-2-butene (**10b**). The geometrical isomers **9a** and **9b** were obtained after deprotection of the enol ether and elimination of oxazolidinone with tetra-*n*-butylammonium fluoride.

Department of Chemistry, Princeton University, Frick Chemical Laboratory, Princeton, NJ, 08544, USA. E-mail: ejs@princeton.edu

† Electronic supplementary information (ESI) available: Experimental procedures and spectroscopic data; interview with Erik Sorensen. See DOI: 10.1039/c0sc00284d




**Scheme 2** Diels–Alder-mediated synthesis of enones **9a,b**. Conditions: (a) tiglaldehyde, toluene, 120 °C, 80%; (b)  $Ph_3PCH_2Br$ ,  $n-BuLi$ , THF, 0 °C, 67%; (c) **8a**: 9-BBN, THF, reflux; then 10 mol%  $PdCl_2(dppf)$ , aq.  $K_3PO_4$ , **10a**, DMF, 50 °C; **8b**: 9-BBN, THF, reflux; then 10 mol%  $PdCl_2(dppf)$ , aq.  $K_3PO_4$ , **10b**, DMF, 50 °C; (d) **9a**: TBAF, THF, 86% (2 steps); **9b**: TBAF, THF, 100% (2 steps). 9-BBN = 9-borabicyclo(3.3.1)nonane; dppf = 1,1'-bis(diphenylphosphino)ferrocene; TBAF = tetra-*n*-butylammonium fluoride.

After exploring multiple methods for constructing the  $C_7$ – $C_{13}$  bond to provide a [3.3.1] bicycle we found that oxidative cyclization of **9a** with a bimetallic system of manganese(III) acetate and copper(II) acetate in a mixture of acetic acid and benzene (3 : 1) provided the volatile [3.3.1] bicycle **12a** in moderate yield as a single stereoisomer (Scheme 3).<sup>9</sup> This cyclization most likely proceeds through the intermediacy of an A(1,3) minimized intermediate (**11a**) to provide **12a**. Utilizing the same reaction conditions enone **9b** underwent cyclization to provide **12b**, the opposite  $C_{13}$  epimer, albeit with a lower degree of diastereoselectivity. We were pleased to find that we could gain access to both  $C_{13}$  epimers by controlling the geometry of the starting trisubstituted olefin.


It was at this point in our efforts that we began to note discrepancies between the reported  $^{13}C$  shifts of the [3.3.1] core of edaxadiene and our synthetic constructs. To address our

scepticism of the proposed structure, we converted the diastereomeric bicycles **12a,b** into dienes **14a,b**, bearing the full [3.3.1] bicyclic core of structure **1**. Thus, stepwise reduction of each enone to the saturated alcohol (**13a** and **13b**) was followed by dehydration to the corresponding alkene (**14a** and **14b**) with Martin sulfurane in refluxing benzene.<sup>10</sup> The unusually high temperature requirement for the dehydration is attributed to the equatorial disposition of the substrate alcohols. Although the high volatility and non-polarity of these compounds complicated their isolation, we were able to obtain sufficient quantities of each  $C_{13}$  epimer for characterization *via* semi-preparative gas chromatography. The spectroscopic data for each  $C_{13}$  epimer differed dramatically from the reported spectra of edaxadiene; most notably, the reported  $^{13}C$ -NMR chemical shift at  $C_{13}$  was substantially different ( $\Delta\delta$  of >30 ppm) from the  $^{13}C$  chemical shifts observed for bicycles **14a,b** (Fig. 2).


These significant differences between the  $^{13}C$ -NMR data of the isolated **1** and our synthetic bicycles (**14a** and **14b**) led to our re-evaluation of the reported spectroscopic data for **1**. Our consideration of this data led us to propose compound **15** (Fig. 3) as the actual structure of edaxadiene. This proposal is consistent with the  $^{13}C$ -NMR data, most notably the deshielded nature of  $C_{13}$ . Observation of the parent  $[M - H_2O]^+$  ion by EI mass spectrometry is presumably due to the facile fragmentation of the



**Fig. 2** Discrepancies in  $^{13}C$ -NMR shifts that indicate the structure of edaxadiene has been incorrectly assigned.



**Scheme 3** Synthesis of the [3.3.1] bicyclic core of **1** *via* a stereodivergent intramolecular oxidative ketone allylation. Conditions: (a)  $Mn(OAc)_3 \cdot 2H_2O$ ,  $Cu(OAc)_2 \cdot H_2O$ , 80–100 °C,  $AcOH$ –benzene (1 : 3), **12a**: 57%, **12b**: 51% (1.4 : 1 dr); (b)  $NaBH_4$ , MeOH; (c)  $LiAlH_4$ ,  $Et_2O$ , 0 °C, **13a**: 87% (2 steps), **13b**: 73% (2 steps); (d) Martin sulfurane, benzene, 80 °C, **14a**: 23%, **14b**: 34%.



**Fig. 3** Structures of nosyberkol and tuberculosinol. Stereochemistry is unassigned at the starred (\*) carbon.

tertiary allylic alcohol at C<sub>13</sub> of **15**. We further suspect that edaxadiene, as reported, is likely not a pure compound, but rather a diastereoisomeric mixture at the remote C<sub>13</sub> stereogenic center, as indicated by fine doubling of the olefinic protons in the <sup>1</sup>H spectrum of **1**. Our confidence in this reassignment was bolstered by a comparison of this structure to known terpenes in the literature, which indicated that this was a previously identified halimane diterpene, nosyberkol (**15**), isolated in 2004 from extracts of the Red Sea sponge *Raspailia* sp.<sup>11</sup> Furthermore nosyberkol (**15**, also referred to as isotuberculosinol) was previously speculated to be the product of the Rv3378c enzyme produced by *M. tuberculosis*.<sup>12</sup>

This proposed structural reassignment led us to develop a synthesis of nosyberkol (**15**) for structural verification.<sup>13</sup> Our efforts began with an *exo*-selective Diels–Alder cycloaddition<sup>14</sup> of diene **17** (available in two steps from 2,2-dimethylcyclohexanone)<sup>15</sup> and ethyl tiglate to give the desired cycloadduct as an inseparable mixture of diastereoisomers (2 : 1) (Scheme 4). Subsequent reduction of the esters and separation of the primary

alcohols provided the desired C<sub>10</sub> epimer.<sup>16</sup> Aldehyde **18** was obtained in good yield using the conditions of Parikh and Doering.<sup>17</sup> Our attempts to perform a homologation of **18** through reaction with a phosphonate or phosphonium ylide led to no product formation; however, an aldol condensation with the sodium enolate of acetone and conjugate reduction of the resultant enone with Wilkinson's catalyst<sup>18</sup> allowed the desired homologation to ketone **20**. Our synthesis of nosyberkol was completed by the addition of vinylmagnesium bromide to the ketone to provide the desired compound as a 1.5 : 1 diastereomeric mixture at the tertiary allylic alcohol-bearing carbon. The spectroscopic data obtained for synthetic nosyberkol (**15**) were identical with those reported for both natural nosyberkol and edaxadiene (Fig. 2, see ESI†).<sup>19</sup>

We were also intrigued by the reported biomimetic conversion of tuberculosinol<sup>20</sup> (**16**) (Fig. 3) into edaxadiene (**15**) by treatment with a mixture of copper(II) chloride and *N,N'*-dicyclohexylcarbodiimide (DCC).<sup>3</sup> To examine this conversion we pursued a short synthesis of tuberculosinol from aldehyde **18**. Thus methylation of the aldehyde provided diene **19** in good yield. Subsequent regioselective hydroboration with 9-BBN and a palladium-mediated cross-coupling with (*E*)-3-iodobut-2-en-1-ol<sup>21</sup> (**21**) (Scheme 4) provided tuberculosinol (**16**). On exposure to catalytic copper(II) chloride tuberculosinol (**16**) was converted into nosyberkol (**15**) (Scheme 4). We found that addition of DCC is unnecessary for this transformation and propose that this is a Lewis acid-mediated allylic transposition.<sup>22</sup> In addition, we observed elimination of the allylic alcohol to provide a mixture of dehydrated products (34% as an *E/Z* mixture at the C<sub>12</sub>–C<sub>13</sub> olefin); however, formation of the proposed structure **1** was never observed.


In conclusion, a rapid and stereodivergent synthesis of the core of the originally proposed structure of edaxadiene (**1**) raised questions regarding its assignment. Re-evaluation of the spectral data led to the proposal that edaxadiene is actually nosyberkol (**15**), a known diterpene previously isolated from *Raspailia* sp. Furthermore, an independent synthesis of nosyberkol has unambiguously established this structural revision.<sup>23</sup> This synthesis of nosyberkol should provide sufficient quantities of this compound to elucidate its role in the infectivity and virulence of *M. tuberculosis*.

## Acknowledgements

We gratefully acknowledge Princeton University, the National Science Foundation (Graduate Research Fellowship for J. E. S.), the Natural Sciences and Engineering Research Council of Canada (Postdoctoral Fellowship for C. A. C.), and the Merck Research Laboratory for supporting this work. For invaluable technical assistance during the course of this project we gratefully acknowledge Lotus Separations (preparative SFC) and John Eng (high resolution EI mass spectrometry).

## Notes and references

- 1 World Health Organization, Fact Sheet 104, March 2010.
- 2 D. G. Russell, *Nat. Rev. Microbiol.*, 2007, **5**, 39–47.
- 3 F. M. Mann, M. Xu, X. Chen, D. B. Ulton, D. G. Russell and R. J. Peters, *J. Am. Chem. Soc.*, 2009, **131**, 17526–17527; F. M. Mann, S. Prisc, H. Hu, M. Xu, R. M. Coates and R. J. Peters, *J. Biol. Chem.*, 2009, **284**, 23574–23579.



**Scheme 4** Syntheses of nosyberkol and tuberculosinol via an *exo*-selective Diels–Alder reaction. Conditions: (a) ethyl tiglate, neat, 160 °C, 71% (2 : 1 *exo* : *endo*); (b) LiAlH<sub>4</sub>, THF, 40 °C, 56% (+24% *endo* isomer); (c) SO<sub>3</sub>·pyridine, NEt<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>–DMSO, 0 °C, 86%; (d) acetone, NaHMDS, THF, –78 → 23 °C, 87%; (e) 10 mol% Rh(PPh<sub>3</sub>)<sub>3</sub>Cl, HSiEt<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, 40 °C, 83%; (f) vinylmagnesium bromide, THF, 0 °C, 93% (1.5 : 1 dr at the starred (\*) carbon); (g) Ph<sub>3</sub>PCH<sub>3</sub>Br, KHMDS, 0 °C, THF, 91%; (h) 9-BBN, THF, 80 °C; then 10 mol% PdCl<sub>2</sub>(dppf), Ph<sub>3</sub>As, Cs<sub>2</sub>CO<sub>3</sub>, **21**, DMF, 73%; (i) 20 mol% CuCl<sub>2</sub>, acetone, 20% (dr = 1 : 1 at the starred (\*) carbon). 9-BBN = 9-borabicyclo(3.3.1)nonane; dppf = 1,1'-bis(diphenylphosphino)ferrocene.

- 4 K. Pethe, D. L. Swenson, S. Alonso, J. Anderson, C. Wang and D. G. Russell, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 13642–13647.
- 5 S. D. Tilley, K. P. Reber and E. J. Sorensen, *Org. Lett.*, 2009, **11**, 701–703.
- 6 E. Breitmaier, *Terpenes*, J. Wiley-VCH, Germany, 2006.
- 7 Y. Huang, T. Iwama and V. H. Rawal, *J. Am. Chem. Soc.*, 2000, **122**, 7843–7844; J. M. Janey, T. Iwama, S. A. Kozmin and V. H. Rawal, *J. Org. Chem.*, 2000, **65**, 9059–9068.
- 8 N. Miyaara, T. Ishiyama, H. Sasaki, M. Ishikawa, M. Satoh and A. Suzuki, *J. Am. Chem. Soc.*, 1989, **111**, 314–321.
- 9 S. A. Kates, M. A. Dombroski and B. B. Snider, *J. Org. Chem.*, 1990, **55**, 2427–2436; B. B. Snider and B. McCarthy Cole, *J. Org. Chem.*, 1995, **60**, 5376–5377; B. McCarthy Cole, L. Han and B. B. Snider, *J. Org. Chem.*, 1996, **61**, 7832–7847; B. B. Snider, *Tetrahedron*, 1996, **52**, 6073–6084.
- 10 J. C. Martin and R. J. Arhart, *J. Am. Chem. Soc.*, 1971, **93**, 4327–4329.
- 11 A. Rudi, M. Akinin, E. Gaydou and Y. Kashman, *J. Nat. Prod.*, 2004, **67**, 1932–1935.
- 12 C. Nakano, T. Sato and T. Hoshino, *Koen Yoshishu – Koryo, Terupen oyobi Seiyu Kagaku ni kansuru Toronkai*, 2005, **49**, 247–249.
- 13 Unfortunately the published spectrum of nosyberkol (**15**) was provided in CDCl<sub>3</sub>, whereas the published spectrum of the proposed edaxadiene (**1**) was provided in C<sub>6</sub>D<sub>6</sub>, preventing a direct spectral comparison.
- 14 D. S. de Miranda, G. J. A. de Conceição, J. Zukerman-Schpector, M. C. Guerrero, U. Schuchardt, A. C. Pinto, C. M. Rezende and J. Marsaioli, *J. Braz. Chem. Soc.*, 2001, **12**, 391–402.
- 15 S. P. Tanis and Y. M. Abdallah, *Synth. Commun.*, 1986, **16**, 251–259.
- 16 We could further separate the two enantiomers of the *exo* primary alcohols to provide enantioenriched material (>99% ee) through purification with supercritical fluid chromatography (see ESI†).
- 17 J. R. Parikh and W. E. Doering, *J. Am. Chem. Soc.*, 1967, **89**, 5505–5507.
- 18 I. Ojima and T. Kogure, *Tetrahedron Lett.*, 1972, **13**, 5035–5038.
- 19 The stereochemistry of C<sub>13</sub> was unassigned in the original isolation of nosyberkol (ref. 11). It is also unclear whether the original isolate was a single diastereoisomer at this stereogenic center.
- 20 C. Nakano, T. Okamura, T. Sato, T. Dairi and T. Hoshino, *Chem. Commun.*, 2005, 1016–1018.
- 21 S. Chen, R. F. Horvath, J. Joglar, M. J. Fisher and S. J. Danishefsky, *J. Org. Chem.*, 1991, **56**, 5834–5845.
- 22 K. Arata and C. Matsuura, *Chem. Lett.*, 1989, 1797; W. Yu, M. Wen, L. Yang and Z. L. Liu, *Chin. Chem. Lett.*, 2002, **13**, 495–496.
- 23 Snider and co-workers independently discovered that the original structural assignment for edaxadiene is actually nosyberkol. Their studies, which also include a synthesis of nosyberkol, were published in *Organic Letters* (see: N. Mangel, F. M. Mann, M. L. Hillwig, R. J. Peters and B. B. Snider, *Org. Lett.*, 2010, **12**, 2626–2629) after this manuscript was accepted for publication.