

Tripotric 2,4,6-tris(organoamino)-1,3,5-triazenes as precursors to multi-site triazenate ligands

Frederic Rivals and Alexander Steiner*

Department of Chemistry, University of Liverpool, Crown Street, Liverpool, UK L69 7ZD.
E-mail: A.Steiner@liv.ac.uk

Received (in Cambridge, UK) 20th June 2001, Accepted 6th September 2001

First published as an Advance Article on the web 1st October 2001

2,4,6-Tris(2-fluoroanilino)-1,3,5-triazene successively undergoes one-, two- and three-fold deprotonation in the presence of BuⁿLi; the dilithiated triazenate exists as the dimeric complex (thf)₆Li₄[(RN)₂(RNH)C₃N₃]₂ in the solid state (R = 2-F-C₆H₄) featuring bidentate N_{endo}-C-N_{exo} chelation sites.

The resemblance of monoanionic amidinates [RC(NR)₂]^{−1} and bis-imino phosphinates [R₂P(NR)₂]^{−2} in reaction and coordination behaviour corroborate the diagonal relationship between carbon and phosphorus. Both are sterically demanding, electron-rich bidentate chelates, which can be obtained from the protic precursor [RC(NR)(NHR)] and [R₂P(NR)(NHR)], respectively. Recently, we have shown that hexakis(organoamino)cyclotriphosphazenes (RNH)₆P₃N₃ act as multiprotic acids in the presence of strong bases yielding trianionic [(RNH)₃(RN)₃P₃N₃]^{3−} and hexaanionic phosphazenes [(RN)₆P₃N₃]^{6−}.³ Considering the diagonal C-P relationship, analogous tripotric 2,4,6-tris(organoamino)-1,3,5-triazenes 1H₃ are expected to be metallated correspondingly to give anionic triazenates (Scheme 1), which promise interesting coordination modes, due to the potential three-fold symmetry of the anticipated multi-site ligands.

Here, we present initial lithiation studies of 2,4,6-tris(2-fluoroanilino)-1,3,5-triazene (1H₃), which was generated by the reaction of cyanuric chloride with an excess of 2-fluoroaniline. We monitored the reaction of 1H₃ with BuⁿLi in THF using ¹H and ¹³C NMR spectroscopy (Fig. 1). Upon addition of BuⁿLi the NH signal gradually shifts up-field and decreases in intensity until it vanishes once three equivalents of BuⁿLi have been added. The ¹³C NMR shift of C(1) moves down-field from 127 in 1H₃ to 144 ppm in Li₃1, and that of C(6) up-field from 124 (1H₃) to 115 ppm (Li₃1) (see Fig. 1 for numbering scheme). The signal of the ¹³C nuclei of the central triazene ring shifts only slightly from 165 (1H₃) to 169 ppm (Li₃1). Throughout lithiation the ¹⁹F NMR consists of a single peak, which also shifts only marginally from 130 to 132 ppm. Both ¹H and ¹³C NMR of partially deprotonated species display only one set of signals for both 2-fluoroanilino and triazene moieties, indicating rapid proton transfer at room temperature. However, two

sets of signals with an intensity ratio of 2:1 appear in the ¹³C NMR spectrum of a THF solution containing the dilithiated species Li₂1H, which was recorded at −80 °C. The chemical shifts of the set of lower intensity are similar to the spectrum of 1H₃ and can be attributed to the non-deprotonated anilino substituent, whereas those of the set of higher intensity, which resemble the pattern observed in the spectrum of Li₃1, are caused by the two deprotonated substituents. At −80 °C the ¹³C nuclei of the triazene ring also give two resonances (at 164 and 169 ppm).

Single crystals were obtained from the reaction of 1H₃ with two equivalents of BuⁿLi in THF. X-Ray structure determination revealed that two-fold deprotonation had occurred and the resulting Li₂1H exists as the centrosymmetric dimer (thf)₆Li₄(1H)₂ 2 in the solid state (Fig. 2).[†] Both ligands in 2 are linked by the two centrally arranged lithium ions (Li1, Li1a) forming an eight-membered [Li-N_{exo}-C-N_{endo}]₂ ring core. In addition, Li1 and Li1a undergo weak cross-ring interactions to N_{exo}-sites and each is coordinated to an F-aryl atom and a thf molecule resulting in distorted trigonal bipyramidal coordination geometries. The other two lithium ions (Li2, Li2a) are accommodated in neighbouring N_{exo}-C-N_{endo} chelates and are additionally coordinated to two thf molecules giving tetrahedral metal surroundings. 2 combines coordination modes of monomeric and dimeric lithium complexes of ligands featuring bidentate N-E-N chelates [E = (R)C,⁴ (R₂)P,⁵ (R)S⁶]. Li2





Fig. 1 ¹³C NMR (top) and ¹H NMR (bottom) spectra of stepwise reaction of 1H₃ with *n* equivalents of BuⁿLi in thf at 20 °C. In ¹³C NMR '+' refers to the C(1) (= C-N) and '*' to the C(6) position of 2-fluoroanilino substituents; in ¹H NMR '*' refers to the NH signal.




Fig. 2 X-Ray structure of **2**. Selected bond lengths (Å) and angles (°): N1–C1 1.339(9), N1–C3 1.351(9), N2–C1 1.343(9), N2–C2 1.359(9), N3–C3 1.336(9), N3–C2 1.341(9), C1–N10 1.353(9), C2–N20 1.334(9), C3–N30 1.374(10), Li1–N1 2.03(2), Li1–N10 2.32(2), Li1–N10a 2.07(2), Li1–F1a 2.15(2), Li2–N2 2.11(2), Li2–N20 1.99(2), Li2...F2 3.072(15); C1–N1–C3 113.1(7), C1–N2–C2 115.7(7), C2–N3–C3 114.3(7), N1–C1–N2 125.5(8), N2–C2–N3 124.0(8), N3–C3–N1 127.3(7), N1–C1–N10 111.6(8), N2–C2–N20 110.5(7), N3–C3–N30 118.5(8).

corresponds to the 'monomeric' mode, where the lithium ion is chelated by a single N–E–N unit, and Li1 to the often encountered N–E–N dimer mode, where two ligands encapsulate two lithium ions in both mono- and bi-dentate fashion. A similar motif is observed in the polymeric trilithium salt of trithiocyanuric acid.⁷

In contrast to phosphazenes, where the metallation causes a marked impact on the structural parameters of the central P₃N₉ core, bonding parameters within the central C₃N₆ ligand core of **2** are not very much affected upon metallation. The C–N_{exo} bond involving the non-metallated anilino group is only marginally longer [C3–N30 1.374(10) Å] than those of the deprotonated substituents [C1–N10 1.353(9), C2–N20 1.334(9) Å], which are of similar lengths as C–N_{endo} bonds ranging from 1.336(9) to 1.359(9) Å. Accordingly, in neutral 2,4,6-tris-(amino)-1,3,5-triazenes C–N_{exo} bonds are on average only 0.02 Å longer than C–N_{endo} bonds.⁸

1H₃ is deprotonated without undergoing side reactions. The lithiation is reversible as 1H₃ is recovered after protolysis of Li₃**1**. On the other hand, 1,3,5-triazene and 2,4,6-tris(organo)-1,3,5-triazenes are prone to nucleophilic attack by organolithium reagents at C_{endo} positions, which leads to either nucleophilic addition, substitution or ring cleavage.⁹ F(aryl) functions in **2**, initially introduced to act as ¹⁹F NMR spectator sites, are inert on lithiation. The absence of line broadening of ¹⁹F signals indicates that there are no considerable ⁷Li–¹⁹F interactions in solution.

The electron deficiency of the central C₃N₃ ring in **1**^{3–} is compensated for by the π-donating character of exocyclic N centres, which allow delocalisation of negative charge across the entire C₃N₆ core. In contrast to the non-aromatic tri- and

hexa-anionic triphosphazenes, triazenes **1**^{2–} and **1**^{3–} feature an aromatic ligand core. This enables electronic interaction between accommodated metal centres across the aromatic ligand system. In addition, the D_{3h} symmetry of the C₃N₆ ligand core and the straightforward introduction of organo amino groups with various functionalities promise the generation of metal complexes with interesting properties.

This work was supported by the EPSRC and the Royal Society.

Notes and references

† Crystal data for 2·C₄H₈O·C₆H₁₄ were collected on a Stoe-IPDS at 200 K using Mo-Kα radiation (λ = 0.71073 Å). Full-matrix, least squares refinements on F² using all data.¹⁰ C₆₆H₇₄F₆Li₄N₁₂O₆·C₄H₈O·C₆H₁₄, M = 1431.41, triclinic, space group P1̄, a = 10.121(6), b = 14.415(3), c = 15.282(3) Å, α = 68.66(3), β = 72.98(5), γ = 79.06(5)°, U = 1977.0(13) Å³, Z = 1, μ(MoKα) = 0.087 mm^{−1}, R1 [I > 2σ(I)] = 0.095, wR2 (all 3514 data) = 0.231. **2** crystallises with one molecule of thf and one molecule of hexane as lattice solvent per formula unit. Both coordinated and non-coordinated thf molecules as well as the hexane molecule are disordered and were split in the refinement on two positions using similar distance and similar U restraints. All non-hydrogen atoms were refined anisotropically with the exception of disordered atoms which were treated isotropically. **2** forms thin and highly fragile plates, which are prone to solvent loss in the absence of the mother-liquor, and give a rather weak diffraction pattern of low resolution (2θ_{max} = 40°).

CCDC reference number 166359. See <http://www.rsc.org/suppdata/cc/b1/b105435j/> for crystallographic data in CIF or other electronic format.

- 1 F. T. Edlmann, *Coord. Chem. Rev.*, 1994, **137**, 403; J. Barker and M. Kilner, *Coord. Chem. Rev.*, 1994, **133**, 219.
- 2 M. Witt and H. W. Roesky, *Chem. Rev.*, 1994, **94**, 1163.
- 3 A. Steiner and D. S. Wright, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 636; G. T. Lawson, C. Jacob and A. Steiner, *Eur. J. Inorg. Chem.*, 1999, 1881; G. T. Lawson, F. Rivals, M. Tascher, C. Jacob, J. F. Bickley and A. Steiner, *Chem. Commun.*, 2000, 341.
- 4 D. Stalke, M. Wedler and F. T. Edlmann, *J. Organomet. Chem.*, 1992, **431**, C1; J. Barker, D. Barr, N. D. R. Barnett, W. Clegg, I. Cragg-Hine, M. G. Davidson, R. P. Davies, S. M. Hodgson, J. A. K. Howard, M. Kilner, C. W. Lehmann, I. Lopez-Solera, R. E. Mulvey, P. R. Raithby and R. Snaith, *J. Chem. Soc., Dalton Trans.*, 1997, 951.
- 5 A. Steiner and D. Stalke, *Inorg. Chem.*, 1993, **32**, 1777; R. Fleischer and D. Stalke, *Inorg. Chem.*, 1997, **36**, 2413; S. Wingerter, M. Pfeiffer, A. Murso, C. Lustig, T. Stey, V. Chandrasekhar and D. Stalke, *J. Am. Chem. Soc.*, 2001, **123**, 1381.
- 6 F. Pauer and D. Stalke, *J. Organomet. Chem.*, 1991, **418**, 127; S. Freitag, W. Kolodziejski, F. Pauer and D. Stalke, *J. Chem. Soc., Dalton Trans.*, 1993, 3779.
- 7 D. R. Armstrong, J. E. Davies, N. Feeder, E. Lamb, J. J. Longridge, J. M. Rawson, R. Snaith and A. E. H. Wheatley, *J. Mol. Model.*, 2000, **6**, 234.
- 8 A. R. Katritzky, I. Ghiviriga, P. J. Steel and D. C. Oniciu, *J. Chem. Soc., Perkin Trans. 2*, 443, 1996.
- 9 W. M. Boesveld, P. B. Hitchcock and M. F. Lappert, *J. Chem. Soc., Dalton Trans.*, 1999, 4041; D. R. Armstrong, K. W. Henderson, M. MacGregor, R. E. Mulvey, M. J. Ross, W. Clegg and P. A. O'Neil, *J. Organomet. Chem.*, 1995, **486**, 79.
- 10 G. M. Sheldrick, SHELX97, X-ray structure determination program, Universität Göttingen, Germany, 1997.