

## An Efficient Stereoselective Synthesis of Co-enzyme Q<sub>10</sub>

Kikumasa Sato,\* Osamu Miyamoto, Seiichi Inoue, Tomoya Yamamoto, and Yukihiro Hirasawa

Department of Applied Chemistry, Faculty of Engineering, Yokohama National University, Tokiwadai, Hodogayaku, Yokohama 240, Japan


Co-enzyme Q<sub>10</sub> was efficiently synthesised by stereo- and regio-selective prenylation of the protected hydroquinone (**2**) with isoprene epoxide and solanesyl *p*-tolyl sulphone in a good overall yield.

Co-enzyme Q<sub>10</sub> (**1**) plays a pivotal role in several metabolic sequences and there is an increasing need for an efficient preparative method for this substance owing to its remarkable physiological and clinical activity.<sup>1</sup> Most of the existing methods involve alkylation of a protected or unprotected hydroquinone or quinone precursor with decaprenyl compounds.<sup>2,3</sup> However, the alkylating agents, such as decaprenol and decaprenyl bromide, from which other alkylating agents can be prepared, are usually obtained as a mixture of *cis* and *trans* isomers from natural solanesol.<sup>2</sup> Therefore the utility of these methods is diminished by the difficulty in isolating a single isomer of the pure alkylating reagent. Another approach to (**1**) was reported by Terao *et al.*<sup>4</sup> using a sulphone-functionalised prenylhydroquinone and solanesyl bromide; however, the stereoselective synthesis of the former component requires multi-step procedures.


Recently<sup>5</sup> we reported a stereoselective synthetic route to all-*trans*-decaprenol from geraniol *via* the coupling of poly-prenyl sulphones and a halide, and subsequent reductive elimination of the sulphone group. This methodology has now been applied to the stereoselective synthesis of (**1**).

The bromide (**2a**)<sup>6</sup> was converted into the Grignard reagent (**3a**) and treated with isoprene epoxide in the presence of a catalytic amount of copper(i) chloride in tetrahydrofuran (THF) at  $-50^{\circ}\text{C}$  to afford the *trans*-allylic alcohol (**4a**) in 77% yield. The stereochemistry of the alcohol was confirmed by the n.m.r. spectrum of the aldehyde ( $\delta_{\text{CHO}}$  9.31)<sup>7</sup> obtained by Collins oxidation of (**4a**). The alcohol (**4a**) was converted (BuLi, *p*-MeC<sub>6</sub>H<sub>4</sub>SO<sub>2</sub>Cl, and LiBr; 89% yield) into the bromide (**5a**), which was then coupled with the anion of solanesyl *p*-tolyl sulphone to give the product (**6a**) in a good yield.

The reductive elimination of the sulphone group in (**6a**)



by the usual method<sup>8</sup> (lithium-ethylamine,  $-78^{\circ}\text{C}$ ) did not afford any desired product. The normal product (**7a**), however, was obtained when (**6a**) was subjected to the modified Bouvaul-Blanc reduction<sup>9</sup> (8 equiv. of metallic sodium and 10 equiv. of ethanol in THF at room temp.). Simple acid-catalysed deprotection of (**7a**) followed by neutralisation and air oxidation furnished co-enzyme Q<sub>10</sub> (**1**) in nearly quantitative yield. Pure all-*trans*-(**1**) was obtained by silica gel column chromatography (10% THF-hexane) and recrystallisation from ethanol, m.p.  $48-49^{\circ}\text{C}$ , in 83% yield.



A similar reaction sequence was successfully applied to give another synthesis of (**1**) starting from the bromide (**2b**). The final step consisted of the modified Bouvaul-Blanc reduction (*vide supra*) of (**6b**), and reductive elimination of the benzyl protecting group in (**7b**) (Li-EtNH<sub>2</sub>,  $-78^{\circ}\text{C}$ ), followed by mild oxidation of the *p*-methoxyphenol (FeCl<sub>3</sub>, ethyl acetate-isopropyl ether).

Considering that (**2a**) is made from 2,3-dimethoxy-5-methylbenzoquinone (**9**) in three steps, and that (**2b**) is obtained in two steps from 2,3,4-trimethoxy-6-methylphenol (**8**),<sup>10</sup> an intermediate in the synthesis of (**9**), the latter route, i.e. (**2b**)  $\rightarrow$  (**4b**)  $\rightarrow$  (**6b**)  $\rightarrow$  (**1**), seems to provide the most effective synthesis of co-enzyme Q<sub>10</sub> (**1**).

Received, 2nd November 1981; Com. 1281

## References

- 1 R. H. Thomson, 'Naturally Occurring Quinones,' 2nd edn., Academic Press, New York, 1971; G. P. Littaru, L. Ho, and K. Folkers, *Int. J. Vitam. Nutr. Res.*, 1972, **42**, 291, 413; A. B. Combs, D. Acosta, and K. Folkers, *I.R.C.S. Med. Sci.: Libr. Compend.*, 1976, **4**, 403.
- 2 R. Rüegg, U. Gloor, N. R. Ryser, O. Wiss, and O. Isler, *Helv. Chim. Acta*, 1959, **42**, 2616.
- 3 S. Inoue, R. Yamaguchi, K. Saito, and K. Sato, *Bull. Chem. Soc. Jpn.*, 1974, **47**, 3098; Y. Naruta and K. Maruyama, *Chem. Lett.*, 1979, 885; *J. Org. Chem.*, 1980, **45**, 4097.
- 4 S. Terao, K. Kato, M. Shiraishi, and H. Morimoto, *J. Org. Chem.*, 1979, **44**, 868; Y. Fujita, M. Ishiguro, T. Onishi, and T. Nishida, *Synthesis*, 1981, 469.
- 5 K. Sato, S. Inoue, A. Onishi, N. Uchida, and N. Minowa, *J. Chem. Soc., Perkin Trans. 1*, 1981, 761.
- 6 K. Sato, S. Inoue, and R. Yamaguchi, *J. Org. Chem.*, 1972, **37**, 1889.
- 7 K. C. Chan, R. A. Jewell, W. H. Nutting, and H. Rapoport, *J. Org. Chem.*, 1968, **33**, 3382.
- 8 E. M. Kaiser, *Synthesis*, 1972, 391.
- 9 K. Ogura, T. Noguchi, S. Mita, and G. Tsuchihashi, 7th Congress of Organo Sulphur and Phosphorus Compounds, Kyoto, Feb. 1979, Abstr. No. 22.
- 10 K. Sato, S. Inoue, and H. Sato, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 3455.