

The first enantioselective syntheses of vicinal difluoropyrrolidines and the first catalytic asymmetric synthesis mediated by the C_2 symmetry of a –CHFCHF– unit

Charles M. Marson*† and Robert C. Melling


Department of Chemistry, Queen Mary and Westfield College, University of London, London, UK E1 4NS

The first enantiopure vicinal difluorides of C_2 symmetry have been prepared by the introduction of fluorine at both centres in a single operation; the first asymmetric synthesis using a catalyst whose chirality depends on organofluorine asymmetry is described.

The stereoselective synthesis of organofluorine compounds^{1,2} is of major importance in many fields, including pharmaceuticals,^{1–3} nucleoside and carbohydrate chemistry,^{1b} biochemistry,^{4,5} liquid crystals⁶ and polymers.⁷ Many fluorinated α -amino acids are potent antitumour and antiviral agents.^{1a,4a} The importance of monofluoro analogues as antimetabolites is illustrated by (2*R*,3*R*)-fluorocitric acid, an aconitase inhibitor that blocks the citric acid cycle.^{2,5} Additionally, organofluoro ligands can be more powerful than oxygen ligands in coordinating metals.⁸

Whereas enantiocontrolled syntheses of monofluoroorganic compounds are well established, synthesis of an enantiopure vicinal difluoro compound, especially of C_2 symmetry, has not to the best of our knowledge been reported prior to this communication.^{9,10} Generally, molecular fluorine adds to alkenes with *syn*-stereoselection, thereby precluding the formation of C_2 symmetric difluorides;¹¹ where *trans*-addition is observed yields are usually low.¹² For example diethylamino-sulfur trifluoride (DAST),¹³ one of the most commonly used reagents for the conversion of alcohols into fluorides, gives merely a trace of 1,2-difluorocyclohexanes, and with loss of stereointegrity compared with the initial cyclohexane-1,2-diol.¹⁴ SF_4 acts on (+)- or (–)-tartaric acid, exchanging both hydroxy groups for fluorine, but with complete loss of optical activity, by formation of only the *meso*-difluoroacid.^{15a} With tartrate esters, XeF_2 was similarly unsuccessful.^{10c} Despite those previous accounts, we here report the enantiocontrolled introduction of fluorine at two adjacent carbon stereocentres in a single operation, and describe syntheses of enantiopure vicinal difluorides **5**, and an asymmetric process using some of those difluorides as catalysts.


In view of reports¹⁵ that double vicinal displacements of tartaric acid derivatives by fluoride do not proceed with enantiocontrol, displacements on cyclic systems were investigated. (3*R*,4*R*)-Diacetoxysuccinic anhydride **1**¹⁶ was reacted with a primary amine (1 equiv., 12 h, 20 °C), and the intermediate amido acid treated directly with $SOCl_2$ (2 equiv., 24 h, 20 °C) to give the diacetoxypyrrolidin-2,5-dione **2** (**2a**, R = Ph; **2b**, R = *n*-C₈H₁₇; **2c**, R = cyclohexyl) (Scheme 1). The pyrrolidin-2,5-diones **2** were reduced with $NaBH_4$ –I₂ in THF (12 h) and the diols **3** liberated by a two-stage work-up involving stirring with 1:1 AcOH–HCl (10 M) for 10 h, followed by washing with methanolic KOH (2 M). Reaction of the diols **3** with Tf_2O (2 equiv., 4 h, –80 °C) in the presence of pyridine (2 equiv.) afforded the bis(trifluoromethanesulfonates) **4**. These were isolable in the cases of **4a** (R = Ph) and **4b** (R = *n*-octyl) but **4c** (R = cyclohexyl) decomposed rapidly during column chromatography. The bis(trifluoromethanesulfonates) **4** were reacted with Bu_4NF (3 equiv., 16 h, –80 to 20 °C) in THF, resulting in stereoselective introduction of



Scheme 1

fluorine with clean inversion at both centres to give the difluoropyrrolidines **5a–c**, in respective yields of 76, 83 and 40%.[‡] To the best of our knowledge, 3,4-difluoropyrrolidines have not been previously prepared, either in racemic or enantiopure form.

Catalysis of the epoxidation of allylic alcohols by difluorides **5** was investigated; reactions were conducted in CH_2Cl_2 using 15 mol% of $Ti(OPr)_4$ and 10 mol% of catalyst (Scheme 2, Table 1, entries 2–7). In the absence of a catalyst, racemic **7** was obtained in 81% yield. Diol **3a** afforded 2,3-epoxygeraniol **7** (97%) in 25% ee in favour of the (2*S*,3*S*)-enantiomer (entry 2).



Scheme 2

Table 1 Asymmetric epoxidation of geraniol (1.6 mmol) with *tert*-butyl hydroperoxide, titanium tetraisopropoxide (15 mol%) and the difluorinated catalyst **5b** (10 mol%)

Entry	Catalyst	<i>T</i> /°C	<i>t</i> /h	Yield (%)	Ee (%)	Configuration
1	—	–20 to 20	12	81	—	racemic
2	3a	–20 to –10	0.67	97	25	(<i>S,S</i>)
3	5b	–20 to 20	1	68	50	(<i>R,R</i>)
4	5b	0	1	74	51	(<i>R,R</i>)
5	5b	–20 to 20	12	90	66	(<i>R,R</i>)
6	5b	–80	3	23	27	(<i>R,R</i>)
7	5c	–20 to 20	12	87	10	(<i>R,R</i>)

The use of **5c** (−20 to 20 °C over 12 h) afforded 2,3-epoxygeraniol (87%) in 10% ee in favour of the (2*R*,3*R*)-enantiomer (entry 7). However, **5b** afforded a 90% yield of 2,3-epoxygeraniol **7** in 66% ee in favour of the (2*R*,3*R*)-enantiomer (entry 5). Entries 3–5 suggest that fluoro groups may provide greater enantioselection than hydroxy groups (entry 2), at least in the case of a C₂ vicinal unit which is part of a heterocyclic ring. The reversal of the major enantiomer of 2,3-epoxygeraniol when using catalyst **3** compared with catalyst **5** would be expected if the modes of binding of the hydroxy and fluoro catalysts had important features in common. Samples of alcohol **7** were converted into the acetate (1 equiv. Ac₂O, 1 equiv. pyridine, 10 mol% DMAP in CH₂Cl₂ at 0 to 20 °C over 2 h), and the ee determined by observation of ¹H NMR peak of the acetate methyl group upon treatment with Eu(hfc)₃;¹⁷ the acetate (10 mg in 0.5 ml of C₆D₆) was treated with consecutive portions of 10–20 ml of a filtered solution of 35 mg of Eu(hfc)₃ in 0.5 ml of C₆D₆.

The presence of fluorine ligands in organic reactions mediated by catalysis is an emerging area of importance.¹⁸ To date, however, the chirality has not been a consequence of the spatial arrangement of the fluorine atoms, but of the asymmetry of an unrelated organic ligand (*e.g.* BINOL).¹⁸ Consequently, the present examples are, to the best of our knowledge, the first examples of asymmetric synthesis catalyzed by a compound whose chirality depends upon organofluorine asymmetry.

In the catalytic asymmetric Sharpless epoxidation,¹⁹ free hydroxy groups on the catalyst (dialkyl tartrate) are a prerequisite for enantioselectivity. In marked contrast to such Sharpless catalysts, the difluorides **5** lack hydroxy groups and are incapable of deprotonation that could lead to ligand exchange, and yet **5a–c** are viable catalysts for asymmetric epoxidation.

Compounds **5a** and **5c** are particularly suitable substructures for liquid crystal applications, and difluoropyrrolidines **5** and their derivatives are currently being evaluated for use as liquid crystals and other new materials; additional catalytic processes are also under investigation.

Support from the EPSRC for a studentship (to R. C. M.) under the ROPA initiative is gratefully acknowledged.

Notes and References

† E-mail: c.m.marson@qmw.ac.uk.

‡ All compounds gave satisfactory spectral data (NMR, IR, MS), and all new compounds gave satisfactory elemental analyses or HRMS. *Selected data for 4a*: prisms, mp 126.5–127 °C (hexane), [α]_D +46.2 (c 1, CHCl₃); δ_{H} (250 MHz, CDCl₃) 7.30 (m, 2 H), 6.88 (t, *J* 9.0, 1 H), 6.60 (d, *J* 9.0, 2 H), 5.52 (t, *J* 2.5, 2 H), 3.95 (dd, *J* 11.0, 5.0, 2 H), 3.65 (dd, *J* 11.0, 3.0, 2 H); δ_{C} (62.2 MHz, CDCl₃) 145.5 (d), 129.7 (d), 118.9 (s), 118.5 (q), 112.6 (d), 85.4 (d), 51.3 (t). For **5a**: needles, mp 89.5 °C (hexane), [α]_D −40.6 (c 3.5, CHCl₃); δ_{H} (600 MHz, CDCl₃) 7.30 (m, 2 H), 6.76 (t, *J* 7.0, 1 H), 6.60 (d, *J* 7.0, 2 H), 5.30 (dm, ²*J*_{HF} 49.3, ³*J*_{HF} 12.6, 2 H), 3.70 (m, 4 H); δ_{C} (150.9

MHz, CDCl₃) 146.6 (s), 129.4 (d), 117.1 (d), 112.0 (d), 92.8 (ddd, ¹*J*_{CF} 180, ²*J*_{CF} 33), 51.6 (m); δ_{F} (564.8 MHz, CDCl₃, internal CFCl₃) −190.3 (m).

- (a) V. Aoloshonok, in *Biomedical Frontiers of Fluorine Chemistry*, ed. I. Ojima, J. R. McCarthy and J. T. Welch, American Chemical Society Symposium Series, Washington, DC, 1996, vol. 639, pp. 26–41; (b) J. R. McCarthy, P. S. Sunkasa, D. P. Matthews, A. J. Bitonti, E. T. Jarvi, J. S. Sabol, R. J. Resvick, E. W. Huber, W. A. van der Donk, G. Yu and J. Stubbe, *ibid.*, pp. 246–264; (c) M. Namchuk, C. Braun, J. D. McCarter, and S. G. Withers, *ibid.*, pp. 265–278; M. Namchuk, C. Braun, J. D. McCarter and S. G. Withers, *ibid.*, pp. 279–293.
- P. Bravo and G. Resnati, *Tetrahedron Asymmetry*, 1990, **1**, 661.
- D. Bouzard, P. Dicesare, M. Essiz, J. P. Jacquet, J. R. Kiechel, P. Remuzon, A. Weber, T. Oki, M. Masuyoshi, R. E. Kessler, J. Fungtomc and J. Desiderio, *J. Med. Chem.*, 1990, **33**, 1344.
- (a) J. T. Welch and S. Eswarakrishnan, *Fluorine in Bio-organic Chemistry*, Wiley, New York, 1991; (b) J. A. Wilkinson, *Chem. Rev.*, 1992, **92**, 505.
- E. Kun, E. Kirsten and M. L. Sharma, *Proc. Natl. Acad. Sci. USA*, 1977, **74**, 4942.
- H. Liu and H. Nohira, *Liq. Crystal.*, 1996, 581.
- C. K. Chen, Y. L. Hu, M. Spears, J. W. Hodby, B. M. Wanklyn, A. V. Narlikar and S. B. Samanta, *J. Mater. Sci. Lett.*, 1996, **15**, 886.
- H. Plenio, R. Diodone and D. Badura, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 156.
- For a single example of a racemic *vic*-difluoro-2,3-dihydrobenzo[*b*]furan, see: R. Ruzziconi and G. V. Sebastiani, *J. Heterocycl. Chem.*, 1980, **17**, 1147.
- For *meso*-difluorides see: (a) M. Hudlicky, *J. Fluorine Chem.*, 1987, **36**, 373; (b) A. Baklouti and R. El Gharbi, *J. Fluorine Chem.*, 1979, **13**, 297; (c) T. B. Patrick, S. Khazaeli, S. Nadji, K. Hering-Smith and D. Reif, *J. Org. Chem.*, 1993, **58**, 705; (d) G. A. Olah, J. T. Welch, Y. D. Vankar, M. Nojima, I. Kerekes and J. A. Olah, *J. Org. Chem.*, 1979, **44**, 3872.
- S. Rozen and M. Brand, *J. Org. Chem.*, 1986, **51**, 3607.
- M. Sato, T. Hirokawa, A. Hattori, A. Toyota and C. Kaneko, *Tetrahedron: Asymmetry*, 1994, **5**, 975.
- W. J. Middleton, *J. Org. Chem.*, 1975, **40**, 574; M. Hudlicky, *Org. React.*, 1988, **35**, 513.
- D. F. Shellhamer, D. T. Austine, K. M. Gallego, B. R. Ganesh, A. A. Hanson, K. A. Hanson, R. D. Henderson, J. M. Prince and V. L. Heasley, *J. Chem. Soc., Perkins Trans. 2*, 1995, 861.
- (a) A. I. Burmakov, L. A. Motnyak, B. V. Kunshenko, L. A. Alexeva and L. M. Yagupolskii, *J. Fluorine Chem.*, 1981, **19**, 151; (b) for a racemic route to mixtures of difluorosuccinic acid derivatives from maleic anhydride and F₂, see: R. G. Syvret, D. L. Vassilaros, D. M. Parees and G. P. Pez, *J. Fluorine Chem.*, 1994, **67**, 277.
- R. L. Shriner and C. L. Furrow, *Org. Synth.*, 1963, **Coll. Vol. IV**, 242.
- Y. Gao, R. M. Hanson, J. M. Klunder, S. Y. Ko, H. Masamune and K. B. Sharpless, *J. Am. Chem. Soc.*, 1987, **109**, 5765.
- R. O. Duthaler and A. Hafner, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 43.
- T. Katsuki and K. B. Sharpless, *J. Am. Chem. Soc.*, 1980, **102**, 5974.

Received in Liverpool, UK, 2nd March 1998; 8/01718B