

Cite this: *Chem. Commun.*, 2012, **48**, 7380–7382

www.rsc.org/chemcomm

## COMMUNICATION

# Practical synthesis of 4,4,4-trifluorocrotonaldehyde: a versatile precursor for the enantioselective formation of trifluoromethylated stereogenic centers *via* organocatalytic 1,4-additions†


Kazutaka Shibatomi,\* Akira Narayama, Yoshiyuki Abe and Seiji Iwasa

Received 18th April 2012, Accepted 1st June 2012

DOI: 10.1039/c2cc32757k

The practical synthesis of 4,4,4-trifluorocrotonaldehyde (**1**) and its application to enantioselective 1,4-additions are described. The organocatalytic 1,4-addition of **1** with several nucleophiles such as heteroarenes, alkylthiols and aldokimes afforded the corresponding products, each bearing a trifluoromethylated stereogenic center with high optical purity. A resulting product was converted into an MAO-A inhibitor, beflouxatone.


Incorporation of perfluoroalkyl substituents into organic molecules is a highly valuable process in pharmaceutical, agrochemical and material sciences. In particular, introduction of a trifluoromethyl (CF<sub>3</sub>) group into biologically active compounds often modifies their physical and/or biological properties such as lipophilicity, metabolic stability, and bioavailability.<sup>1</sup> Although a large number of pharmaceuticals contain the CF<sub>3</sub> group(s), developing flexible methods for the highly enantioselective construction of trifluoromethylated stereogenic centers remains a challenging task. Trifluoromethylated chiral carbon centers are mostly constructed by either of these approaches: (i) asymmetric direct trifluoromethylation of a prochiral carbon; (ii) asymmetric functionalization of a trifluoromethylated prochiral carbon. Although the former approach is simpler and straightforward, only a few catalytic methods that provide high enantioselectivity are known.<sup>2</sup> Therefore, the latter approach, the so-called building block approach, has often been adopted to obtain a chiral center bearing a CF<sub>3</sub> moiety, mainly by using trifluoromethyl ketones, trifluoroacetaldehyde, trifluoropyruvates or their derivatives as precursors.<sup>3</sup> Intrigued by the fact that 4,4,4-trifluorocrotonates are simple building blocks of various trifluoromethyl compounds, some research groups, including our group, have applied them to asymmetric transformations.<sup>4</sup> During the course of our continuous research in this direction, we envisaged that 4,4,4-trifluorocrotonaldehyde (**1**) would be a versatile precursor for the construction of trifluoromethylated stereogenic centers (Scheme 1). However, to the best of our knowledge,



Scheme 1 Synthetic strategy.

there are only four reports on the synthesis of **1**,<sup>5</sup> but no report on the isolation of this compound or its use in enantioselective transformations, although  $\alpha,\beta$ -enals have been widely used as precursors in numerous organocatalytic asymmetric transformations.<sup>6</sup> One possible reason for this is the high volatility of **1** and the consequent difficulty in its purification. In the present research, we successfully isolated **1** for the first time and applied it to enantioselective 1,4-addition with several nucleophiles in the presence of the Jørgensen–Hayashi-type prolinol ether catalyst **2**.<sup>7</sup>

We first focused on the development of an efficient method for the synthesis of **1**. Since **1** is highly volatile, we decided to avoid the use of liquid–liquid extraction and column chromatography for the isolation and purification of this compound. Thus, we planned to synthesize **1** by the oxidation of 4,4,4-trifluoro-2-butenol (**4**) with manganese dioxide, which can be removed from the reaction system by a simple filtration process. First, **4** was synthesized in 61% yield by the reduction of ethyl 4,4,4-trifluorocrotonate (**3**) with lithium aluminum hydride and aluminum trichloride.<sup>8</sup> Fortunately, the manganese dioxide oxidation of **4** proceeded smoothly to afford the desired product with high conversion (85%, determined by <sup>1</sup>H NMR analysis using an internal standard). After the removal of manganese dioxide by filtration, the filtrate was distilled to afford nearly pure **1** (98% purity) in 47% yield on a 5 g scale (Scheme 2).






Scheme 2 Synthesis of 4,4,4-trifluorocrotonaldehyde.

Department of Environmental and Life Sciences, Toyohashi University of Technology, 1-1 Hibarigaoka, Tempaku-cho, Toyohashi 441-8580, Japan. E-mail: shiba@ens.tut.ac.jp; Fax: +81 532 48 5833; Tel: +81 532 44 6810

† Electronic supplementary information (ESI) available: Experimental procedures and characterization of all new compounds. See DOI: 10.1039/c2cc32757k


**Table 1** Enantioselective Friedel–Crafts arylation of **1**<sup>a</sup>





						
Entry	Pyrrole or indole	R <sup>1</sup>	R <sup>2</sup>	Product	Time/h	Yield <sup>b</sup> (%)
1		H	H	<b>5a</b>	10	92
2 <sup>d</sup>		H	H	<b>5a</b>	4	92
3		Me	H	<b>5b</b>	12	96
4		Bn	H	<b>5c</b>	16	85
5 <sup>e</sup>		H	Me	<b>5d</b>	11	61
6		H	H	<b>6a</b>	12	98
7		Me	H	<b>6b</b>	14	83
8		H	OMe	<b>6c</b>	13	71

<sup>a</sup> All reactions were carried out using 1.5 equiv. pyrrole or indole in the presence of 10 mol% of **2**. <sup>b</sup> Isolated yield. <sup>c</sup> Determined by chiral HPLC analysis. <sup>d</sup>  $\alpha,\alpha$ -Diphenyl-2-pyrrolidinemethanol trimethylsilyl ether was used instead of **2**. <sup>e</sup> The product was obtained as a mixture of 2,4- and 2,3-disubstituted forms.

With the method for the practical synthesis of **1** in hand, we attempted the enantioselective transformation of **1**. First, organocatalytic Friedel–Crafts arylation of **1** was carried out with pyrroles and indoles as the nucleophiles.<sup>9</sup> The reaction of an unprotected pyrrole with **1** in the presence of 10 mol% **2** afforded the corresponding 2-substituted pyrrole **5a** in 96% ee (Table 1, entry 1).<sup>10</sup> High enantioselectivity was also observed when *N*-substituted pyrroles were used (97% and 99% ee, entries 3 and 4). The reaction of 3-methylpyrrole with **1** afforded the corresponding 2,4-disubstituted product **5d** in 61% yield with good enantioselectivity, along with a 2,3-disubstituted product in 25% yield (entry 5). When indoles were employed, the corresponding 3-substituted indoles **6** were obtained with high enantioselectivities, irrespective of the substituent on the indole ring (entries 6–8). We also examined the use of other electron-rich aromatic compounds. The reaction with 3-methoxy-*N,N*-dimethylaniline<sup>11</sup> afforded the desired compound **7** with 99% ee. On the other hand, the reaction with 2-methoxyfuran afforded the corresponding product **8** with 70% ee (Scheme 3).

We next attempted the enantioselective Michael addition of **1** with heteroatom nucleophiles. As per the reported procedure,<sup>12</sup>

**Scheme 3** Enantioselective Friedel–Crafts arylation of **1** with electron-rich aromatic compounds.**Table 2** Enantioselective Michael addition of heteroatom nucleophiles<sup>a</sup>


						
Entry	Nucleophile	Product	Yield <sup>b</sup> (%)	ee <sup>c</sup> (%)		
1	RSH		<b>9a</b> : R = Ph	97	94	
2 <sup>d</sup>			<b>9a</b>	92	90	
3			<b>9b</b> : R = CH <sub>2</sub> Ph	98	95	
4			<b>9c</b> : R = <i>c</i> -Hex	99	96	
5			<b>9d</b> : R = <i>n</i> -Bu	93	93	
6			<b>9e</b> : R = CH <sub>2</sub> CO <sub>2</sub> Et	87	92	
7 <sup>e</sup>	1,2,4-Triazole		<b>10a</b> : X = N	77	86	
8 <sup>e</sup>	1,2,3-Triazole		<b>10b</b> : X = CH	74	83	
			Y = N			
9 <sup>f,g</sup>	PhCH=N–OH		<b>11</b>	73	95	

<sup>a</sup> All reactions were carried out using 1.5 equiv. of **1** (based on the nucleophile) in the presence of 10 mol% **2** and benzoic acid, unless otherwise noted. <sup>b</sup> Isolated yield. <sup>c</sup> Determined by chiral HPLC analysis. <sup>d</sup>  $\alpha,\alpha$ -Diphenyl-2-pyrrolidinemethanol trimethylsilyl ether was used instead of **2**. <sup>e</sup> Reaction was carried out at –20 °C for 40 h with 20 mol% of **2** and benzoic acid. <sup>f</sup> *ent*-**2** was used instead of **2**. <sup>g</sup> Reaction was carried out using 3 equiv. of aldoxime (based on **1**).

**1** was allowed to react with several nucleophiles in the presence of 10–20 mol% **2** and benzoic acid (Table 2). The reaction of **1** with alkylthiols proceeded smoothly to yield the corresponding adducts, which were subsequently reduced to the primary alcohols **9** in good yields. The optical purity of **9** was found to be 92–96% ee (entries 1, 3–6). The reactions of **1** with triazoles also afforded the corresponding adducts with good enantioselectivities (entries 7 and 8). Unfortunately, the reactions with succinimide and *N*-methoxycarbamates were not productive. Furthermore, the oxa-Michael reaction of **1** was carried out using benzaldehyde oxime as the nucleophile (entry 9). The reaction yielded the corresponding  $\beta$ -oxime ether with good conversion, which was isolated after reduction to **11** (73% yield, 95% ee).

Finally, we demonstrated the asymmetric synthesis of a reversible monoamine oxidase A (MAO-A) inhibitor, bexlofloxone,<sup>13</sup> using a literature procedure (Scheme 4).<sup>13e</sup> First, 4,4,4-trifluorobutane-1,3-diol **12** was synthesized by palladium-catalyzed hydrogenolysis of **11**.<sup>14,15</sup> Selective tosylation of the primary alcohol in **12** afforded tosylate **13** in 53% yield. Next, oxazolidinone **17** was synthesized. Carbamate **14** was prepared by the treatment of 4-benzyloxylaniline with methyl chloroformate. Subsequent condensation with commercially available (*S*)-4-methoxymethyl-1,3-dioxolane-2-one (**15**) in the presence of K<sub>2</sub>CO<sub>3</sub> afforded **16** in 83% yield.<sup>16</sup> Then, deprotection of **16** by palladium-catalyzed hydrogenolysis yielded oxazolidinone **17**. Finally, etherification of **17** with **13** yielded bexlofloxone in 84% yield with 95% de, as shown in Scheme 4.

In conclusion, we developed a practical synthetic route to 4,4,4-trifluorocrotonaldehyde (**1**), which is a versatile precursor for organocatalytic enantioselective 1,4-additions. Reactions with



**Scheme 4** Asymmetric synthesis of befloxtone. Reaction conditions: (a) Pd(OH)<sub>2</sub> (20 mol% Pd), MeOH, H<sub>2</sub>, rt, 3 h; (b) TsCl (1.5 equiv.), pyridine, rt, 24 h; (c) K<sub>2</sub>CO<sub>3</sub> (3 equiv.), DMF, reflux, 3 h; (d) Pd/C (10 mol% Pd), MeOH, H<sub>2</sub> (1 atm), rt, 3 h; (e) K<sub>2</sub>CO<sub>3</sub> (2 equiv.), DMF, 85 °C, 6 h.

various nucleophiles such as alkylthiols, pyrroles, indoles, triazoles, and aldoximes afforded a variety of trifluoromethylated chiral stereogenic centers in a highly enantioselective manner. We also demonstrated the application of this method to the asymmetric synthesis of an MAO-A inhibitor, befloxtone.

This study was supported by the Takeda Science Foundation and a Grant-in-Aid for Young Scientists (B) (23750111) from MEXT. The partial support from the Asahi Glass Foundation is also acknowledged.

## Notes and references

- (a) *Fluorine in medicinal chemistry and chemical biology*, ed. I. Ojima, Wiley & Sons, New York, 2009; (b) J.-P. Bégue and D. Bonnet-Delphon, *Bioorganic and Medicinal Chemistry of Fluorine*, Wiley & Sons, Hoboken, 2008.
- For reviews, see: (a) G. Valero, X. Companyó and R. Rios, *Chem.-Eur. J.*, 2011, **17**, 2018; (b) J.-A. Ma and D. Cahard, *Chem. Rev.*, 2008, **108**, PR1; (c) N. Shibata, S. Mizuta and H. Kawai, *Tetrahedron: Asymmetry*, 2008, **19**, 2633; (d) T. Billard and B. R. Langlois, *Eur. J. Org. Chem.*, 2007, 891.
- For reviews, see: (a) J. Nie, H.-C. Guo, D. Cahard and J.-A. Ma, *Chem. Rev.*, 2011, **111**, 455; (b) K. Mikami, Y. Itoh and M. Yamanaka, *Chem. Rev.*, 2004, **104**, 1.
- For the catalytic asymmetric reaction with 4,4,4-trifluorocrotonates, see: (a) K. Shibatomi, F. Kobayashi, A. Narayama, I. Fujisawa and S. Iwasa, *Chem. Commun.*, 2012, **48**, 413; (b) X.-Q. Dong, X. Fang and C.-J. Wang, *Org. Lett.*, 2011, **13**, 4426; (c) Q.-H. Li, M.-C. Tong, J. Li, H.-Y. Tao and C.-J. Wang, *Chem. Commun.*, 2011, **47**, 11110; For the reactions with 4,4,4-trifluorocrotonoyl amide or oxazolidinone, see: (d) Y. Huang, E. Tokunaga, S. Suzuki, N. Shibata and M. Shiro, *Org. Lett.*, 2010, **12**, 1136; (e) T. Konno, T. Tanaka, T. Miyabe, A. Morigaki and T. Ishihara, *Tetrahedron Lett.*, 2008, **49**, 2106; For related building block approach with trifluoroethylidene malonates, see: (f) L. Wen, Q. Shen, X. Wan and L. Lu, *J. Org. Chem.*, 2011, **76**, 2282; (g) L. Wen, Q. Shen and L. Lu, *Org. Lett.*, 2010, **12**, 4655.
- There are four reports describing the synthesis of **1**. All of them mentioned the difficulty involved in the isolation of **1**: (a) D. M. Pearson, N. R. Conley and R. M. Waymouth, *Organometallics*, 2011, **30**, 1445; (b) L. Dumitrescu, D. T. M. Huong, N. V. Hung, B. Crousse and D. Bonnet-Delphon, *Eur. J. Med. Chem.*, 2010, **45**, 3213; (c) H. Plenkiewicz, W. Dmowski and M. Lipinski, *J. Fluorine Chem.*, 2001, **111**, 227; (d) E. T. McBee, O. R. Pierce and D. D. Smith, *J. Am. Chem. Soc.*, 1954, **76**, 3725.
- For recent reviews, see: (a) M. Nielsen, D. Worgull, T. Zweifel, B. Gschwend, S. Bertelsen and K. A. Jørgensen, *Chem. Commun.*, 2011, **47**, 632; (b) A. Córdova, *Catalytic Asymmetric Conjugate Reactions*, Wiley-VCH, Weinheim, 2010; (c) S. Bertelsen and K. A. Jørgensen, *Chem. Soc. Rev.*, 2009, **38**, 2178.
- For early reports, see: (a) M. Marigo, T. C. Wabnitz, D. Fielenbach and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2005, **44**, 794; (b) Y. Hayashi, H. Gotoh, T. Hayashi and M. Shoji, *Angew. Chem., Int. Ed.*, 2005, **44**, 4212.
- N. Ishikawa, M. G. Koh, T. Kitazume and S. K. Choi, *J. Fluorine Chem.*, 1984, **24**, 419.
- For early reports on the enantioselective Friedel-Crafts alkylation of pyrroles and indoles with MacMillan catalyst, see: (a) J. F. Austin and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2002, **124**, 1172; (b) N. A. Paras and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2001, **123**, 4370; For the report with Jørgensen-Hayashi catalyst, see: (c) L. Hong, L. Wang, C. Chen, B. Zhang and R. Wang, *Adv. Synth. Catal.*, 2009, **351**, 772.
- Recently, Soloshonok revealed that compounds containing a trifluoromethyl group directly bonded to a stereogenic carbon center are prone to an enantiomer self-disproportionation effect on achiral silica gel. We therefore confirmed that the optical purity of selected products, **6a**, **9b** and **11**, do not change even after chromatographic purification using achiral silica gel. For the self-disproportionation effect of the enantiomers, see: (a) H. Ueki, M. Yasumoto and V. A. Soloshonok, *Tetrahedron: Asymmetry*, 2010, **21**, 1396; (b) V. A. Soloshonok, H. Ueki, M. Yasumoto, S. Mekala, J. S. Hirschi and D. A. Singleton, *J. Am. Chem. Soc.*, 2007, **129**, 12112; (c) V. A. Soloshonok and D. O. Berbasov, *J. Fluorine Chem.*, 2006, **127**, 597; (d) V. A. Soloshonok, *Angew. Chem., Int. Ed.*, 2006, **45**, 766.
- N. A. Paras and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2002, **124**, 7894.
- For the addition of alkylthiols, see: (a) S. Brandau, E. Maerten and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2006, **128**, 14986; (b) M. Marigo, T. Schulte, J. Franzén and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2005, **127**, 15710; For the addition of triazoles, see: (c) P. Dinér, M. Nielsen, M. Marigo and K. Anker Jørgensen, *Angew. Chem., Int. Ed.*, 2007, **46**, 1983; For the addition of succinimide and *N*-methoxycarbamates, see: (d) H. Jiang, J. B. Nielsen, M. Nielsen and K. A. Jørgensen, *Chem.-Eur. J.*, 2007, **13**, 9068; (e) J. Vesely, I. Ibrahim, R. Rios, G.-L. Zhao, Y. Xu and A. Córdova, *Tetrahedron Lett.*, 2007, **48**, 2193; (f) For the addition of aldoximes, see: (g) S. Bertelsen, P. Dinér, R. L. Johansen and K. A. Jørgensen, *J. Am. Chem. Soc.*, 2007, **129**, 1536.
- For selected papers on the pharmacology of befloxtone or [<sup>11</sup>C]befloxtone, see: (a) M. Bottlaender, H. Valette, J. Delforge, W. Saba, I. Guenther, O. Curet, P. George, F. Dollé and M.-C. Grégoire, *J. Cereb. Blood Flow Metab.*, 2010, **30**, 792; (b) C. Leroy, V. Bragulat, I. Berlin, M.-C. Grégoire, M. Bottlaender, D. Roumenov, F. Dollé, S. Bourgeois, J. Penttilä, E. Artiges, J.-L. Martinot and C. Trichard, *J. Clin. Psychopharmacol.*, 2009, **29**, 86; (c) F. Dolle, H. Valette, Y. Bramoulle, I. Guenther, C. Fuseau, C. Coulon, C. Lartizien, S. Jegham, P. George, O. Curet, J.-L. Pinquier and M. Bottlaender, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 1771; (d) J. Wouters, F. Moureau, G. Evrard, J. J. Koenig, S. Jegham, P. George and F. Durant, *Bioorg. Med. Chem.*, 1999, **7**, 1683; (e) X. Rabasseda, L. A. Sorbera and J. Castañer, *Drugs Future*, 1999, **24**, 1057; (f) N. Haddjeri, C. De Montigny, O. Curet and P. Blier, *Eur. J. Pharmacol.*, 1998, **343**, 179; (g) O. Curet, G. Damoiseau-Ovens, C. Sauvage, N. Sontag, P. Avenet, H. Depoortere, D. Caille, O. Bergis and B. Scatton, *J. Affective Disord.*, 1998, **51**, 287; (h) O. Curet, G. Damoiseau, N. Aubin, N. Sontag, V. Rovei and F.-X. Jarreau, *J. Pharmacol. Exp. Ther.*, 1996, **277**, 253; (i) V. Rovei, D. Caille, O. Curet, D. Ego and F.-X. Jarreau, *J. Neural. Transm. Suppl.*, 1994, **41**, 339.
- For earlier reports describing the synthesis of optically active **12**, see: (a) T. Yamazaki, N. Okamura and T. Kitazume, *Tetrahedron: Asymmetry*, 1990, **1**, 521; (b) T. Kitazume, J. T. Lin and T. Yamazaki, *Tetrahedron: Asymmetry*, 1991, **2**, 235; (c) See also ref. 13e.
- The absolute configuration of **12** was determined to be *R* by comparing the optical rotation with a value described in ref. 14a.
- S. Jegham, A. Nedelec, Ph. Burnier, Y. Guminski, F. Puech, J. J. Koenig and P. George, *Tetrahedron Lett.*, 1998, **39**, 4453.