

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/23308711>

Amine-Borane σ -Complexes of Rhodium. Relevance to the Catalytic Dehydrogenation of Amine-Boranes

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · NOVEMBER 2008

Impact Factor: 12.11 · DOI: 10.1021/ja806582n · Source: PubMed

CITATIONS

97

READS

22

3 AUTHORS, INCLUDING:



Andrew Weller

University of Oxford

177 PUBLICATIONS 3,224 CITATIONS

SEE PROFILE

Amine-Borane σ -Complexes of Rhodium. Relevance to the Catalytic Dehydrogenation of Amine-Boranes


Thomas M. Douglas, Adrian B. Chaplin, and Andrew S. Weller*

Department of Chemistry, Inorganic Chemistry Laboratories, University of Oxford, U.K., OX1 3QR

Received August 19, 2008; E-mail: andrew.weller@chem.ox.ac.uk

Chemical hydrogen storage in amine-boranes (e.g., $\text{H}_3\text{B}\cdot\text{NH}_3$, 19.6 wt% H) is a possible solution to the transport of hydrogen for future energy requirements due to their high hydrogen content.¹ Although solution and solid-state dehydrogenations have been reported, there is much interest in transition-metal-catalyzed dehydrogenation or hydrolytic² reactions due to favorable kinetics and lower reaction temperatures. Catalysts for dehydrogenation include Cp_2Ti derivatives,^{3,4} Re-nitrosyls,⁵ Ir-pincer complexes,^{6,7} and Ni-NHC complexes;⁸ colloidal-Rh has also been shown to be an active catalyst,^{9,10} for which in situ EXAFS suggests that the active species might actually be smaller Rh_4 and Rh_6 “clusters”.¹¹

Scheme 1. Amine-Borane Dehydrogenation



Scheme 1 illustrates the accepted reaction course for homogeneous systems. Computational studies indicate a number of mechanistic scenarios for the dehydrogenation step: NH proton transfer to a coordinated ligand followed by transfer to the metal (Ni-NHC),¹² intermolecular stepwise transfer of NH then BH (Cp_2Ti -derivatives),¹³ and concerted NH/BH activation at the metal center (Ir-pincer complexes).¹⁴ Oxidative addition of the BH bond followed by NH β -elimination has also been suggested.⁸ All routes implicate σ -complexes of amine-borane in the reaction, and while details of intermediate species remain scarce,^{4,7} materials that represent catalyst deactivation products have been isolated.^{4,6,7} We report here η^2 -amine-borane σ -complexes of rhodium that are models for such intermediate complexes, and also catalysts for the dehydrogenation of $\text{H}_3\text{B}\cdot\text{NHMe}_2$ (DMAB), a close analogue of $\text{H}_3\text{B}\cdot\text{NH}_3$. Borane σ -complexes have been reported previously¹⁵ as have σ -amine-borane complexes,¹⁶ but as far as we are aware, there is only a brief report of such species' involvement in catalytic dehydrogenation.¹⁷ No examples involving Rh have been reported.



Scheme 2. Synthesis of New Amine-Borane σ -Complexes

Figure 1. Cationic portion of **2-Me** (left) and **3-H** (right). Selected distances (Å) and angles (deg): (**2-Me**): Rh1–B1, 2.180(4); P1–Rh1–P2, 97.35(4). (**3-H**): Rh1–B1, 2.318(8); P1–Rh1–P2, 163.65(7).

Addition of DMAB (2 equiv) to $[\text{Rh}(\text{P}^t\text{Bu}_3)_2][\text{BAR}^{\text{F}_4}]$ **1**¹⁸ ($\text{Ar}^{\text{F}_4} = 3,5\text{-C}_6\text{H}_3(\text{CF}_3)_2$) in 1,2-difluorobenzene results in the immediate formation of a purple Rh(I) species $[\text{Rh}(\text{P}^t\text{Bu}_3)_2(\eta^2\text{-H}_3\text{B}\cdot\text{NHMe}_2)]\text{-}[\text{BAR}^{\text{F}_4}]$ **2-H**. **2-H** is short-lived ($t_{1/2} \sim 1$ min) and evolves to give yellow Rh(III) $[\text{Rh}(\text{H})_2(\text{P}^t\text{Bu}_3)_2(\eta^2\text{-H}_3\text{B}\cdot\text{NHMe}_2)]\text{-}[\text{BAR}^{\text{F}_4}]$ **3-H** and the cyclic dimer $[\text{BH}_2\text{NMe}_2]_2$.¹⁰ Addition of smaller amounts of DMAB to **1** resulted in the formation of mixtures of **2-H**, **3-H**, and **1** in varying proportions, meaning that **2-H** could not be isolated free of **3-H**. **2-H** was longer-lived under these conditions allowing for its full characterization. Both complexes have been characterized by NMR spectroscopy, ESI-MS/MS, and, for **3-H**, also in the solid state (Figure 1).

¹H NMR spectra show the coordinated borane group as a broad 3H signal, relative to ^tBu and NH groups, at δ –2.13 (**2-H**) and δ –0.77 (**3-H**), which sharpen on ¹¹B decoupling. This suggests rapid exchange of terminal and bound hydrides. Cooling **3-H** to 190 K arrests this process (δ –3.15, 2H, Rh–H–B). **2-H** was not stable in suitable low-temperature solvents (CD_2Cl_2). The two hydrido ligands in **3-H** are observed as a 2H dt, δ –17.42 [$J(\text{PH})$ 20, $J(\text{RhH})$ 17], while the NH signals appear at δ 4.67 (**2-H**) and δ 3.87 (**3-H**). ³¹P{¹H} NMR spectra indicate a Rh(I) species **2-H** δ 35.9 [$J(\text{RhP})$ 174] and a Rh(III) species **3-H** δ 22.3 [$J(\text{RhP})$ 105]. ¹¹B{¹H} NMR spectroscopy shows broad signals at δ 19.3 (**2-H**) and δ 2.23 (**3-H**), shifted significantly downfield from DMAB (δ –13.4). The solid-state structure of **3-H** shows a pseudo-octahedral Rh(III) center with *trans* phosphines, *cis* hydrides, and an η^2 - $\text{H}_3\text{B}\cdot\text{NMe}_2\text{H}$ ligand [Rh1–B1 2.318(8) Å] (Figure 1). NMR data and structural metrics indicate a Rh(III) center ligated with a σ -borane rather than an alternative Rh(V) tetrahydridoboryl structure;¹⁹ a bond-indices analysis on calculated structures confirms this (see SI). **3-H** probably forms *via* dehydrogenation of the bound DMAB in **2-H** to give $[\text{Rh}(\text{H})_2(\text{P}^t\text{Bu}_3)_2][\text{BAR}^{\text{F}_4}]$ **4**,¹⁸ which combines with a further equivalent of DMAB. Consistent with this, **3-H** can be formed by addition of DMAB to **4**. **3-H** slowly loses H_2 under vacuum to reform (unstable) **2-H**, establishing a plausible dehydrogenation cycle for DMAB mediated by **1**.

As complex **2-H** is short-lived and undergoes dehydrogenation to give **3-H** by NH/BH scission, blocking this route should afford




Figure 2. DMAB dehydrogenation by **1** (5 mol%, C₆H₄F₂) in a sealed NMR tube. Inset: ¹¹B NMR spectrum after 90 min.

a stable complex. This is the case, with H₃B·NMe₃ (TMAB) affording a stable (under Ar) analogue **2-Me**. The resulting complex **2-Me** has a solid-state structure that shows a coordinated TMAB ligand with a pseudo square-planar Rh(I) center (Figure 1) and is structurally similar to related hydridoborate complexes of Rh(I).²⁰ **3-Me** can be prepared by adding H₂ to **2-Me** or addition of TMAB to **4**. Spectroscopic and ESI-MS/MS data are in full accord with these structures and are also similar to **2-H/3-H** underscoring their own structural assignment. Interestingly **3-Me** loses H₂ much more rapidly than **3-H** (simply by flushing with Ar), and we speculate that this is a steric effect arising from the additional *N*-methyl group, forcing the Rh center to adopt a less crowded Rh(I) square-plane configuration.

Complexes **1**, **4**, and **3-H** are active catalysts for the dehydrogenation of DMAB. In an open system under Ar a modest^{4,7,8} overall turnover frequency (34 h⁻¹, 298 K, 5 mol%, 100% conversion) is achieved to ultimately afford the cyclic dimer [H₂BNMe₂]₂ {δ(¹¹B) 5.4 [t, J(BH) 113]}.¹⁰ Repeating this reaction in a sealed NMR tube resulted in a lower TOF (2 h⁻¹) indicating inhibition by H₂ released during catalysis. Under these attenuated conditions a time/concentration plot (Figure 2) showed no evidence of sigmoidal kinetics. Addition of Hg did not inhibit catalysis. Both observations suggest nanoparticle formation is not occurring in catalysis.⁹ A species that shows characteristic intermediate time/concentration dependence is also observed by ¹¹B NMR spectroscopy in both the open and closed systems, δ 2.4 [t, J(BH) 112], tentatively identified as [H₂BNMe₂]₃. This species has also been identified during the dehydrogenation of DMAB by “Cp₂Ti”.³ A small amount of H₂B=NMe₂, δ 38 [d, J(BH) 123], following a similar concentration/time profile, was also observed.¹¹

Monitoring the “closed” system during catalysis by NMR spectroscopy identified a number of metal containing species, including **3-H** (ca. 20%). Other species currently elude definitive identification. At the end of catalysis only two compounds are observed in a ca. 1:1 ratio: **3-H** and another that is currently only partially characterized. ³¹P{¹H} NMR spectroscopy suggests a Rh(III) center, while ¹H NMR data indicate 2 Rh–H, 2 Rh–H–B groups and no NH. These data fit an empirical formula [Rh(H)₂–(P^tBu₃)₂(η²-H₂B=NMe₂)]⁺.¹⁵ In support of this assignment, addition of H₂B=NCy₂ to **4** results in a complex with similar NMR spectroscopic characteristics (see Supporting Information). We discount assignment as a [H₂BNMe₂]₂ adduct, as addition of this fragment¹⁰ to **4** is followed by immediate H₂ loss and the isolation of a different complex in quantitative yield: [Rh(P^tBu₃)₂(η²-(H₂BNMe₂)₂)] [BAF₄]⁻ **5** (Figure 3), a σ-complex of a cyclic amino-borane. Addition of excess DMAB to **5** or the postcatalysis mixture gives **3-H** and the resumption of catalysis. Addition of H₂ (1 atm) to **5** gives a mixture of **5**, **4**, and [H₂BNMe₂]₂.




Figure 3. Cationic portion of **5** from asymmetric cell. Selected distance (Å) and angle (deg): Rh1–B1, 2.161(6); P1–Rh1–P2, 98.31(6).

In conclusion we have isolated Rh(I) and Rh(III) σ-amine-borane complexes of H₃B·NMe₂R, and although the details of the dehydrogenation mechanism currently remain unresolved, these complexes provide useful insight into the likely intermediates. Given the isoelectronic relationship between alkane and amine-boranes, complexes **2** and **3** are also analogues of σ-alkane complexes of late-transition metals.¹⁶ **5** is thus an analogue of a transition metal bound to a cyclic alkane, complexes that have previously been observed in solution at low temperatures by NMR spectroscopy or by time-resolved IR spectroscopy.²¹

Acknowledgment. The EPSRC for support. Dr. Simon Aldridge for stimulating discussions and the gift of H₂B=NCy₂.

Supporting Information Available: Full experimental details, kinetic and characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Marder, T. B. *Angew. Chem., Int. Ed.* **2007**, *46*, 8116–8118. (b) Stephens, F. H.; Pons, V.; Baker, R. T. *Dalton Trans.* **2007**, 2613–2626.
- (2) (a) Yoon, C. W.; Sneddon, L. G. *J. Am. Chem. Soc.* **2006**, *128*, 13992–13993. (b) Yan, J.-M.; Zhang, X.-B.; Han, S.; Shioyama, H.; Xu, Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 2287–2289.
- (3) Clark, T. J.; Russell, C. A.; Manners, I. J. *Am. Chem. Soc.* **2006**, *128*, 9582–9583.
- (4) Pun, D.; Lobkovsky, E.; Chirik, P. J. *Chem. Commun.* **2007**, 3297–3299.
- (5) Jiang, Y.; Berke, H. *Chem. Commun.* **2007**, 3571–3573.
- (6) Hebden, T. J.; Denney, M. C.; Pons, V.; Piccoli, P. M. B.; Koetzle, T. F.; Schultz, A. J.; Kaminsky, W.; Goldberg, K. I.; Heinekey, D. M. *J. Am. Chem. Soc.* **2008**, *130*, 10812–10820.
- (7) Denney, M. C.; Pons, V.; Hebden, T. J.; Heinekey, D. M.; Goldberg, K. I. *J. Am. Chem. Soc.* **2006**, *128*, 12048–12049.
- (8) Keaton, R. J.; Blacquiere, J. M.; Baker, R. T. *J. Am. Chem. Soc.* **2007**, *129*, 1844–1845.
- (9) Jaska, C. A.; Manners, I. J. *Am. Chem. Soc.* **2004**, *126*, 9776–9785.
- (10) Jaska, C. A.; Temple, K.; Lough, A. J.; Manners, I. J. *Am. Chem. Soc.* **2003**, *125*, 9424–9434.
- (11) Fulton, J. L.; Linehan, J. C.; Autrey, T.; Balasubramanian, M.; Chen, Y.; Szymczak, N. K. *J. Am. Chem. Soc.* **2007**, *129*, 11936–11949.
- (12) Yang, X.; Hall, M. B. *J. Am. Chem. Soc.* **2008**, *130*, 1798–1799.
- (13) Luo, Y.; Ohno, K. *Organometallics* **2007**, *26*, 3597–3600.
- (14) Paul, A.; Musgrave, C. B. *Angew. Chem., Int. Ed.* **2007**, *46*, 8153–8156.
- (15) (a) Alcaraz, G.; Clot, E.; Helmstedt, U.; Vendier, L.; Sabo-Etienne, S. *J. Am. Chem. Soc.* **2007**, *129*, 8704–8705. (b) Crestani, M. G.; Muñoz-Hernández, M.; Arévalo, A.; Acosta-Ramírez, A.; García, J. J. *J. Am. Chem. Soc.* **2005**, *127*, 18066–18073. (c) Hartwig, J. F.; Muhoro, C. N.; He, X.; Eisenstein, O.; Bosque, R.; Maseras, F. *J. Am. Chem. Soc.* **1996**, *118*, 10936–10937. (d) Lin, Z. *Struct. Bonding (Berlin)* **2008**, *130*, 123–148.
- (16) Shimoi, M.; Nagai, S.-I.; Ichikawa, M.; Kawano, Y.; Katoh, K.; Uruichi, M.; Ogino, H. *J. Am. Chem. Soc.* **1999**, *121*, 11704–11712.
- (17) Shimoi, M.; Kawano, Y.; Taeko, T. S. 224th ACS National Meeting, Boston, 2002.
- (18) Douglas, T. M.; Chaplin, A. B.; Weller, A. S. *Organometallics* **2008**, *27*, 2918–2921.
- (19) Hartwig, J. F.; De Gala, S. R. *J. Am. Chem. Soc.* **1994**, *116*, 3661–3662.
- (20) Wescott, S. A.; Marder, T. B.; Baker, R. T.; Harlow, R. L.; Calabrese, J. C.; Lam, K. C.; Lin, Z. *Polyhedron* **2004**, *23*, 2665–2677.
- (21) Ball, G. E.; Brookes, C. M.; Cowan, A. J.; Darwish, T. A.; George, M. W.; Kawanami, H. K.; Portius, P.; Rourke, J. P. *Proc. Natl. Acad. Sci. U.S.A.* **2007**, *104*, 6927–6932, and references therein.

JA806582N