J Am Chem Soc. Author manuscript; available in PMC 2011 April 7.

Published in final edited form as:

J Am Chem Soc. 2010 April 7; 132(13): 4562-4563. doi:10.1021/ja100949e.

anti-Diastereo- and Enantioselective Carbonyl (Hydroxymethyl) allylation from the Alcohol or Aldehyde Oxidation Level: Allyl Carbonates as Allylmetal Surrogates

Yong Jian Zhang^{†,‡}, Jin Haek Yang[†], Sang Hoon Kim[†], and Michael J. Krische^{†,*}
[†]University of Texas at Austin, Department of Chemistry and Biochemistry, Austin, TX 78712

[‡]School of Chemistry and Chemical Engineering, Shanghai Jiao Tong University, Shanghai 200240, China

Abstract

Reactant Alcohols Dehydrogenate - Product Alcohols Do Not

Enantioselective transfer hydrogenation of carbonate **1a** in the presence of aromatic, allylic or aliphatic alcohols **2a–2i** employing a cyclometallated iridium *C,O*-benzoate derived from allyl acetate, 4-cyano-3-nitrobenzoic acid and (*S*)-SEGPHOS delivers products of (hydroxymethyl) allylation **4a–4i** in good isolated yields (60–74%), good *anti*-diastereoselectivities (5:1–10:1 dr) and exceptional levels of enantiocontrol (93–99% ee). Under identical reaction conditions, but in the presence of isopropanol, aldehydes **3a–3i** are converted to an equivalent set of adducts **4a–4i** in good isolated yields (58–74%), good *anti*-diastereoselectivities (4:1–14:1 dr) and exceptional levels of enantiocontrol (95–99% ee). Thus, identical sets of adducts **4a–4i** are produced with equal facility from the alcohol or aldehyde oxidation level. These studies represent the first general method for enantioselective carbonyl (hydroxymethyl)allylation, a process that has no highly stereoselective counterpart in conventional allylmetal chemistry.

The hydroxymethyl 1,3-diol motif appears in numerous natural products, 1^{-4} yet asymmetric methods for carbonyl (hydroxymethyl)allylation are largely unexplored. The most cases, catalytic carbonyl hydroxymethylation has been accomplished through umpolung of palladium π -allyl complexes derived from 2-butene-1,4-diol carboxylates or vinyl epoxides 6 in combination with metallic reductants, such as SnCl₂ or InI. However, control of regio- and diastereoselectivity has proven challenging. Nakajima, as well as Cozzi and Umani-Ronchi, each report a single example of catalytic *syn*-(hydroxymethyl)allylation but only moderate

mkrische@mail.utexas.edu.

enantioselectivities were observed. To our knowledge, corresponding protocols for enantioselective *anti*-(hydroxymethyl)allylation are unknown. 8

We have found that chiral *ortho*-cyclometallated iridium C,O-benzoates catalyze carbonyl allylation,9a,b,e—h crotylation9c,f and *tert*-prenylation9d,f employing allyl acetate, α -methyl allyl acetate and 1,1-dimethylallene as allyl donors, respectively. For such C-C bond forming transfer hydrogenations, ¹⁰ alcohols function both as hydrogen donors and carbonyl precursors, enabling identical sets of carbonyl addition products to be generated from either the alcohol or aldehyde oxidation level. ^{1,2} In more recent work, it was found that use of the isolated iridium C,O-benzoate complex was essential for efficient reductive couplings of allylic *gem*-dibenzoates. ⁹ⁱ This outcome prompted us to reexamine processes that failed using *in situ* generated catalysts, including reactions of allylic carbonates.

Here, we report that complex (*S*)-**I**, which is modified by the chiral phosphine ligand (*S*)-SEGPHOS, ¹¹ serves as a single-component catalyst for the coupling of cyclic carbonate **1a** to alcohols **2a–2i** to furnish (hydroxymethyl)allylation products **4a–4i** in highly enantiomerically enriched form. Under similar conditions in the presence of isopropanol, cyclic carbonate **1a** couples to aldehydes **3a–3i** to furnish an identical set of adducts **4a–4i** with comparable levels of selectivity. These studies represent the first general method for enantioselective carbonyl (hydroxymethyl)allylation - a process that has no highly stereoselective counterpart in conventional allylmetal chemistry.

Reactant Alcohols Dehydrogenate - Product Alcohols Do Not

A principle concern regarding use of cyclic carbonate 1a is the requirement that alcohols 2 selectively dehydrogenate in the presence of diol-containing products 4. To probe this issue and to explore the feasibility of utilizing allylic carbonates as allyl donors, cyclic carbonate 1a was exposed to benzyl alcohol 2a in the presence of the cyclometallated complex derived from $[Ir(cod)Cl]_2$, 4-cyano-3-nitrobenzoic acid, allyl acetate and BIPHEP (2,2'-bis (diphenylphosphino)biphenyl). Remarkably, decarboxylative anti-(hydroxymethyl)allylation occurs smoothly to furnish the desired diol 4a in good isolated yield. Dehydrogenation of the diol product is not observed as the homoallylic olefin of 4a binds the coordination site required for β -hydride elimination. 10d,11 Exclusive formation of the branched regioisomer and anti-diastereoselectivity are consistent with carbonyl addition from the primary (E)- σ -allyl iridium haptomer by way of a chair-like transition structure. Finally, unlike analogous reactions of allylic acetates which require added base, 9a-c, e-i the decarboxylative process occurs in the absence of base or any other additive.

This result prompted an assay of chiral iridium C,O-benzoates. Among the complexes assayed, (S)-I, which is modified by the chiral phosphine ligand (S)-SEGPHOS, ¹² was superior. By simply combining carbonate **1a** with alcohols **2a–2i** in the presence of (S)-I in THF solvent at 90 °C, products of (hydroxymethyl)allylation **4a–4i** are generated with good *anti*-diastereoselectivities (5:1-10:1 dr) and exceptional levels of enantiocontrol (93–99% ee). The isolated yields were moderate (60-74%) due to incomplete consumption of alcohols **2a–2i** (Table 1). Higher yields are obtained if reaction time is extended.

Aldehydes **3a–3i** are converted to an equivalent set of adducts **4a–4i** under similar conditions employing isopropanol as the terminal reductant. Comparable isolated yields (58–74%), *anti*-diastereoselectivities (4:1–14:1 dr) and enantioselectivities (95–99% ee) are observed (Table 2). Thus, identical adducts **4a–4i** are produced with equal facility from the alcohol or aldehyde oxidation level. Construction of oxetane **5c** in 2 steps from adduct **4c** serves to illustrate the utility of the (hydroxymethyl)allylation process. Similarly, pyrans **6c** and **7c** are prepared in 3 and 2 steps from adduct **4c**, respectively (Scheme 1).

The ability of allylic carbonate **1a** to participate in intermolecular decarboxylative C-C bond forming transfer hydrogenation prompted us to investigate the decarboxylative C-C coupling of allyl-benzyl carbonates **1b** and **1c**. Remarkably, using the achiral iridium catalyst BIPHEP-**I**, the desired products of CC bond formation **8** and **9** were produced in modest yield along with recovered benzyl alcohol. As a molar excess of allyl donor is required to enforce high conversion in the iridium catalyzed carbonyl allylations we describe, high-yielding decarboxylative C-C coupling of allyl carbonates will require improved second generation catalysts.

In summary, we report the first general method for enantioselective carbonyl (hydroxymethyl) allylation. Future studies will focus on the development of related C-C couplings of alcohols and π -unsaturated reactants.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Acknowledgment is made to the Robert A. Welch Foundation and the NIH-NIGMS (RO1-GM069445). Dr. Yasunori Ino and Dr. Wataru Kuriyama of Takasago are thanked for the generous donation of (S)-SEGPHOS. YJZ acknowledges partial financial support from Shanghai Jiao Tong University.

References

Tedanolide, 13-deoxytedanolide and the myriaporones incorporate the hydroxymethyl 1,3-diol motif. For total syntheses, see: (a) Pérez M, del Pozo C, Reyes F, Rodríguez A, Francesch A, Echavarren AM, Cuevas C. Angew. Chem., Int. Ed 2004;43:1724. (b) Taylor RE, Fleming KN. Angew. Chem., Int. Ed 2004;43:1728. (c) Julian LD, Newcom JS, Roush WR. J. Am. Chem. Soc 2005;127:6186. [PubMed: 15853320] (d) Ehrlich G, Hassfeld J, Eggert U, Kalesse M. J. Am. Chem. Soc 2006;128:14038. [PubMed: 17061881] (e) Smith AB III, Lee D. J. Am. Chem. Soc 2007;129:10957. [PubMed: 17691787] (f) Dunetz JR, Julian LD, Newcom JS, Roush WR. J. Am. Chem. Soc 2008;130:16407. [PubMed: 18980317] (g) Ehrlich G, Hassfeld J, Eggert U, Kalesse M. Chem. Eur. J 2008;14:2232.

2. Tylosin, its aglycone tylonolide, and O-mycinosyltylonolide incorporate the hydroxymethyl 1,3-diol motif. For total syntheses, see: (a) Tatsuta K, Amemiya Y, Kanemura Y, Kinoshita M. Tetrahedron Lett 1981;22:3997. (b) Tatsuta K, Amemiya Y, Kanemura Y, Takahashi H, Kinoshita M. Tetrahedron Lett 1982;23:3375. (c) Masamune S, Lu LD-L, Jackson WP, Kaiho T, Toyoda T. J. Am. Chem. Soc 1982;104:5523. (d) Grieco PA, Inanaga J, Lin N-H, Yanami T. J. Am. Chem. Soc 1982;104:5781. (e) Nicolaou KC, Seitz SP, Pavia MR. J. Am. Chem. Soc 1982;104:2030. (f) Tanaka T, Oikawa Y, Hamada T, Yonemitsu O. Tetrahedron Lett 1986;27:3651. (g) Tanaka T, Oikawa Y, Hamada T, Yonemitsu O. Chem. Pharm. Bull 1987;35:2219.

- 3. The mycinamicins incorporate the hydroxymethyl 1,3-diol motif. For total syntheses, see: (a) Suzuki K, Matsumoto T, Tsuchihashi G-i. Chem. Lett 1987:113. (b) Matsumoto T, Maeta H, Suzuki K, Tsuchihashi G-i. Tetrahedron Lett 1988;29:3575.
- 4. For other natural products that incorporate the hydroxymethyl 1,3-diol motif, see Supporting Information.
- 5. For (hydroxymethyl)allylation via palladium catalyzed reductive coupling of allylic carboxylates, see: (a) Masuyama Y, Takahara JP, Kurusu Y. J. Am. Chem. Soc 1988;110:4473. (b) Masuyama Y, Otake K, Kurusu Y. Tetrahedron Lett 1988;29:3563. (c) Takahara JP, Masuyama Y, Kurusu Y. J. Am. Chem. Soc 1992;114:2577.
- 6. For (hydroxymethyl)allylation via palladium catalyzed reductive coupling of vinyl epoxides, see: (a) Araki S, Kameda K, Tanaka J, Hirashita T, Yamamura H, Kawai M. J. Org. Chem 2001;66:7919. [PubMed: 11701061] (b) Gagliardo M, Selander N, Mehendale NC, van Koten G, Klein Gebbink RJM, Szabó KJ. Chem. Eur. J 2008;14:4800.
- For catalytic enantioselective (hydroxymethyl)allylation, see: (a) Bandini M, Cozzi PG, Licciulli S, Umani-Ronchi A. Synthesis 2004:409. (b) Nakajima M, Saito M, Hashimoto S. Chem. Pharm. Bull 2000;48:306. [PubMed: 10705529]
- For selected reviews on enantioselective carbonyl allylation and crotylation, see: (a) Hoffmann RW. Angew. Chem., Int. Ed 1982;21:555. (b) Yamamoto Y, Asao N. Chem. Rev 1993;93:2207. (c) Ramachandran PV. Aldrichim. Acta 2002;35:23. (d) Kennedy JWJ, Hall DG. Angew. Chem., Int. Ed 2003;42:4732. (e) Denmark SE, Fu J. Chem. Rev 2003;103:2763. [PubMed: 12914480] (f) Yu C-M, Youn J, Jung H-K. Bull. Korean Chem. Soc 2006;27:463. (g) Marek I, Sklute G. Chem. Commun 2007:1683. (h) Hall DG. Synlett 2007:1644.
- 9. For enantioselective carbonyl allylation, crotylation and reverse prenylation via iridium catalyzed C-C bond forming transfer hydrogenation, see: (a) Kim IS, Ngai M-Y, Krische MJ. J. Am. Chem. Soc 2008;130:6340. [PubMed: 18444616] (b) Kim IS, Ngai M-Y, Krische MJ. J. Am. Chem. Soc 2008;130:14891. [PubMed: 18841896] (c) Kim IS, Han S-B, Krische MJ. J. Am. Chem. Soc 2009;131:2514. [PubMed: 19191498] (d) Han SB, Kim I-S, Han H, Krische MJ. J. Am. Chem. Soc 2009;131:6916. [PubMed: 19453190] (e) Lu Y, Kim I-S, Hassan A, Del Valle DJ, Krische MJ. Angew. Chem., Int. Ed 2009;48:5018. (f) Itoh J, Han SB, Krische MJ. Angew. Chem., Int. Ed 2009;48:6316. (g) Lu Y, Krische MJ. Org. Lett 2009;11:3108. [PubMed: 19586066] (h) Hassan A, Lu Y, Krische MJ. Org. Lett 2009;11:3112. [PubMed: 19586067] (i) Han SB, Han H, Krische MJ. J. Am. Chem. Soc 2010;132:1760. [PubMed: 20099821] (j) For a recent application in total synthesis, see: Harsh P, O'Doherty GA. Tetrahedron 2009;65:5051. [PubMed: 20161297]
- For recent reviews on C-C bond forming transfer hydrogenation, see: (a) Patman RL, Bower JF, Kim IS, Krische MJ. Aldrichim. Acta 2008;41:95. (b) Shibahara F, Krische MJ. Chem. Lett 2008;37:1102.
 (c) Bower JF, Kim IS, Patman RL, Krische MJ. Angew. Chem. Int. Ed 2009;48:34. (d) Han SB, Kim IS, Krische MJ. Chem. Commun 2009:7278.
- 11. In response to a reviewers comment: the olefin moiety of homoallylic alcohol products **4a–4i**, appears to bind the iridium center more strongly than the olefin moiety of allylic alcohols such as cinnamyl alcohol **2f** (5-membererd versus 4-membered chelate). Thus, alcohol reactants **2a–2i** dehydrogenate whereas homoallylic alcohol products do not.
- 12. Saito T, Yokozawa T, Ishizaki T, Moroi T, Sayo N, Miura T, Kumobayashi H. Adv. Synth. Catal 2001;343:264.

Scheme 1.

Conversion of diol **4c** to compounds **5c**, **6c** and **7c**.a aReagents: (a) NaH, TsCl, THF, 82%. (b) *n*-BuLi, THF, 92%. (c) NaH, H2C=CHCH2Br, THF, 82%. (d) Grubbs I, DCM, 90%. (e) TBSCl, Et3N, DMAP, DCM, 88%. (f) NaH, H2C=CHCH2Br, THE, 90%. (c) Grabbs I, DCM, 91%. See Supporting Information for further

 $H2C = CHCH2Br, THF, 90\%. \ (g) \ Grubbs \ I, DCM, 91\%. \ See \ Supporting \ Information \ for further \ details.$

Table 1

Enantioselective (hydroxymethyl)allylation from the alcohol oxidation level.^a

66% Yield, 10:1 dr 95% ee, 4g

70% Yield, 5:1 dr

98% ee, 4d

HC

72% Yield, 9:1 dr 99% ee, 4e H₽

67% Yield, 9:1 dr 97% ee, 4h

74% Yield, 7:1 dr 93% ee, 4f

60% Yield, 9:1 dr 98% ee, 4i

^aYields are of material isolated by silica gel chromatography. Enantiomeric excess was determined by chiral stationary phase HPLC analysis. See Supporting Information for further details.

Table 2

Enantioselective (hydroxymethyl)allylation from the aldehyde oxidation level. a

3a, R = Ph	3b, $R = m$ -MeOPh 3e, $R = 2$ -Furyl 3h, $R = (CH2)2Ph$		3c, $R = p$ -BrPh 3f, $R = CH = CHPh$ 3i, $R = (CH2)7Me$
3d, $R = p-(CO_2Me)Ph$			
3g, R = Geranyl			
	HO	HO	HO
		OMe	
	63% Yield, 5:1 dr	74% Yield, 6:1 dr	HO Br 64% Yield, 5:1 dr
	99% ee, 4a HO	98% ee, 4b HO	95% ee, 4c HO
			Ph
	HO CO ₂ Me	но	но
	70% Yield, 4:1 dr 98% ee, 4d	67% Yield, 5:1 dr 97% ee, 4e	68% Yield, 7:1 dr 99% ee, 4f
	HQ Me	HQ	HQ
	Me Me	Ph	(CH ₂) ₇ Me
	HO HO	HO	HO
	63% Yield, 14:1 dr 97% ee, 4g	61% Yield, 8:1 dr 99% ee, 4h	58% Yield, 8:1 dr 97% ee, 4i

^aAs described for Table 1.