

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/255756116>

Diaza-18-crown-6 hydroxyquinoline derivatives as flexible tools for the assessment and imaging of total intracellular magnesium

ARTICLE *in* CHEMICAL SCIENCE · FEBRUARY 2012

Impact Factor: 9.21 · DOI: 10.1039/C1SC00751C

CITATIONS

8

READS

17

10 AUTHORS, INCLUDING:



Giovanna Farruggia

University of Bologna

73 PUBLICATIONS 871 CITATIONS

SEE PROFILE



Luca Prodi

University of Bologna

207 PUBLICATIONS 6,955 CITATIONS

SEE PROFILE



Federica I Wolf

Catholic University of the Sacred Heart

77 PUBLICATIONS 2,331 CITATIONS

SEE PROFILE



Stefano Iotti

University of Bologna

111 PUBLICATIONS 2,296 CITATIONS

SEE PROFILE

Supporting Information

Title

Diaza-18-crown-6 hydroxyquinoline derivatives as flexible tools for the assessment and imaging of total intracellular magnesium

Author List

Chiara Marraccini,^{1*} Giovanna Farruggia,² Marco Lombardo,³ Luca Prodi,³ Massimo Sgarzi,³ Valentina Trapani,⁴ Claudio Trombini,³ Federica I. Wolf,⁴ Nelsi Zaccheroni³ and Stefano Iotti.^{1,5**}

Institution Address

¹Università di Bologna, Dipartimento di Medicina Interna, dell’Invecchiamento e Malattie Nefrologiche, Bologna, Italy

² Università di Bologna, Dipartimento di Biochimica “G. Moruzzi”, Bologna, Italy

³ Università di Bologna, Dipartimento di Chimica “G. Ciamician”, Bologna, Italy

⁴Università Cattolica del Sacro Cuore, Istituto di Patologia Generale e Centro di Ricerche Oncologiche Giovanni XXIII, Facoltà di Medicina e Chirurgia “A. Gemelli”, Roma, Italy

⁵ Istituto Nazionale Biostrutture e Biosistemi, Roma, Italy

Corresponding Authors

* Chiara Marraccini - Email: chiara.marraccini@unibo.it

** Stefano Iotti - Email: stefano.iotti@unibo.it

1. General Information and Instrumentation

General Spectroscopic Methods. All reagents were Ultrapure grade. The fluorescent probes were synthesized as reported in ¹ and were dissolved in dimethyl sulfoxide (DMSO) to a final concentration of 1 mg/mL (DCHQ1 1.7 mM; DCHQ3 1.4 mM; DCHQ4 1.4 mM; DCHQ5 1.4 mM; DCHQ6 1.3 mM). Aliquots were kept at 4°C in the dark.

Photophysical measurements. Absorption spectra were recorded on a Perkin-Elmer Lambda 45 spectrophotometer. For the fluorescence spectroscopy measurements, uncorrected emission and corrected excitation spectra were obtained with a Perkin-Elmer LS 55 spectrofluorimeter.

Cell Culture. HL60 (Human Promyelocytic Leukemia) cells were grown at 37°C and 5% CO₂ in RPMI 1640 medium supplemented with 10% heat-inactivated fetal calf serum (FCS), 2 mM L-glutamine, 1000 units/mL penicillin and 1 mg/mL streptomycin. Differentiation of HL60 cells was induced by treatment with 1.3 % DMSO for at least 3 days, as reported in ².

Flow Cytometric Assay of DNA content in DMSO-treated HL60 cells. Control and DMSO-differentiated HL60 cells were collected following the Nüsse protocol for the assessment of total DNA content.³ Briefly, cell suspensions were washed and resuspended at a final concentration of 2×10^6 cells/mL in Nüsse-1 Solution (NaCl 0.584 g/L, trisodic citrate 1.139 g/L, DNA free RNase 10 mg/L, Nonidet P-40 0.03 % v/v).

After 30 minutes of incubation at 4°C, an equal volume of Nüsse-2 Solution was added (sucrose 85.5 g/L, citric acid 15 g/L, propidium iodide 100 mg/L). Samples were analyzed on an Epics-XL cytometer (Beckman Coulter, USA) with excitation band centered on 488 nm and emission band on 605 nm in linear scale.

2. Photophysical properties of DCHQ1




Figure 1 Titration with Mg^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 324 \text{ nm}$, panel B) spectra of DCHQ1 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 2 equivalents) of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.66 mM).




Figure 2 Titration with Zn^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 324 \text{ nm}$, panel B) spectra of DCHQ1 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 3 equivalents) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (1.89 mM).




Figure 3 Titration with Cd^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 330$ nm, panel B) spectra of DCHQ1 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 2.5 equivalents) of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ solution (1.60 mM).

3. Photophysical properties of DCHQ3




Figure 4 Titration with Mg^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 336 \text{ nm}$, panel B) spectra of DCHQ4 ($25 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 1.5 equivalents) of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 5 Titration with Zn^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 336 \text{ nm}$, panel B) spectra of DCHQ4 ($25 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 2.5 equivalents) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 6 Titration with Cd^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 336$ nm, panel B) spectra of DCHQ4 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 3 equivalents) of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ solution (1.60 mM).

4. Photophysical properties of DCHQ4




Figure 7 Titration with Mg^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 330 \text{ nm}$, panel B) spectra of DCHQ4 ($25 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 1.5 equivalents) of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 8 Titration with Zn^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 330 \text{ nm}$, panel B) spectra of DCHQ4 ($25 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 2 equivalents) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 9 Titration with Cd^{2+} . Absorption (panel A) and emission ($(\lambda_{\text{exc}} = 330 \text{ nm})$, panel B) spectra of DCHQ4 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 2.5 equivalents) of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ solution (1.60 mM).

5. Photophysical properties of DCHQ5




Figure 10 Titration with Mg^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 344$ nm, panel B) spectra of DCHQ5 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 4 equivalents) of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 11 Titration with Zn^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 344$ nm, panel B) spectra of DCHQ5 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 2 equivalents) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 12 Titration with Cd^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 344$ nm, panel B) spectra of DCHQ5 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 1.5 equivalents) of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ solution (1.60 mM).

6. Photophysical properties of DCHQ6




Figure 13 Titration with Mg^{2+} . Absorption (panel A) and emission ($\lambda_{\text{ex}} = 337 \text{ nm}$, panel B) spectra of DCHQ6 ($18 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 1 equivalent) of $\text{Mg}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 14 Titration with Zn^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 337 \text{ nm}$, panel B) spectra of DCHQ6 ($25 \mu\text{M}$) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at $\text{pH}=7.4$ with MOPS at room temperature upon addition of increasing amounts (up to 2 equivalents) of $\text{Zn}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ solution (2.50 mM).




Figure 15 Titration with Cd^{2+} . Absorption (panel A) and emission ($\lambda_{\text{exc}} = 335$ nm, panel B) spectra of DCHQ6 (25 μM) in 1:1 $\text{CH}_3\text{OH}:\text{H}_2\text{O}$ mixture buffered at pH=7.4 with MOPS at room temperature upon addition of increasing amounts (up to 1.5 equivalents) of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ solution (1.60 mM).

7. Flow Cytometric Assay of DNA content of HL60 treated with DMSO




Figure 16 Assessment of the total DNA content of control (panel A) and DMSO-differentiated (panel B) HL60 cells. Hystograms show the PI fluorescence (expressed in fluorescence channels), that is directly proportional to the DNA content, with respect to the number of events acquired. Results from a typical experiment.

8. References

- ¹A. Sanz-Medel, R. Fernandez de la Campa and J. I. Garcia-Alonso. *Analyst*, 1987, 112:493–497.
- ²F. I. Wolf, V. Covacci, N. Bruzzese, A. Di Francesco, A. Sacchetti, D. Corda and A. Cittadini. *J. Cell. Biochem.*, 1998, 71:441–448.
- ³M. Nusse and J. Kramer. *Cytometry*, 1984, 5:20-5.