

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/43506531>

# Structure and Absolute Stereochemistry of the Anticancer Agent EBC-23 from the Australian Rainforest

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · OCTOBER 2008

Impact Factor: 12.11 · DOI: 10.1021/ja807133p · Source: OAI

CITATIONS

20

READS

87

## 9 AUTHORS, INCLUDING:



**Victoria A Gordon**

University of Queensland

17 PUBLICATIONS 106 CITATIONS

SEE PROFILE



**Paul Reddell**

EcoBiotics Limited

83 PUBLICATIONS 1,472 CITATIONS

SEE PROFILE



**Heiko Schill**

Agilent Technologies

24 PUBLICATIONS 284 CITATIONS

SEE PROFILE



**Craig M Williams**

University of Queensland

142 PUBLICATIONS 1,222 CITATIONS


SEE PROFILE

## Structure and Absolute Stereochemistry of the Anticancer Agent EBC-23 from the Australian Rainforest

Lin Dong,<sup>†</sup> Victoria A. Gordon,<sup>§</sup> Rebecca L. Grange,<sup>†</sup> Jenny Johns,<sup>‡</sup> Peter G. Parsons,<sup>‡</sup>  
Achim Porzelle,<sup>†</sup> Paul Reddell,<sup>§</sup> Heiko Schill,<sup>†</sup> and Craig M. Williams<sup>\*,†</sup>*School of Molecular and Microbial Sciences, University of Queensland, Brisbane, Queensland, Australia,  
Queensland Institute of Medical Research, PO Royal Brisbane Hospital, Brisbane, Queensland, Australia, and  
EcoBiotics Ltd, Yungaburra, Queensland, Australia*

Received September 15, 2008; E-mail: c.williams3@uq.edu.au

Australian tropical rainforests are megadiverse and offer an opportunity for discovery of new bioactive compounds. During a screening program to identify biologically active candidates as potential treatments for cancer, the spiroacetal EBC-23 (**1**) was found in the fruit of *Cinnamomum laubatii* (family Lauraceae).<sup>1</sup> EBC-23 (**1**) displayed noticeable in vitro anticancer activity against a range of cancer cell lines [i.e., melanoma (MM96L), breast carcinoma (MCF7), prostate cancer (DU145)]. More significantly, xenografts of a human prostate tumor cell line (DU145) in immunodeficient mice when treated with **1** inhibited tumor growth with no observable side effects, suggesting the potential for the treatment of refractory solid tumors in adults.<sup>1</sup> Critical to the development of this lead candidate would be an implicit understanding of its chemical structure; however, only software generated<sup>2</sup> bond connectivity (i.e., **1**) was reported.<sup>1</sup> It was therefore imperative to confirm the gross structure and deduce both relative and absolute stereochemistry of this unique fused  $\alpha,\beta$ -unsaturated lactone spiroacetal.



Comparison of **1** with the known small lactone natural product osumundalactone<sup>3</sup> and the synthetic *syn*-isomer<sup>4</sup> confirmed the presence of a *syn* disubstituted lactone. Utilizing NOESY and 1D NOE experiments H5 ( $\delta_H$  4.5) correlated with H13 ( $\delta_H$  4.4), which in turn correlated with H16 ( $\delta_H$  3.8). This established the spiroacetal carbon (C8) as *EE*, and the side chain hydroxyl was  $\alpha$  configured. Even though a *syn* correlation was present between H13 ( $\delta_H$  4.4) and H12 $\beta$  ( $\delta_H$  1.8) the corresponding correlation to H11 ( $\delta_H$  4.1) was ambiguous. When a more diluted sample was measured by NMR a sharp doublet ( $\delta_H$  3.05) and singlet ( $\delta_H$  2.95) appeared for the two hydroxyl groups. With this development NOE correlations between H13 ( $\delta_H$  4.4), the C11 hydroxyl hydrogen, and H16 ( $\delta_H$  3.8) defined both configurations at C11 and C16 (i.e., H16 $\beta$ ) suggesting structure **2** (Figure 1).

Total synthesis was now key to authenticating structure **2** and determining absolute stereochemistry; however, six chiral centers partly imbedded in a unique fused  $\alpha,\beta$ -unsaturated lactone consisting of an acid labile spiroacetal unit required careful synthetic planning. Based on a plausible biosynthetic pathway, the retrosynthetic analysis initiated by ring opening of the spiroacetal moiety




Figure 1. Deduced structure **2** and retrosynthetic analysis.


arrived at the polyketide **3**. The keto functionality in **3** serves as an acyl equivalent junction (i.e., **4**) potentially suited to a Tietze<sup>5</sup>–Smith<sup>6</sup> linchpin (i.e., dithiane **5**) convergence strategy (i.e., **6** and **7**). Ring closing metathesis (RCM) was employed as a feasible choice for lactone installation via epoxide **7**. Left hand fragment **6** requires significant stereochemical flexibility built into the synthetic approach in the event the relative stereochemical assignment of **2** is incorrect or structure–activity relationship studies are deployed. To achieve this goal linchpin methodology is again considered from enantiopure epoxides **8** and **9**, which would allow construction of ketone **10** in either enantiopode or desired diastereoisomer (i.e., **6**) selectively (Figure 1).

The all *R* isomer was targeted in the first instance. Linchpin coupling of the anion of TMS-dithiane (**11**) with (*R*)-pentadecene oxide,<sup>7</sup> and then (*R*)-epi-chlorohydrin afforded the disubstituted dithiane **12** in 61% yield. Although the TBS derivative (**13**) is reported to outperform **11** in linchpin reactions (i.e., affording higher yields),<sup>8</sup> *O*-TMS protection (and not TBS protection) was fortuitously found to undergo simultaneous deprotection when the keto functionality was unmasked [Hg(ClO<sub>4</sub>)<sub>2</sub>], giving hydroxyketone **14** in high yield (Scheme 1). *Syn* selective reduction,<sup>9</sup> followed by ketal protection,<sup>10</sup> proceeded smoothly in 80% yield (over 2 steps) affording the desired left-hand fragment **15** (i.e., *RRR* enantiomer of **6**). The second linchpin coupling, that is, reaction of **13** and **15** with **16**,<sup>11</sup> gave the coupled product **17** in 78% yield. Reaction of **17** with acryloyl chloride proceeded smoothly affording acrylate **18** in 82% yield. Surprisingly, RCM required considerable effort to facilitate the formation of the desired lactone (i.e., **19**). Six RCM catalysts (**A**–**E**,<sup>14a–c</sup> **20**<sup>14f</sup>) were extensively investigated, but only the Hoveyda–Grubbs second generation catalyst (i.e., **20**) under microwave irradiation afforded lactone **19** in substantial yield (65%). Even though RCM reactions have on occasion been performed in the presence of a dithiane ring system,<sup>15</sup> the

<sup>†</sup> University of Queensland.


<sup>§</sup> EcoBiotics Ltd.

<sup>‡</sup> PO Royal Brisbane Hospital.



Comparison of the optical rotation of both the synthetic material **22** (−16.6) and the natural material (+16.6) confirmed the non-natural enantiomer had been synthesized. The synthesis was then repeated starting from (*S*)-pentadecene oxide (Scheme 2), which afforded synthetic material (+16.3) matching (<sup>1</sup>H and <sup>13</sup>C NMR are identical) both the relative and absolute stereochemistry of natural EBC-23 (**2**) (Figure 1).

**Scheme 2.** Total Synthesis of EBC-23 (**2**)



**Acknowledgment.** We thank EcoBiotech Ltd and the University of Queensland for financial support; Prof. W. Kitching for constructive discussions; Dr. L. Lambert for assistance with NMR experiments; Dr. A. Savchenko for attempts to crystallize EBC-23.

**Supporting Information Available:** Full experimental and characterization data,  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra, complete ref 4, and structures of RCM catalysts **A–E**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Reddell, P. W.; Gordon, V. A. WO 200707984 A1 20070628 PCT Int. Appl. 2007.
- (2) ACD/Structure Elucidator, Advanced Chemistry Development, Inc., Toronto ON, Canada.
- (3) Hashimoto, T.; Arakawa, T.; Tanaka, M.; Asakawa, Y. *Heterocycles* **2002**, 56, 581–588.
- (4) Ley, S. V.; et al. *J. Chem. Soc., Perkin Trans. 1* **1991**, 667–692.
- (5) Tietze, L. F.; Geissler, H.; Gewert, J. A.; Jakobi, U. *Synlett* **1994**, 511–512.
- (6) Smith, A. B., III; Adams, C. M. *Acc. Chem. Res.* **2004**, 37, 365–377.
- (7) (R)- and (S)-Pentadecene oxide were obtained from (R)- and (S)-epichlorohydrin using the method of Lepoittevin; see: Choukchou-Braham, N.; Asakawa, Y.; Lepoittevin, J.-P. *Tetrahedron Lett.* **1994**, 35, 3949–3952.
- (8) Smith, A. B., III; Pitram, S. M.; Boldi, A. M.; Gaunt, M. J.; Sfougataakis, C.; Moser, W. H. *J. Am. Chem. Soc.* **2003**, 125, 14435–14445.
- (9) Chen, K.-M.; Hardtmann, G. E.; Prasad, K.; Repic, O.; Shapiro, M. J. *Tetrahedron Lett.* **1987**, 28, 155–158.
- (10) Lipshutz, B. H.; Kozlowski, J. A. *J. Org. Chem.* **1984**, 49, 1149–1151.
- (11) Acquisition of **16** was achieved through direct transformation of a known epoxide,<sup>12</sup> via Mitsunobu inversion,<sup>13</sup> subsequent hydrolysis, and TBS protection.
- (12) For example: Schreiber, S. L.; Schreiber, T. S.; Smith, D. B. *J. Am. Chem. Soc.* **1987**, 109, 1525–1529.
- (13) Albert, B. J.; Sivaramakrishnan, A.; Naka, T.; Czaicki, N. L.; Koide, K. *J. Am. Chem. Soc.* **2007**, 127, 2648–2659.
- (14) (a) Trnka, T. M.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, 34, 18–29. (b) Scholl, M.; Ding, S.; Woo Lee, C.; Grubbs, R. H. *Org. Lett.* **1999**, 1, 953–956. (c) Kingsbury, J. S.; Harrity, J. P. A.; Bonitatebus, P. J. J.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1999**, 121, 791–99. (d) Jafarpour, L.; Schanz, H.-J.; Stevens, E. D.; Nolan, S. P. *Organometallics* **1999**, 18, 5416–5419. (e) van der Schaaf, P. A.; Hafner, A.; Mühlebach, A. US 640719 02002. (f) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, 122, 8168–8179.
- (15) For example, see: (a) Fürstner, A.; Fasching, B.; O’Neil, G. W.; Fenster, M. D. B.; Goudbot, C.; Cecon, J. *Chem. Commun.* **2007**, 304, 5–3047. (b) Nicolaou, K. C.; Kofitis, T. V.; Vyskocil, S.; Petrovic, G.; Ling, T.; Yamada, Y. M. A.; Tang, W.; Frederich, M. O. *Angew. Chem., Int. Ed.* **2004**, 43, 4318–4324. (c) Garbaccio, R. M.; Stachel, S. J.; Baeschlin, D. K.; Danishefsky, S. J. *J. Am. Chem. Soc.* **2001**, 123, 10903–10908.
- (16) Smulik, J. A.; Giessert, A. J.; Diver, S. T. *Tetrahedron Lett.* **2002**, 43, 209–211.
- (17) Aho, J. E.; Pihko, P. M.; Rissa, T. K. *Chem. Rev.* **2005**, 105, 4406–4440.
- (18) Terauchi, T.; Terauchi, T.; Sato, I.; Tsukada, T.; Kanoh, N.; Nakata, M. *Tetrahedron Lett.* **2000**, 41, 2649–2653.
- (19) Ho, T.-L.; Ho, H. C.; Wong, C. M. *J. Chem. Soc., Chem. Commun.* **1972**, 791.
- (20) (a) Peuchmaur, M.; Wong, Y.-S. *Synlett* **2007**, 2902–2906. (b) Jacobs, M. F.; Glenn, M. P.; McGrath, M. J.; Zhang, H.; Brereton, I.; Kitching, W. *ARKIVOC [online computer file]* **2001**, 2 (vii), 114–137.
- (21) Smith, A. B., III; Doughty, V. A.; Lin, Q.; Zhuang, L.; McBriar, M. D.; Boldi, A. M.; Moser, W. H.; Murase, N.; Nakayama, K.; Sobukawa, M. *Angew. Chem., Int. Ed.* **2001**, 40, 191–195.
- (22) (a) Evans, D. A.; Trotter, B. W.; Coleman, P. J.; Côté, B.; Dias, L. C.; Rajapakse, H. A.; Tyler, A. N. *Tetrahedron* **1999**, 55, 8671–8726. (b) Evans, D. A.; Coleman, P. J.; Diaz, L. C. *Angew. Chem., Int. Ed. Engl.* **1997**, 36, 2738–2741.

JA807133P