

Erratum: On non-RRKM unimolecular kinetics: Molecules in general, and CH3NC in particular

William L. Hase

Citation: The Journal of Chemical Physics 69, 4711 (1978); doi: 10.1063/1.436427

View online: http://dx.doi.org/10.1063/1.436427

View Table of Contents: http://scitation.aip.org/content/aip/journal/jcp/69/10?ver=pdfcov

Published by the AIP Publishing

Articles you may be interested in

Publisher's Note: "Coriolis coupling as a source of non-RRKM effects in ozone molecule: Lifetime statistics of vibrationally excited ozone molecules" [J. Chem. Phys.132, 224305 (2010)]

J. Chem. Phys. 137, 159902 (2012); 10.1063/1.4759494

Direct dynamics simulation of dioxetane formation and decomposition via the singlet ·O–O–CH2–CH2· biradical: Non-RRKM dynamics

J. Chem. Phys. 137, 044305 (2012); 10.1063/1.4736843

Coriolis coupling as a source of non-RRKM effects in ozone molecule: Lifetime statistics of vibrationally excited ozone molecules

J. Chem. Phys. 132, 224305 (2010); 10.1063/1.3430514

On non-RRKM unimolecular kinetics: Molecules in general, and CH3NC in particular

J. Chem. Phys. 59, 4621 (1973); 10.1063/1.1680672

RRKM Theory, the CH3NC Paradox, and the Decomposition of Hot-Atom Substitution Products

J. Chem. Phys. 57, 332 (1972); 10.1063/1.1677967






FIG. 3. The zero-field reduced mobility of CO_3^* ions in oxygen gas and of the ion X^* , whose mass is 51 amu and whose identity is not positively known. The data are presented as a function of the gas temperature.

the measurements was the determination of the gas temperature. The drift tube was operated under continuous gas flow. The temperature was controlled by heating the drift tube at three places along its length, the tube having been wrapped with several layers of $\frac{1}{2}$ in. thick

insulation. The incoming gas line was wrapped several times around the drift tube, inside the insulation, and the flow rate was low enough to permit the gas to reach approximate thermal equilibrium before entering the drift tube. The temperature was measured by a small (No. 30 gauge wire) thermocouple suspended in the gas, but outside the cylindrical electrodes establishing the electric field. We believe that our accuracy in measuring the temperature is $\pm 0.5\%$ at temperatures near 300 °K, but perhaps is no better than $\pm 2\%$ at 470 °K.

Additional uncertainty resulted from weak ion signals at particular temperatures. We believe that our total experimental error does not exceed $\pm 2\%$ at temperatures below 400 °K and $\pm 3\%$ for higher temperatures. Our value of K_0 for CO_3 in O_2 at 297 °K agrees with the zero-field value of Elford and Rees³ at 293 °K to within 1%.

³M. T. Elford and J. A. Rees, Aust. J. Phys. 27, 333 (1974).

ERRATA

Erratum: On non-RRKM unimolecular kinetics: Molecules in general, and CH₃NC in particular [J. Chem. Phys. 59, 4621 (1973)]

William L. Hase

Department of Chemistry, Wayne State University, Detroit, Michigan 48202

Equation (A14) is printed incorrectly. The correct expression is

$$x_{n-1} = \left(1 - \sum_{j=2}^{n-2} x_j^2\right)^{1/2} \sin(\frac{1}{2}\pi R).$$

The above expression was used in the calculations reported in this paper.

a) Research supported by the Atmospheric Sciences Division of the National Science Foundation through Research Grant No. ATM 76-03742.

b) Present address: Eckerd College, St. Petersburg, FL 33733.
¹H. W. Ellis, R. Y. Pai, I. R. Gatland, E. W. McDaniel, R. Wernlund, and M. J. Cohen, J. Chem. Phys. 64, 3935 (1976).
²E. W. McDaniel and E. A. Mason, The Mobility and Diffusion of Ions in Gases (Wiley, New York, 1973).