

The structure of C₄ as studied by the Coulomb explosion method

M. Algranati, H. Feldman, D. Kella, E. Malkin, E. Miklazky, R. Naaman, Z. Vager, and J. Zafman

Citation: *The Journal of Chemical Physics* **90**, 4617 (1989); doi: 10.1063/1.456597

View online: <http://dx.doi.org/10.1063/1.456597>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/90/8?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

Structures of N₂Ar, O₂Ar, and O₂Xe dimers studied by Coulomb explosion imaging

J. Chem. Phys. **137**, 104308 (2012); 10.1063/1.4750980

Nucleosynthesis driven by Coulomb explosion of nanodroplets

Phys. Plasmas **14**, 123102 (2007); 10.1063/1.2815792

Molecular structure by Coulomb explosion imaging of stored molecular ions

AIP Conf. Proc. **457**, 185 (1999); 10.1063/1.57458

Theoretical study of the Coulombic explosion in doubly charged Xe clusters

J. Chem. Phys. **107**, 9894 (1997); 10.1063/1.475287

The structures of C₂H⁺ and C₂H⁺ 2 as measured by Coulomb explosion imaging

J. Chem. Phys. **94**, 6377 (1991); 10.1063/1.460316



The Letters to the Editor section is subdivided into four categories entitled Communications, Notes, Comments, and Errata. The textual material of each Letter is limited to 1200 words minus the following: (a) 200 words for a square figure one-column wide. Larger figures are scaled in proportion to their area; (b) 50 words for each displayed equation; (c) 7 words for each line of table including headings and horizontal rulings. Proof will be sent to authors. See the issue of 1 January 1989 for a fuller description of Letters to the Editor.

COMMUNICATIONS

The structure of C_4 as studied by the Coulomb explosion method

M. Algranati, H. Feldman, D. Kella, E. Malkin, and E. Miklazky

Department of Nuclear Physics, Weizmann Institute of Science, Rehovot, 76100 Israel

R. Naaman

Department of Isotope Research, Weizmann Institute of Science, Rehovot, 76100, Israel

Z. Vager and J. Zajfman

Department of Nuclear Physics, Weizmann Institute of Science, Rehovot, 76100, Israel

(Received 13 January 1989; accepted 9 February 1989)

The structure of carbon clusters in general and C_4 in particular has been investigated recently both theoretically¹ and experimentally.^{2,3} Although a number of high level calculations indicated that even clusters that contain only four atoms have a stable cyclic (rhombic) form, experimentally it was found that only when the number of atoms exceeds nine can a cyclic form be detected. In the present work the Coulomb explosion imaging (CEI) method was used to probe the structure of C_4 . In the results presented here one observes for the first time a rhombic C_4 .

Yang *et al.*³ applied the laser sputtering source to study the photoelectron spectroscopy of negatively charged carbon clusters. They found that clusters with less than eleven carbon atoms exhibit an even-odd alternation in their electron affinity value, the odd having lower electron affinity. For clusters having more than ten carbon atoms it was observed that a new series can be drawn in which the reverse odd-even dependence of the electron affinity exists. In conjunction with theory, the two series were attributed to chain and ring structures, respectively. By varying the cluster formation conditions, the authors could observe, in the case of C_{11}^- , both the linear and the ring form.

Bernholdt *et al.*¹ in a theoretical investigation predicted, however, that a cyclic rhombic form exists and is isoenergetic with the linear form. It was further claimed that while the electron affinity of the linear C_4 is about 3.3 eV, the electron affinity of the rhombic form is about 2 eV.

Recently we developed a system which combines a laser photodetachment setup with the Coulomb explosion imaging method.⁴ At the Pelletron accelerator at the Weizmann Institute, a chopped negatively charged carbon cluster beam was produced by sputtering with Cs. After mass selection the clusters were accelerated by a field of 12 MV. When reaching a field free region at the middle of the accelerator,

the chopped ion beam interacted with a frequency doubled Nd:Yag laser (Quantell-535 nm, 25 mJ/cm², 20 ns long pulse). Electrons were photodetached and the neutral clusters drifted 30 m till they impinged upon a thin Formvar foil (about 30 Å thick). Following the foil, the charges of the four "Coulomb exploding" carbon ions were separated by a magnetic field. Their position and time of arrival were measured by a detector that combines a microchannel plate with hybridized anode and digitized video.⁵ It is evident from the two dimensional projection of the "exploded" structure, that the neutral C_4 enters the foil as a ring. Hence we present here the analysis of two-dimensional projections in which four C^{3+} ions were selected. This information is sufficient for distinguishing between the linear and cyclic forms.

For the analysis we used special body coordinates⁶ for the two-dimensional projection. These are

$$E_b = \frac{1}{\sqrt{12}} \{2(r_{12} + r_{34}) - (r_{13} + r_{14} + r_{23} + r_{24})\},$$

$$E_t = \frac{1}{2} (r_{13} - r_{14} + r_{24} - r_{23}),$$

where r_{ij} ($i, j = 1, \dots, 4$) are the carbon-carbon distances.

In Fig. 1(A) the data is presented using these two coordinates. For comparison in Figs. 1(B) and 1(C) simulations of the results for a rhombic and linear forms, respectively, are shown. Since the molecules are expected to contain internal energy,⁷ some internal motion was invoked in the simulation. The data prove unambiguously that many of the neutrals measured have the rhombic form. The experiment was repeated with a laser energy of about 3.6 eV (third harmonic of a Nd:Yag laser), and identical results were obtained.

The apparent inconsistency between the results present-




FIG. 1. The data presented in the body coordinates as discussed in the text (A), and simulation of the data assuming rhombus configuration (B) and linear shape (C).

ed here and those shown in Ref. 3 can be resolved as follows. Yang *et al.* applied the second harmonic of a Nd:Yag laser to sputter carbon, and an excimer laser for producing photoelectrons in the nozzle's throat. In both processes species with electron affinity below 2.34 (corresponding to 535 nm of the doubled Nd:Yag), can be easily neutralized. In our source on the other hand no discrimination against the low electron affinity species exists, and hence it is more probable to observe them. Thus, it is probable that cyclic C_n with $n < 10$ exist, although so far only the cyclic structure of C_4 has been established experimentally. Further work is on the way to obtain the structure of larger carbon clusters, and the full three-dimensional structure as a function of the laser frequency.

This work was partially supported by the US-Israel Binational Foundation, and by the MINERVA Fund, Munich, Germany.

¹D. E. Bernholdt, D. H. Magers, and R. J. Bartlett, *J. Chem. Phys.* **89**, 3612 (1988), and references cited therein.

²R. J. van Zee, R. F. Ferrante, K. J. Zeringue, and W. Weltner, Jr., *J. Chem. Phys.* **86**, 5212 (1987).

³S. Yang, K. J. Taylor, M. J. Craycraft, J. Conceicao, C. L. Pettiette, O. Cheshnovsky, and R. E. Smalley, *Chem. Phys. Lett.* **144**, 431 (1988).

⁴J. Zajtman, H. Kovner, Z. Vager, and R. Naaman (to be published).

⁵M. Algranati, H. Kovner, Z. Vager, and J. Zajtman (to be published).

⁶Z. Vager, in *The Structure of Small Molecules and Ions*, edited by R. Naaman and Z. Vager (Plenum, New York, 1989).

⁷I. Plesser, Z. Vager, and R. Naaman, *Phys. Rev. Lett.* **56**, 1559 (1986).