

Photography of Crystal Structures

Maurice L. Huggins

Citation: *The Journal of Chemical Physics* **12**, 520 (1944); doi: 10.1063/1.1723902

View online: <http://dx.doi.org/10.1063/1.1723902>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/12/12?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Photography](#)

Phys. Today **8**, 32 (1955); 10.1063/1.3062033

[Crystal Settings for Upper Level Photography, Precession Method](#)

Rev. Sci. Instrum. **25**, 928 (1954); 10.1063/1.1771222

[Streak Photography](#)

J. Appl. Phys. **21**, 445 (1950); 10.1063/1.1699682

[Elementary Photography](#)

Am. J. Phys. **17**, 523 (1949); 10.1119/1.1989685

[The Teaching of Photography](#)

Am. J. Phys. **7**, 394 (1939); 10.1119/1.1991491



Letters to the Editor

P*PROMPT publication of brief reports of important discoveries in physics may be secured by addressing them to this department. The closing date for this department is the third of the month. Because of the late closing date for the section no proof can be shown to authors. The Board of Editors does not hold itself responsible for the opinions expressed by the correspondents. Communications should not in general exceed 600 words in length.*

Photography of Crystal Structures

MAURICE L. HUGGINS

Eastman Kodak Company, Rochester, New York

November 1, 1944

AS shown by the writer¹ several years ago, Sir Lawrence Bragg's² method of photographic summation of Fourier series by addition of the proper patterns of light and dark bands can be improved in accuracy and speed by the use of a previously prepared set of masks.




FIG. 1. Photograph, looking along one of the crystallographic axes, of the body-centered cubic unit of α -iron, from x-ray data by Armstrong.

With the assistance of the Physics Department of these Laboratories, an improved set of 316 masks for this purpose has recently been made. Using these, electron density photographs of crystal structures and Patterson summations are easily and rapidly made from the appropriate x-ray data. Examples are shown in Figs. 1-3. The magnification is of the order of 100,000,000 in each case.

Most of the background density, between the atoms, in Fig. 3³ is not real, but is a result of inaccuracies in the x-ray data and in the masks and incompleteness of the former. The relative positions and approximate relative densities of the atomic peaks are correctly shown, however. In Fig. 1,⁴ the background was removed by a short treatment with Farmer's reducer solution. In Fig. 2⁵ the background has been practically all removed by reprinting.

The shape of the molecule in Fig. 2 is somewhat dis-




FIG. 2. Photograph of the hexamethylbenzene molecule, using x-ray data from the crystal, by Brockway and Robertson.

torted, both because the plane of the molecule is not parallel to the plane of the projection and because the projection, as obtained, is a square, whereas the true projection of the unit cell is not. The latter cause of distortion can readily be removed, if desired, by adding another optical step or in other ways. For purposes of structure analysis, however, this is not necessary.




FIG. 3. Photograph of a β -resorcinol (*meta*-dihydroxybenzene) crystal, using x-ray data by Robertson and Ubbelohde. Two complete molecules, with parts of others, are shown.

We hope soon to be able to furnish duplicate copies of the improved set of masks, at a nominal cost, to others engaged in crystal structure analysis.

¹ M. L. Huggins, J. Am. Chem. Soc. **63**, 66 (1941).

² W. L. Bragg, Zeits. f. Krist. **A70**, 475 (1929); *The Crystalline State* (The Macmillan Company, London and New York, 1934), p. 229.

³ J. M. Robertson and A. R. Ubbelohde, Proc. Roy. Soc. **A167**, 122 (1938).

⁴ A. H. Armstrong, Phys. Rev. **34**, 931 (1929).

⁵ L. O. Brockway and J. M. Robertson, J. Chem. Soc., p. 1324 (1939).