

Geometry of the Molecules CH₃Cl and CH₃Br

Börge Bak

Citation: [The Journal of Chemical Physics](#) **14**, 698 (1946); doi: 10.1063/1.1724087

View online: <http://dx.doi.org/10.1063/1.1724087>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/14/11?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Dissociative adsorption of CH₃X \(X = Br and Cl\) on a silicon\(100\) surface revisited by density functional theory](#)

J. Chem. Phys. **141**, 174701 (2014); 10.1063/1.4899841

[Photochemistry of adsorbed molecules. XIV. Photoejection of CH₃X \(X=Cl, Br\) from CH₃Br/LiF\(001\)](#)

J. Chem. Phys. **103**, 461 (1995); 10.1063/1.469611

[Theoretical study of geometries and dipole moments of AsX₃ and SbX₃ \(X=F, Cl, Br, and I\) molecules](#)

J. Chem. Phys. **89**, 4452 (1988); 10.1063/1.454783

[Reaction rates of O+CH₃Br and O+CH₃Cl](#)

J. Chem. Phys. **62**, 4477 (1975); 10.1063/1.430354

[The Geometry of CH₃X Molecules](#)

J. Chem. Phys. **16**, 553 (1948); 10.1063/1.1746941



Letters to the Editor

THIS section will accept reports of new work, provided these are terse and contain few figures, and especially few halftone cuts. The Editorial Board will not hold itself responsible for opinions expressed by the correspondents. Contributions to this section should not exceed 600 words in length and must reach the office of the Managing Editor not later than the 15th of the month preceding that of the issue in which the letter is to appear. No proof will be sent to the authors. The usual publication charge (\$3.00 per page) will not be made and no reprints will be furnished free.

Geometry of the Molecules CH_3Cl and CH_3Br

BÜRGE BAK

University of Copenhagen, Copenhagen, Denmark
October 7, 1946

IN organic-chemical problems one often meets with the question to what extent the substitution of one or more hydrogen atoms by so-called "electronegative" groups alters the original stereochemistry of the compound. The simplest case which can be studied in this respect is the substitution of one of the hydrogen atoms of methane by a halogen atom. Among the methyl halides conditions for studying the geometry are most favorable in the case of CH_3Cl and CH_3Br . Here, comparatively exact spectroscopic and interferometric data are available, which, in connection with thermodynamic data, give the best possible information on intramolecular distances and angles.

Penney¹ has given an approximative quantum-mechanical treatment of the bond energies and the valency angles of the methyl halides. To quote Penney, the result is that "if some or all of the hydrogen atoms in methane are replaced by other monovalent groups, the resulting deviations from the tetrahedral angle are remarkably small and can hardly exceed a few degrees."

If the greatest moment of inertia of a methyl halide is denoted by I_A , the small one by I_C , the C-X distance by a (X being a halogen atom), the C-H distance by d , the valency angle H-C-X by α and the mass of the atom or molecule Y by m_Y , we have:

$$I_C = 3m_H d^2 \sin^2 \alpha \quad (\text{I})$$

$$(I_A - \frac{1}{2}I_C)m_{\text{CH}_3\text{X}} = a^2 m_X m_C + 3(a-d \cos \alpha)^2 m_X m_H + 3d^2 \cos^2 \alpha m_C m_H. \quad (\text{II})$$

The quantities to be determined are a , d , and α . Sutherland² has discussed the matter from the point of view that the methyl group of the methyl halides has the same shape as in methane, e.g., that $d=1.09\text{\AA}$ and $\alpha=110^\circ$. In the case of CH_3Cl , I_A and I_C were known from spectroscopic work. Thus, Sutherland could calculate a , the C-Cl distance. He found $a=1.66\text{\AA}$ with a maximum possible error of 0.05\AA . However, from electronic diffraction experiments³ a had been found to be $1.77\text{\AA} \pm 0.02\text{\AA}$. A serious discrepancy seemed to exist. As is easily seen this discrepancy will vanish if we assume that the methyl group of the methyl halides and of methane may have a different shape. Indeed, new experimental data, published since the paper of Sutherland, seem to support this point of view.

In the case of CH_3Cl we can take $I_A=57.9 \cdot 10^{-40} \text{ gcm}^2$ from the paper by Nielsen.⁴ The value is good to 1-2 percent. $I_C=5.44 \cdot 10^{-40} \text{ gcm}^2$ is good to about 5 percent.^{5,6} Setting $a=1.77 \pm 0.02\text{\AA}$ we get:

$I_A \cdot 10^{40}$	$I_C \cdot 10^{40}$	$a \text{ \AA}$	α°	$d \text{ \AA}$	$\langle \text{HCH} \rangle^\circ$
57.9	5.44	1.77	93	1.05	118
59.0	5.17	1.75	102	1.04	114
Val. for CH_3 group in CH_4			110	1.09	110

$\alpha=93^\circ$ is the value obtained by using middle numbers of all experimental data. $\alpha=102^\circ$ is the maximum possible value. To some extent a similar result can be established in the case of CH_3Br . Taking $I_C=5.37 \cdot 10^{-40} \text{ gcm}^2 \pm 5$ percent^{5,6} and $a=1.91 \pm 0.06\text{\AA}$ ⁷ we only miss a value for I_A to be able to calculate α and d . Egan and Kemp⁸ have determined the standard entropy of CH_3Br at 276.66°K . Besides being a function of a number of quantities, such as the vibrational frequencies of the molecule, which are all known practically exact, the entropy is a function of $I_A^2 \cdot I_C$. We get:

$$I_A^2 I_C = 33,400 \cdot 10^{-120} \text{ g}^3 \text{ cm}^6 \pm 10 \text{ percent.}$$

Substituting $I_C=5.37 \cdot 10^{-40}$ we get $I_A=78 \cdot 10^{-40} \pm 8$ percent. Another estimate of I_A can be obtained from the doublet separation, $\delta\nu$, of the \parallel bands in infra-red. Taking the average value of the $\delta\nu$'s which have been determined by several authors,^{9,9-11} as $25.0 \text{ cm}^{-1} \pm 1 \text{ cm}^{-1}$ we get: $I_A=82 \cdot 10^{-40} \pm 10$ percent.

In a paper to be published the author¹² has studied the equilibrium $\text{CH}_3\text{Br} + \text{HCl} \rightleftharpoons \text{CH}_3\text{Cl} + \text{HBr}$ in a temperature interval from 70°C to 335°C . The exact knowledge of the equilibrium constant at different temperatures permits of calculating the ratio $I_A(\text{CH}_3\text{Br}):I_A(\text{CH}_3\text{Cl})$, as $I_C(\text{CH}_3\text{Br})$ to a good approximation is equal to $I_C(\text{CH}_3\text{Cl})$. The result is that $I_A=76.0 \cdot 10^{-40} \pm 4$ percent. From these three determinations we conclude that $I_A=77.5 \cdot 10^{-40} \text{ gcm}^2 \pm 4$ percent. The corresponding values of α and d are given below.

$I_A \cdot 10^{40}$	$I_C \cdot 10^{40}$	$a \text{ \AA}$	α°	$d \text{ \AA}$	$\langle \text{HCH} \rangle^\circ$
77.5	5.37	1.91	90	1.04	120
80.5	5.11	1.85	111	1.08	110
Val. for CH_3 group in			110	1.09	110

$\alpha=90^\circ$ is the value obtained using middle numbers of all experimental data. $\alpha=111^\circ$ is the maximum possible value.

For CH_3Cl it thus seems firmly established that α deviates considerably from the tetrahedral angle. For CH_3Br the same is very probably the case. The methyl group in CH_3Cl and CH_3Br thus seems to be considerably more 'flat' than in CH_4 .

¹ Penney, Trans. Faraday Soc. **31**, 734 (1935).

² Sutherland, Trans. Faraday Soc. **34**, 325 (1938).

³ Sutton and Brockway, J. Am. Chem. Soc. **57**, 473 (1935).

⁴ Nielsen, Phys. Rev. **56**, 847 (1939).

⁵ Bennett and Mayer, Phys. Rev. **32**, 888 (1928).

⁶ Johnston and Dennison, Phys. Rev. **48**, 868 (1935).

⁷ Levy and Brockway, J. Am. Chem. Soc. **59**, 1662 (1937).

⁸ Egan and Kemp, J. Am. Chem. Soc. **60**, 2097 (1938).

⁹ Moorhead, Phys. Rev. **39**, 788 (1932).

¹⁰ Sleator, Phys. Rev. **38**, 147 (1932).

¹¹ Barker and Nielsen, Phys. Rev. **46**, 970 (1934).

¹² Bak, Dansk. Vidensk. Selsk. math.-fys. Medd. (1947).