

Infrared Spectra of HF Complexes in Noble-gas Solids

Lester Andrews

Department of Chemistry, University of Virginia, Charlottesville, Virginia 22901, U.S.A.

The very weak hydrogen-bonded complex of molecular hydrogen and HF has been prepared by condensing the neon-diluted reagents at 5 K. Infrared spectra of the $\text{H}_2 \cdots \text{HF}$ complex revealed a $\nu_s(\text{HF})$ stretching mode at 3938 cm^{-1} and $\nu(\text{H}_2)$ mode at 4155 cm^{-1} , which are bathochromically shifted from the isolated molecule values of 15 and 6 cm^{-1} , respectively. The very strong $\text{NH}_3 \cdots \text{HF}$ complex has been studied in solid neon and solid argon. The $\nu_s(\text{HF})$ mode at 3215 cm^{-1} in the gas phase is shifted to 3106 cm^{-1} in solid neon and 3041 cm^{-1} at solid argon. The $\nu_1(\text{HF})$ librational mode appeared at 916 cm^{-1} with $2\nu_1$ at 1679 cm^{-1} , while $\nu_2(\text{NH}_3)$ was hypsochromically shifted to 1093 cm^{-1} in solid argon. The antihydrogen-bonded arrangement is the more stable structure for the $\text{HF} \cdots \text{Cl}_2$ complex in the gas phase; however, solid argon and neon samples reveal both antihydrogen-bonded and hydrogen-bonded complexes. In the $\text{Cl}_2 \cdots \text{HCl}$ case, evidence for only the hydrogen-bonded form was observed in solid argon

Matrix-isolation studies of reactive molecules and complexes can make unique contributions to the spectroscopy of these species by taking advantage of the matrix host. These advantages include accumulation of sufficient sample to observe weak absorptions, long cooling time to ensure that minimum temperature is reached by the sample, and fast relaxation of new energized species to prevent decomposition. In addition, a complete low-resolution spectrum can be recorded to provide a useful guide for diode laser studies. This report will describe chemical systems that exploit these advantages of matrix-isolation spectroscopy.

The $\text{NH}_3 \cdots \text{HF}$ Complex

The strong $\text{NH}_3 \cdots \text{HF}$ complex was one of the first hydrogen-bonded complexes studied in our group and it remains one of the most interesting.^{1,2} The $\nu_s(\text{HF})$ fundamental and microwave spectrum have been observed in the gas phase (unpublished work by Thomas and Clements *et al.*), and the complex has three-fold symmetry. A series of matrix experiments was performed with isotopic ammonia and hydrogen fluoride samples in argon at 12 K, and the new absorptions are collected in table 1. The spectrum recorded for DF/HF and NH_3 identifies the HF submolecule modes due to the appearance of DF counterparts and the NH_3 submolecule mode (ν_2^s), which shows only a very small effect for the acid isotope.^{1,2} The strong $\nu_s(\text{DF})$ band at 2278 cm^{-1} (f.w.h.m. = 4 cm^{-1}) has a $\nu_s(\text{HF})$ counterpart at 3041 cm^{-1} (f.w.h.m. = 16 cm^{-1}). Sharp librational ν_1 bands were observed for HF and DF at 916.0 and 696.7 cm^{-1} with their overtones at 1679.4 and 1315.9 cm^{-1} . The HF submolecule fundamentals observed for the complex are the $\nu_s(\text{HF})$ stretching fundamental and the $\nu_1(\text{HF})$ librational mode derived from rotational degrees of freedom from the isolated diatomic. The doubly degenerate nature of this librational motion is maintained by the C_{3v} symmetry of the complex. The $\nu_s(\text{HF})/\nu_s(\text{DF}) = 3041/2278 = 1.334$ ratio is characteristic of the HF stretching mode in a hydrogen-bonded complex. The overtone ratio $2\nu_1(\text{HF})/2\nu_1(\text{DF}) = 1679/1316 = 1.276$ is less than the fundamental ratio $\nu_1(\text{HF})/\nu_1(\text{DF}) = 916/697 = 1.314$ owing to increased


Table 1. Absorptions (cm^{-1}) for the ammonia–hydrogen fluoride complex in solid neon and argon

neon				argon				assignment
$\text{NH}_3 + \text{HF}$	$\text{NH}_3 + \text{DF}$	$\text{NH}_3 + \text{HF}$	$\text{NH}_3 + \text{DF}$	$^{15}\text{NH}_3 + \text{HF}$	$^{15}\text{NH}_3 + \text{DF}$	$\text{ND}_3 + \text{HF}$	$\text{ND}_3 + \text{DF}$	
3106	2309	3041	2278	3042	2277	3017	2269	ν_s
1676	1317	1679	1316	1678	1314	1664	1272	$2\nu_1$
1090	1091	1093	1094	1087	1088	849	849	ν_2^c
912	695	916	697	915	695	906	675	ν_1

anharmonicity in the overtone. The positions of the ν_s and ν_1 modes provide a measure of the interaction between acid and base as the ν_s mode is decreased from the 3919 cm^{-1} stretching fundamental and the ν_1 mode is increased from the 44 cm^{-1} rotational fundamental of the diatomic.

The ammonia submolecule mode ν_2^c in the complex follows $^{15}\text{NH}_3$ and ND_3 substitution, as does ν_2 of ammonia. The ν_2^c mode is not split by inversion as ν_2 for NH_3 is split; this shows that the acid ligand presents inversion of the ammonia submolecule in the complex 1. Note the effect of $^{15}\text{NH}_3$ and ND_3 substitution on the HF submolecule modes; this verifies that a reagent complex has been formed in the matrix.

The $\text{NH}_3 \cdots \text{HF}$ complex has been prepared in solid neon for comparison with the argon-matrix data (table 1). First, the strong ν_s band at 3106 cm^{-1} (f.w.h.m. = 10 cm^{-1}) is sharper than in solid argon and it falls approximately midway between the gas-phase (3215 cm^{-1}) and argon-matrix (3041 cm^{-1}) values. The $2\nu_1$ and ν_1 modes are shifted only a slightly and the ratio is essentially unchanged from the solid-argon value. The ν_2^c band at 1090 cm^{-1} is also shifted only slightly from the solid-argon value, but the shift above NH_3 in solid neon, 122 cm^{-1} , is essentially the same as in solid argon, 120 cm^{-1} . The HF/DF ratios in solid neon are essentially unchanged from solid-argon values. The neon medium is more inert, and the argon matrix effect is clearly seen to be greater on ν_s than on ν_1 and ν_2^c . This probably arises from the greater polarizability of argon and the large dipole change in the ν_s vibration.



$\text{NH}_3 \cdots \text{F}_2$ and $\text{NH}_2\text{F} \cdots \text{HF}$ Complexes

The fluorine–ammonia matrix system yielded two interesting new complexes.³ On codeposition of Ar/NH_3 and Ar/F_2 samples, new bands appeared at 966 and 781 cm^{-1} . These bands increased threefold on sample annealing and decreased markedly on photolysis that produced the $\text{NH}_2\text{F} \cdots \text{HF}$ complex to be described below, hence assignment of the strong 966 and weak 781 cm^{-1} bands to the $\text{NH}_3 \cdots \text{F}_2$ complex 2 follows. The strong 966 cm^{-1} band is within 1 cm^{-1} of the median of the inversion-split Q branches of NH_3 in solid argon,⁴ which demonstrates that even the weak $\text{NH}_3 \cdots \text{F}_2$ interaction prevents inversion of the NH_3 submolecule in the complex. The weak 781 cm^{-1} band is below the 892 cm^{-1} Raman fundamental of F_2 in solid argon, which suggests a weak charge-transfer interaction between NH_3 and F_2 .

The major photolysis product bands in the $\text{NH}_3 \cdots \text{F}_2$ system appeared at 3389 cm^{-1} (ν_s), 3269 cm^{-1} (ν_1), 1568 cm^{-1} (ν_2), 1314 cm^{-1} ($2\nu_1$), 1244 cm^{-1} (ν_3), 934 cm^{-1} (ν_4) and


Table 2. Infrared absorptions (cm^{-1}) for the amine-bonded $\text{H}_2\text{FN}\cdots\text{HF}$ complex in solid argon

$\text{H}_2\text{FN}\cdots\text{HF}$	$\text{H}_2\text{F}^{15}\text{N}\cdots\text{HF}$	$\text{HDFN}\cdots\text{HF}$	$\text{D}_2\text{FN}\cdots\text{DF}$	assignment
3389	3388	3384	2505	ν_s
3269	3264		2399	ν_1 ($\text{N}-\text{H}_2$ str)
1568	1566	1429	1151	ν_2 ($\text{N}-\text{H}_2$ bend)
1314	1314	1312	998	$2\nu_1$
1244	1238		968	ν_3 (NH_2 wag)
934	916	933	924	ν_4 ($\text{N}-\text{F}$ str)
750	750		561	ν_1 (in-plane)
723	723		540	ν_1 (out-plane)

750, 723 cm^{-1} (ν_1), which are listed in table 2 along with their ND_3 and $^{15}\text{NH}_3$ counterparts. The ν_s and ν_1 bands are clearly due to acid submolecule and the ν_1 , ν_2 , ν_3 and ν_4 bands to base submolecule modes in a new complex with the structure proposed in 3 based on the spectrum. This nitrogen lone-pair complex $\text{H}_2\text{FN}\cdots\text{HF}$ is weaker than $\text{H}_3\text{N}\cdots\text{HF}$ as ν_s is displaced less (to 3389 cm^{-1}) and the ν_1 modes are increased less (to 750 and 723 cm^{-1}). The base has less than threefold symmetry based on splitting of the librational mode into two bands. The other four bands show isotopic shifts and components for the NH_2F vibrations as assigned in the table. Unfortunately isolated NH_2F has not been characterized owing to its extremely great reactivity, and it is observed here because of efficient quenching of the reaction exothermicity by the cold matrix.

Hydrogen Molecule–HF Complex

Very weak, hydrogen-bonded complexes of molecular hydrogen and HF have been prepared by condensing the neon-diluted reagents at 5 K. Infrared spectra of an $\text{Ne}/\text{H}_2/\text{HF}$ sample in fig. 1 reveal HF monomer, dimer (D) and trimer (T) bands at 3992, 3953, 3919, 3848 and 3706 cm^{-1} , a strong, sharp ν_s band at 3838 cm^{-1} , a weak ν^c band at 4155 cm^{-1} , a weaker ν^{cc} band at 4146 cm^{-1} and split ν_{sa} and ν_{sb} bands at 3889 and 3840 cm^{-1} .^{5,6} Changes in reagent concentration and sample annealing show that the ν^{cc} , ν_{sa} and ν_{sb} bands are due to a higher-order complex than ν^c and ν_s . Accordingly, the primary product complex is identified as $\text{H}_2\cdots\text{HF}$ (4) and the higher-order complex as $\text{H}_2\cdots(\text{HF})_2$ (5). Vibrational assignments are aided by the isotopic data given in table 3. The ν_s band is displaced 15 cm^{-1} below isolated HF in solid neon (3953 cm^{-1}) and exhibits an HF/DF ratio of 1.364; which is appropriate for an HF complex; the ν^c band is shifted 6 cm^{-1} below the H_2 fundamental in the gas phase and exhibits an H_2/D_2 ratio of 1.389, near that for the gaseous diatomics (1.390). Accordingly, the ν_s and ν^c bands are assigned to the H–F and H–H fundamentals in the $\text{H}_2\cdots\text{HF}$ complex, and in like fashion, the ν^{cc} , ν_{sa} and ν_{sb} assignments follow as well. Comparison of fundamentals for 4, 5 and $(\text{HF})_2$ is of chemical interest. The H_a-F and H_b-F fundamentals of $(\text{HF})_2$ at 3919 and 3848 cm^{-1} are displaced to 3889 and 3840 cm^{-1} by the addition of H_2 to form 5; note the greater effect for the H_a-F mode where H_2 is attached. Likewise, when H_b-F binds to 4 the H–F mode is shifted from 3938 to 3889 cm^{-1} owing to the fluoride affinity of H_b-F .






Fig. 1. Infrared spectra ($4200\text{--}3700\text{ cm}^{-1}$) for hydrogen and hydrogen fluoride in solid neon: 11 mmol of $\text{Ne}/\text{H}_2 = 50/1$ and 11 mmol of $\text{Ne}/\text{HF} = 200/1$ codeposited at 4 K.

Table 3. Absorptions (cm^{-1}) for molecular hydrogen and hydrogen fluoride complexes in solid neon at 5 K


$\text{H}_2 + \text{HF}$	$\text{H}_2 + \text{DF}$	$\text{D}_2 + \text{HF}$	$\text{D}_2 + \text{DF}$	assignment
4155	4152	2991	2988	ν^c (4)
4146	4143	2983	2980	ν^{cc} (5)
3938	2887	3938	2886	ν_s (4)
3889	2850	3887	2848	ν_{sa} (5)
3840	2814	3838	2813	ν_{sb} (5)

Note the relatively larger effect of DF on H_2 and D_2 than HF, which is listed in table 3. This is due to the shorter time-average distance of D from H_2 than H owing to the smaller librational amplitude of DF. In the $\text{D}_2\cdots\text{HF}$ complex the ν^c mode was bathochromically shifted only 3 cm^{-1} below the gas-phase fundamental for D_2 . Clearly the D_2 oscillator is less susceptible to electrical interactions than H_2 since D_2 is substantially lower in the potential well.

Initial studies on $\text{H}_2\cdots\text{HF}$ were performed in solid argon. The $\text{H}_2\cdots\text{HF}$ complex gave a ν_s mode shifted hypsochromically $8\text{--}17\text{ cm}^{-1}$ above the HF fundamental in solid argon, and the value of these results was confused by interaction with the argon matrix. Replacing one argon atom by H_2 provided a repulsive contribution to the HF intramolecular potential function and hypsochromic shift resulted since H_2 is only slightly more physically inert than argon. Neon is, however, still more inert and neon-matrix studies of very weak complexes are representative of gas-phase studies. Accordingly, the neon-matrix work, taking the weak $\text{Ar}\cdots\text{HF}$ complex as a model, predicts a gas-phase band at 3950 cm^{-1} for $\text{H}_2\cdots\text{HF}$,⁵ which has subsequently been observed.^{7,8} The gas-phase spectrum has been analysed in terms of a $j = 1$ hindered H_2 rotor in an anisotropic potential with a minimum energy at the T-shaped geometry; no evidence was found for the corresponding *para*- $\text{H}_2\cdots\text{HF}$ complex.⁷ The neon-matrix experiments provide data for only one 1:1 complex, which is most probably the same *ortho*- $\text{H}_2\cdots\text{HF}$ complex characterized in the gas phase. In other words, HF 'traps' H_2 in the *ortho* spin state, whereas in a matrix with trace O_2 present, conversion to the *para* spin state eventually occurs for H_2 molecules isolated in the matrix.⁵

Chlorine-Hydrogen Halide Complexes

The $\text{HF}\cdots\text{Cl}_2$ complex has been studied in a molecular beam⁹ and by theoretical calculations.¹⁰ Both conclude that the anti-hydrogen-bonded $\text{HF}\cdots\text{Cl}_2$ arrangement (6) is more stable than the hydrogen-bonded $\text{Cl}_2\cdots\text{HF}$ structure (7); however, the energy difference is calculated to be small ($<100\text{ cm}^{-1}$). In an argon matrix at 5 K, HF/Cl_2 mixtures¹¹ gave sharp new bands at 3901 and 2864 cm^{-1} (labelled ν_s) above the $\text{N}_2\cdots\text{HF}$ and $\text{N}_2\cdots\text{DF}$ complexes (labelled N)¹² and sharp multiplets at 3860 , 3853 , 3848 and 2831 , 2826 , 2822 cm^{-1} (ν_s) between N and hydrogen fluoride dimer (D), fig. 2(a). In an argon matrix at 12 K, the ν_s bands were stronger relative to ν_s , and on sample annealing to 23 K, both ν_s and ν_s features increased, fig. 2(b) and (c), but the ν_s band increased more than ν_s in the case of HF.



The new absorptions in $\text{HF}/\text{DF}/\text{Cl}_2$ matrix samples are considered for 1:1 complexes between HF and Cl_2 . It is expected that the $\text{HF}\cdots\text{Cl}_2$ arrangement perturbs the $\text{H}\cdots\text{F}$ fundamental less than the $\text{Cl}_2\cdots\text{HF}$ structure since this vibration is primarily a hydrogen motion. The ν_s band at 3901 cm^{-1} is bathochromically shifted 18 cm^{-1} from the HF-induced Q branch² at 3919 cm^{-1} , not as much as the $\text{HF}\cdots\text{HCN}$ complex¹³ at 3891 cm^{-1} , and the strongest ν_s band at 3860 cm^{-1} is bathochromically shifted 59 cm^{-1} , more than $\text{N}_2\cdots\text{HF}$ at 3881 cm^{-1} . These data are consistent with assignment of the ν_s band to 6 and the ν_s bands to 7. Parallel studies with CIF support these assignments as CIF is a




Fig. 2. Infrared spectra in the HF and DF regions for Cl_2 mixtures in solid argon: (a) after codeposition of $\text{Ar}/(\text{HF} + \text{DF}) = 200/1$ (60% DF) and $\text{Ar}/\text{Cl}_2 = 200/1$ at 5 K for 3 h, (b) after deposition of $\text{Ar}/(\text{HF} + \text{DF}) = 200/1$ (90% DF) and $\text{Ar}/\text{Cl}_2 = 200/1$ at 12 K for 4 h, (c) after warming to 23 K and recooling to 12 K.

stronger Lewis acid but a weaker Lewis base than Cl_2 owing to the inductive effect of fluorine.¹¹ The $\text{HF} \cdots \text{ClF}$ complex absorption at 3896 cm^{-1} indicates a stronger interaction than **6**, and $\text{FCl} \cdots \text{HF}$ band at 3870 cm^{-1} denotes a weaker interaction than that found for **7**.

The greater relative yield of the more stable **6** arrangement compared to **7** in 12 K argon matrices is consistent with beam experiments and calculations. However, freezing the samples more quickly at 5 K gives more absorbance for the **7** species, and this is more pronounced for DF than HF. Similar observations¹¹ were found in solid neon at 5 K. Apparently the lower temperature does not allow thermodynamic equilibrium to be established between structures and traps relatively more of the less stable arrangement. The matrix results suggest that **9** is closer in energy to **8** and perhaps even lower and invite gas-phase and theoretical studies on the DF species **8** and **9** to test this hypothesis.




Fig. 3. Infrared spectra ($2900\text{--}2800\text{ cm}^{-1}$) for chlorine and hydrogen chloride in solid argon: (a) 6 mmol of $\text{Ar}/\text{HCl}=200/1$ deposited at 5 K, (b) $\text{Ar}/\text{HCl}=200/1$ codeposited at 12 K with $\text{Ar}/\text{Cl}_2=100/1$ sample passed through microwave discharge and (c) continuation of (a) with 7 mmol of $\text{Ar}/\text{HCl}=200/1$ codeposited at 5 K with 3.5 mmol of $\text{Ar}/\text{Cl}_2=100/1$ sample.

Similar experiments have been performed with the HCl/DCI/Cl₂ system to prepare the analogous complexes.¹⁴ Fig. 3 illustrates spectra for Ar/HCl: the strong HCl R(0) band, the induced Q doublet, the N₂⋯HCl complex, the HCl P(1) and (HCl)₂ bands are in agreement with earlier studies.¹⁵ On deposition with Ar/Cl₂ a new chlorine isotopic doublet was observed at 2842.2, 2840.1 cm⁻¹ [labelled ν_s in fig. 3(c)] between the N doublet (2863.5, 2861.3 cm⁻¹) and (HCl)₂ (2817.5, 2815.4 cm⁻¹ sh); no new features were observed between the Q doublet (2871.0, 2868.7 cm⁻¹) and the N doublet. In a separate experiment the Ar/Cl₂ stream was passed through a microwave discharge to dissociate chlorine and was subsequently codeposited with an Ar/HCl stream, and in addition a new doublet was observed at 2836.4, 2834.5 cm⁻¹ (labelled R) [fig. 3(b)] and HCl₂⁻ appeared at 696 cm⁻¹.¹⁶ The chlorine isotopic doublet between N₂⋯HCl and (HCl)₂ is assigned to the Cl₂⋯HCl complex and the new band in the discharge experiments is due to the Cl⋯HCl complex. Again comparison with the ClF system,¹⁴ which reveals a new doublet at 2849.9, 2848.1 cm⁻¹ for the FCl⋯HCl complex, supports this identification of Cl₂⋯HCl because FCl⋯HCl is expected to be a weaker complex and exhibit a smaller shift in the H—Cl fundamental than the Cl₂⋯HCl complex. Finally, comparison of the HCl fundamental in the Cl₂⋯HCl complex at 2842.2 cm⁻¹ with HCl fundamentals in the Br₂⋯HCl and I₂⋯HCl complexes at 2820.2 and 2783.9 cm⁻¹ in solid argon, respectively,¹⁷ defines a trend of increasing hydrogen-bonding interaction with increasing halogen molecule size and polarizability. In addition the heavy halogen atom complexes with HCl exhibit stronger interactions with HCl than the diatomic,¹⁷ as found for chlorine.

Three points are noteworthy: (1) the chlorine atom interaction with HCl is slightly stronger than the Cl₂ molecule interaction, (2) the anion electron makes a significant difference in the bonding and structure of the weak Cl⋯H—Cl complex and the strongly hydrogen-bonded centrosymmetric anion HCl₂⁻, and (3) with HCl the favoured structure is the Cl₂⋯HCl hydrogen-bonded form, whereas with HF the anti-hydrogen-bonded form HF⋯Cl₂ discussed above is the more stable arrangement. Although HCl forms a weaker hydrogen bond than HF, the much larger covalent radius of chlorine results in a much weaker Lewis acid–base interaction HCl⋯Cl₂, which allows the Cl₂⋯HCl arrangement to be more stable in this system.

The author gratefully acknowledges National Science Foundation support for this work and the contributions of co-workers whose names are given in the references.

References

- 1 G. L. Johnson and L. Andrews, *J. Am. Chem. Soc.*, 1982, **104**, 3043.
- 2 L. Andrews, *J. Phys. Chem.*, 1984, **88**, 2940.
- 3 L. Andrews and R. Lascola, *J. Am. Chem. Soc.*, 1987, **109**, 6243.
- 4 L. Abouaf-Marguin, M. E. Jacox and D. E. Milligan, *J. Mol. Spectrosc.*, 1977, **67**, 34.
- 5 R. D. Hunt and L. Andrews, *J. Chem. Phys.*, 1987, **86**, 3781.
- 6 L. Andrews, V. E. Bondybey and J. H. English, *J. Chem. Phys.*, 1984, **81**, 3452.
- 7 C. M. Lovejoy, D. D. Nelson Jr and D. J. Nesbitt, *J. Chem. Phys.*, 1987, **87**, 5621.
- 8 K. W. Jucks and R. E. Miller, *J. Chem. Phys.*, 1987, **87**, 5629.
- 9 F. A. Baiocchi, T. A. Dixon and W. Klemperer, *J. Chem. Phys.*, 1982, **77**, 1632.
- 10 A. P. L. Rendell, G. G. Bacskay and N. S. Hush, *J. Chem. Phys.*, 1987, **87**, 535.
- 11 R. D. Hunt and L. Andrews, *J. Phys. Chem.*, 1988, **92**, 3769.
- 12 L. Andrews, B. J. Kelsall and R. T. Arlinghaus, *J. Chem. Phys.*, 1983, **79**, 2488.
- 13 G. L. Johnson and L. Andrews, *J. Am. Chem. Soc.*, 1983, **105**, 163.
- 14 R. D. Hunt and L. Andrews, *J. Chem. Phys.*, 1988, **89**, 3502.
- 15 D. Maillard, A. Schriver and J. P. Perchard, *J. Chem. Phys.*, 1979, **71**, 505.
- 16 C. A. Wight, B. S. Ault and L. Andrews, *J. Chem. Phys.*, 1976, **65**, 1244 and references therein.
- 17 A. Engdahl and B. Nelander, *Chem. Phys.*, 1985, **100**, 273.