

Erratum: Jahn–Teller effect in tetrahedral d 1 metal complexes [J. Chem. Phys. 81, 1861 (1984)]

A. Agresti, J. H. Ammeter, and M. Bacci

Citation: [The Journal of Chemical Physics](#) **82**, 5299 (1985); doi: 10.1063/1.448988

View online: <http://dx.doi.org/10.1063/1.448988>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/82/11?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

Erratum: “The dynamic Jahn–Teller effect in Cu(II) doped MgO” [J. Chem. Phys. 130, 104708 (2009)]
J. Chem. Phys. **130**, 199901 (2009); 10.1063/1.3137184

Relativistic $E \times T$ Jahn–Teller effect in tetrahedral systems
J. Chem. Phys. **129**, 224102 (2008); 10.1063/1.3035189

Erratum: Theoretical calculation of the absorption and magnetic circular dichroism spectrum of a Jahn–Teller distorted excited state: The $1E'$ excited state of cyclopropane [J. Chem. Phys. 79, 2951 (1983)]
J. Chem. Phys. **84**, 544 (1986); 10.1063/1.450866

Jahn–Teller effect in tetrahedral d 1 metal complexes
J. Chem. Phys. **81**, 1861 (1984); 10.1063/1.447859

Erratum: Experimental confirmation of the Jahn–Teller distortion of CH_4 + [J. Chem. Phys. 72, 1402 (1980)]
J. Chem. Phys. **72**, 6818 (1980); 10.1063/1.439182

A promotional banner for AIP APL Photonics. The background is a vibrant orange with a sunburst effect. On the left is a small image of the journal cover, which shows a blue and yellow abstract pattern. To the right of the cover is a yellow starburst with the words 'OPEN ACCESS' in red. The main text 'Launching in 2016!' is in a large, white, sans-serif font. Below it, 'The future of applied photonics research is here' is in a smaller, white, sans-serif font. In the bottom right corner, the 'AIP APL Photonics' logo is displayed in white.

collide essentially only with solid surfaces. Let $p(l_s)$ be the probability that Knudsen limit trajectories exceed l_s from random starting positions in the pore space. The probability for no Knudsen limit surface collisions occurring between distances l_s and $l_s + dl_s$ is $1 - Kdl_s$, where K is a positive constant. Since averaging for $P(l_s)$ is taken over a macroscopically homogeneous and isotropic volume, the probability $P(l_s + dl_s)$ is equal to the product of the independent probabilities $P(l_s)$ and $P(dl_s)$. This gives

$$\frac{d}{dl_s} P(l_s) = -KP(l_s) \quad (5)$$

which upon integration results in $P(l_s) = \exp(-Kl_s)$. The distribution of interest is

$$p(l_s) = \frac{-d}{dl_s} P(l_s) = \frac{1}{r_0} e^{-l_s/r_0}, \quad (6)$$

where r_0 , equal to the mean value of l_s , replaces $1/K$.

The porous media mean free path l_e results from the integration of the product of Eqs. (4b) and (6) over all possible l_s ,

$$l_e = \frac{l_0}{r_0} \int_0^\infty \left\{ \exp\left(\frac{-l_s}{r_0}\right) - \exp\left[-l_s\left(\frac{l_0 + r_0}{l_0 r_0}\right)\right] \right\} dl_s \quad (7a)$$

$$= \frac{l_0 r_0}{l_0 + r_0}. \quad (7b)$$

rearranging Eq. (7b) shows $l_e^{-1} = l_0^{-1} + r_0^{-1}$. This form of l_e^{-1} is analogous to those of free path models for electron and phonon diffusion in solids.⁸ In each case additivity of the inverses of interparticle effects and particle-medium effects result.

Combining Eqs. (1) and (7b) provide the final expressions

$$D_e = \frac{1}{3} f(\epsilon) \bar{v} \frac{l_0 r_0}{l_0 + r_0} = \frac{f(\epsilon) D_0}{1 + \text{Kn}} \quad (8a)$$

and

$$D_e^{-1} = [f(\epsilon) D_0]^{-1} + \left[\frac{1}{3} f(\epsilon) \bar{v} r_0 \right]^{-1}. \quad (8b)$$

The Knudsen number Kn in Eq. (8a) is the ratio l_0/r_0 . By equating the last term in Eq. (8b) to D_K^{-1} , the inverse of the Knudsen diffusivity, the Bosanquet result is obtained. The Bosanquet formula, $D_e^{-1} = [f(\epsilon) D_0]^{-1} + D_K^{-1}$, is applicable in the Knudsen limit ($\text{Kn} \gg 1$), the continuum limit ($\text{Kn} \ll 1$), as well as intermediate ranges of Kn . The derivation presented here demonstrates that the Bosanquet result arises through the influence of pore surface collisions on the free path distribution.

I thank Professor T. N. Narasimhan and Professor L. J. Waldron for helpful comments. This work was supported by the Department of Energy under Contract No. DE-AC03-76SF00098.

¹W. G. Pollard and R. D. Present, *Phys. Rev.* **73**, 762 (1948).

²W. Strieder, *J. Chem. Phys.* **54**, 4050 (1971).

³F. G. Ho and W. Strieder, *J. Chem. Phys.* **70**, 5635 (1979).

⁴F. G. Ho and W. Strieder, *J. Chem. Phys.* **73**, 6296 (1980).

⁵R. B. Evans, G. M. Watson, and E. A. Mason, *J. Chem. Phys.* **35**, 2076 (1961).

⁶K. S. Spiegel, *Ind. Eng. Chem. Fundamentals* **5**, 529 (1966).

⁷J. Jeans, *An Introduction to the Kinetic Theory of Gases* (Cambridge University, Cambridge, 1940), pp. 142–145.

⁸C. A. Wert and R. M. Thomas, *Physics of Solids* (McGraw-Hill, New York, 1964), pp. 211–214.

ERRATUM

Erratum: Jahn–Teller effect in tetrahedral d^1 metal complexes [*J. Chem. Phys.* **81**, 1861 (1984)]

A. Agresti, J. H. Ammeter, and M. Bacci

Istituto di Ricerca sulle Onde Elettromagnetiche del CNR, Via Panciatichi 64, 50127 Firenze, Italy

The relationships (10) were erroneously evaluated. The right expressions are

$$A_2 = \frac{1}{R^2} (e_\sigma - e_\pi), \quad (10a)$$

$$b_{ee} = -\frac{\sqrt{3}}{R^2} \left(e_\sigma - \frac{5}{3} e_\pi \right), \quad (10b)$$

$$b_{rr} = \frac{3}{2R^2} \left(e_\sigma - \frac{5}{3} e_\pi \right), \quad (10c)$$

$$c_{rr} = \frac{7}{6R^2} \left(e_\sigma - \frac{1}{3} e_\pi \right). \quad (10d)$$

Accordingly column I of Table I should read

$$e_\sigma = 6655 \text{ cm}^{-1},$$

$$e_\pi = 520 \text{ cm}^{-1},$$

$$A_1 = 458 \text{ cm}^{-1}/\text{\AA},$$

$$A_2 = 1340 \text{ cm}^{-1}/\text{\AA}^2,$$

$$E_{JT} = 37 \text{ cm}^{-1},$$

$$b = -4946 \text{ cm}^{-1}/\text{\AA},$$

$$c = -2397 \text{ cm}^{-1}/\text{\AA},$$

$$c_1 = -4814 \text{ cm}^{-1}/\text{\AA}.$$

All conclusions of the paper remain still valid.