

Rotational effects on pyrazine S 1(1 B 3u) dynamics and the problem of the missing states

Aviv Amirav

Citation: *The Journal of Chemical Physics* **88**, 2840 (1988); doi: 10.1063/1.454715

View online: <http://dx.doi.org/10.1063/1.454715>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/88/4?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

The absolute value of the quantum yield of the fluorescence of the 1 B 3u 0–0 state of pyrazine as a function of the rotational quantum numbers

J. Chem. Phys. **86**, 4004 (1987); 10.1063/1.451909

Rotational state dependence of pyrazine fluorescence: Initial decays for the vibrationless 1 B 3u state

J. Chem. Phys. **82**, 1067 (1985); 10.1063/1.448478

Intermediate level structure, quantum beats, and nuclear spin effects in the electronic relaxation of the 1 B 3u ($n\pi^*$) state of pyrazine

J. Chem. Phys. **76**, 2099 (1982); 10.1063/1.443185

Single vibronic level fluorescence from 1 B 3u pyrazine: The role of Fermi resonance and Duschinski rotation

J. Chem. Phys. **74**, 4893 (1981); 10.1063/1.441742

Electronic relaxation of 1 B 3u ($n\pi^*$) pyrazine in a supersonic jet: Rotational state dependence of the nonradiative rate leading to biexponential decay

J. Chem. Phys. **74**, 3616 (1981); 10.1063/1.441468



The Letters to the Editor section is subdivided into four categories entitled Communications, Notes, Comments, and Errata. The textual material of each Letter is limited to 1200 words minus the following: (a) 200 words for a square figure one-column wide. Larger figures are scaled in proportion to their area; (b) 50 words for each displayed equation; (c) 7 words for each line of table including headings and horizontal rulings. Proof will be sent to authors. See the issue of 1 January 1988 for a fuller description of Letters to the Editor.

COMMUNICATIONS

Rotational effects on pyrazine $S_1(^1B_{3u})$ dynamics and the problem of the missing states

Aviv Amirav

School of Chemistry, Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978 Tel Aviv, Israel

(Received 2 June 1987; accepted 2 December 1987)

The $S_1(^1B_{3u})$ state of pyrazine serves as a touchstone for the theory of radiationless transitions. This theory is confronted with the wealth of experimental results. A full account on the "story of pyrazine" is given elsewhere.¹ One of the major problems, known as the "problem of the missing states," emerges from the number of spectrally observed, mixed singlet-triplet states [molecular eigenstates (MEs)] in each rotational transition. This number is too low and is rotational independent²⁻⁵ and thus is in conflict with the emission quantum yield^{6,7} and time resolved^{8,9} data which require a much larger number of MEs and its rotational dependence. This problem also pertains to the origin of the short component in the time resolved emission after incoherent nanosecond laser excitation.¹⁻⁴ A multitude of weak Coriolis coupled MEs denoted as "grass" were introduced^{10,11} to account for this short component within the intermediate level structure limit.

Coriolis coupling was shown to be the dominant intrastate vibronic coupling leading to vibrational energy redistribution (IVR) in benzene,^{12,13} 9-cyanoanthracene,¹⁴ anthracene,¹⁵ and pyrimidine.¹⁶

It was predicted by the author¹ that the absorption contour of a single rotational transition should be broader in absorption than in excitation, and a low quantum yield absorption should extend to the valley between adjacent rotational transitions where no MEs were observed in the high resolution excitation spectra.¹ The debated grass^{10,11} is real, but in the form of a continuous absorption having a low quantum yield. Here I address the question of the existence and nature of this predicted grass.

Our experimental apparatus^{6,17} was extended to include an excimer pumped dye laser (Lambda Physik EMG 53 MSC + FL2002E) having a 2 GMz spectral resolution using an étalon and pressure tuning. Pulsed planar jets were generated by expansion of seeded argon through a nozzle slit.¹⁸

Direct absorption and fluorescence excitation spectra were simultaneously measured on each laser pulse and then

averaged. The laser repetition rate was doubled as compared with the pulsed nozzle repetition rate and a second differentiation was performed between the timed (with the gas pulses) and untimed laser light pulses. This procedure resulted in a flat, zero wavelength independent absorption base line.

Figure 1 shows the absorption and excitation spectra of pyrazine $S_1(^1B_{3u})$ origin. It is clearly observed that the




FIG. 1. Absorption and fluorescence excitation spectra of pyrazine $S_1(^1B_{3u})$ electronic origin. Spectral resolution is 2 GHz. The two spectra were simultaneously measured using an étalon and pressure tuning. Nozzle temperature was 24 °C, argon backing pressure was 60 Torr, and the laser crossed the pulsed planar jet 16 mm from the nozzle.




FIG. 2. Absorption (full line) and fluorescence excitation (dashed line) spectra of the R_1 rotational transition of pyrazine S_1 ($^1B_{3u}$) electronic origin. Experimental conditions are as in Fig. 1.

emission quantum yield reduces with the rotational quantum number both in P and R branches.⁶ A closer inspection reveals that the absorption signal between rotational transitions is much larger than the corresponding excitation signal.

In Fig. 2, the R_1 transition is magnified. Figure 2 nicely demonstrates that the absorption contour is broader than the excitation contour and a low quantum yield absorption extends to the two sides of this transition. These experimental results constitute and demonstrate two important findings: (a) the dynamics of pyrazine S_1 ($^1B_{3u}$) electronic origin is governed by many more MEs than is deduced from ultrahigh resolution excitation spectra;²⁻⁵ (b) there is a dramatic large variation in the observed emission quantum yield which is much smaller (by a numerical factor of 20!) in the valleys between adjacent rotational transitions.

We are now in the position to state that the problem of the missing states is solved. The missing states are grass, namely a low quantum yield, short time emission, inhomogeneous superposition of homogeneously broadened MEs ($J'', K' \neq 0$).¹ They are characterized by a fast intratriplet IVR. In the excitation spectra they are missing because they are broad and have low (6×10^{-4}) emission quantum yield; thus, they are below the detection sensitivity limit of Kommandeur and co-workers,²⁻⁵ as they are predicted to be $\sim 10^{-3}$ – 10^{-4} of the peak height of MEs ($J'', K'' = 0$) in the ultrahigh resolution spectra.²⁻⁵

Finally we note that these experimental results are in contrast to predictions based on the existence of a high quantum yield near resonance Raman light scattering (NRLS) in the valleys.^{2-4,19} NRLS has 100% quantum yield, much higher than that of the S_1 origin of pyrazine.⁶ NRLS imply a higher excitation spectrum in the valley compared with the absorption spectrum. The experimental results are inconsistent with that prediction.

My deep appreciation is extended to Professor J. Jortner for long, stimulating discussions, and to Professor E.C. Lim for collaboration on the subject of rotational effects. This research was supported by the Israel Academy of Sciences.

¹A. Amirav, Chem. Phys. **108**, 403 (1986).

²K. E. Drabe and J. Kommandeur, in *Excited State*, edited by E. C. Lim and K. K. Innes (Academic, New York, 1987), Vol. 7.

³J. Kommandeur, B. J. van der Meer, and H. Th. Jonkman in *Intramolecular Dynamics*, edited by J. Jortner and B. Pullman (Reidel, Dordrecht, 1982) p. 259.

⁴B. J. van der Meer, H. Th. Jonkman, and J. Kommandeur, Laser Chem. **2**, 77 (1983).

⁵B. J. van der Meer, H. Th. Jonkman, J. Kommandeur, W. L. Meerts, and W. A. Majewski, Chem. Phys. Lett. **92**, 565 (1982).

⁶A. Frad, F. Lahmani, A. Tramer, and C. Tric, J. Chem. Phys. **60**, 4419 (1974).

⁷A. Amirav and J. Jortner, J. Chem. Phys. **84**, 1500 (1986).

⁸Y. Matsumoto, L. H. Spangler, and D. W. Pratt, Laser Chem. **2**, 91 (1983); Chem. Phys. Lett. **98**, 333 (1983).

⁹H. Saigusa and E. C. Lim, J. Chem. Phys. **78**, 91 (1983); Chem. Phys. Lett. **88**, 455 (1982).

¹⁰International Conference on Radiationless Transitions, Newport Beach, California, January 7, 1984, in the panel discussion and private communication. D. W. Pratt claims that the grass exists and J. Kommandeur and L. Meerts denies its existence.

¹¹Y. Matsumoto, L. H. Spangler, and D. W. Pratt, J. Chem. Phys. **80**, 5542 (1984).

¹²E. Riedle, H. J. Neusser, E. W. Schlag, and S. H. Lin, J. Phys. Chem. **88**, 198 (1984).

¹³E. Riedle and H. J. Neusser, J. Chem. Phys. **80**, 4686 (1984).

¹⁴A. Amirav, J. Jortner, M. Terazima, and E. C. Lim, Chem. Phys. Lett. **133**, 179 (1987).

¹⁵A. Amirav, J. Chem. Phys. **86**, 4706 (1987).

¹⁶B. F. Forch and E. C. Lim, Chem. Phys. Lett. **110**, 593 (1984).

¹⁷A. Amirav (in preparation).

¹⁸A. Amirav (in preparation).

¹⁹H. Th. Jonkman, K. E. Drabe, and J. Kommandeur, Chem. Phys. Lett. **116**, 357 (1982).

Photoinitiated $H + CO_2 \rightarrow OH + CO$ reactions: OH distributions and three-body interactions in CO_2H_2S complexes^{a)}

J. Rice,^{b)} G. Hoffmann, and C. Wittig

Department of Chemistry, University of Southern California, Los Angeles, California 90089-0484

(Received 23 September 1987; accepted 4 December 1987)

In previous publications, we reported using weakly bonded complexes as precursors in studies involving oriented and aligned reactants.¹⁻⁵ This method provides a unique environment wherein the V , R wave functions and equilibri-

um geometry determine the initial positions and momenta of the nuclei in the reaction which follows photoexcitation, thereby allowing geometrically constrained entrance channels to be explored. In these studies of precursor geometry