

**Density functional and coupled-cluster study on the  $\text{HNO} \rightarrow \text{HON}$  transition state**

M. E. Alikhani, Christopher E. Dateo, and Timothy J. Lee

Citation: *The Journal of Chemical Physics* **107**, 8208 (1997); doi: 10.1063/1.475123

View online: <http://dx.doi.org/10.1063/1.475123>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/107/19?ver=pdfcov>

Published by the [AIP Publishing](#)

---

**Articles you may be interested in**

[A simple density functional fractional occupation number procedure to determine the low energy transition region of spin-flip reactions](#)

*J. Chem. Phys.* **126**, 124109 (2007); 10.1063/1.2566404

[Computational study of  \$\text{Si}\_2\text{H}\$  and  \$\text{Si}\_2\text{H}\_2\$  species as candidates of interstellar molecules](#)

*AIP Conf. Proc.* **855**, 289 (2006); 10.1063/1.2359567

[Renormalized coupled-cluster methods exploiting left eigenstates of the similarity-transformed Hamiltonian](#)

*J. Chem. Phys.* **123**, 224105 (2005); 10.1063/1.2137318

[Interaction of lead atom with atmospheric hydroxyl radical. An ab initio and density functional theory study of the resulting complexes  \$\text{PbOH}\$  and  \$\text{HPbO}\$](#)

*J. Chem. Phys.* **121**, 7207 (2004); 10.1063/1.1784431

[A coupled-cluster study of the  \$\text{HOBr} \rightarrow \text{HBrO}\$  transition state](#)

*J. Chem. Phys.* **111**, 5780 (1999); 10.1063/1.479875

---



# Density functional and coupled-cluster study on the HNO→HON transition state

M. E. Alikhani<sup>a)</sup>

Laboratoire de Spectrochimie Moléculaire (URA 508), Université P. et M. Curie, Boîte 49, Bâtiment F74,  
4 Place Jussieu Paris Cedex 05, France

Christopher E. Dateo

Thermosciences Institute, MS 230-3, NASA Ames Research Center, Moffett Field, California 94035-1000

Timothy J. Lee

MS230-3, NASA Ames Research Center, Moffett Field, California 94035-1000

(Received 14 August 1997; accepted 14 August 1997)

[S0021-9606(97)03643-X]

Recently, the XNO (X=H, F, Cl) molecules have been investigated using the singles and doubles coupled-cluster method that includes a perturbational estimate of the effect of connected triple excitations, denoted CCSD(T).<sup>1-3</sup> In one of these studies, the transition state structure for the HNO→HON isomerization was reported to have been located,<sup>2</sup> but we have determined that the structure reported therein appears to be the transition state for the HON→H+ON reaction and not the isomerization reaction (note that the other two transition states reported in Ref. 2 for the FNO→FON and ClNO→ClON isomerization reactions are correct). In another previous study,<sup>3</sup> the experimental fundamental vibrational frequencies<sup>4</sup> of HNO are very well reproduced by the CCSD(T) method, which attest to the reliability of the CCSD(T) method for this system.

In this letter, we present the structural, energetic, and vibrational properties of the HNO→HON transition structure (HON(TS)) using density functional theory (DFT) and the CCSD(T) approach. The particular DFT method used here is the Becke3-LYP hybrid method<sup>5</sup> (denoted B3LYP), as implemented within the Gaussian94/DFT quantum chemical package.<sup>6</sup> The 6-311+G(2d) basis set of Pople *et al.*<sup>7</sup> has been used in the DFT calculations. For the CCSD(T) calculations, Dunning's<sup>8</sup> cc-pVTZ and cc-pVQZ basis sets were used, with the quadratic force constants and vibrational frequencies having been determined with the cc-pVTZ basis set. The nitrogen and oxygen core 1s-like molecular orbitals have been required to remain doubly occupied in the CCSD(T) calculations (i.e., the frozen core approximation), and only spherical harmonic components of the *d*, *f*, and *g* functions have been included in the basis.

The DFT calculations have been performed using the Gaussian94/DFT quantum chemical package,<sup>6</sup> while the CCSD(T) calculations have been performed using the TITAN<sup>9</sup> self-consistent field, transformation, and coupled-cluster programs interfaced to the SEWARD<sup>10</sup> integral program.

In Table I, the structural and vibrational properties of HON(TS) are reported. It is important to note that the isomerization reaction is spin forbidden if it is to connect the ground states of HON (<sup>3</sup>A'') and HNO (<sup>1</sup>A').<sup>11</sup> Here we have considered HON, HNO, and HON(TS) in their <sup>1</sup>A' electronic state (closed-shell systems with <*S*<sup>2</sup>>=0). Com-

parison of the CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVQZ geometries shows that the CCSD(T) one-particle basis set limit has nearly been achieved. Indeed, the difference between the CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVQZ HON(TS) geometries is very small. In addition, the  $\mathcal{T}_1$  diagnostic<sup>12</sup> for the CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVQZ calculations (0.036 and 0.034, respectively) indicates that CCSD(T) level of theory should perform well for HNO(TS).

A further inspection of the geometrical parameters shows that the DFT results agree well with the CCSD(T) ones, and are in best agreement with the CCSD(T)/cc-pVQZ quantities although the [DFT-CCSD(T)/cc-pVQZ] difference is quite a bit larger than the [CCSD(T)/cc-pVTZ-CCSD(T)/cc-pVQZ] difference. We note that  $\angle\text{HON} \approx 62^\circ$  is between those for HNO  $\angle\text{HON}(\text{HNO}) \approx 33^\circ$  and HON  $\angle\text{HON}(\text{HON}) \approx 112^\circ$ , as it should be in order to connect the HNO and HON minima. The N-O bond distance varies as NO<HNO<HON<HON(TS), which correlates with the N-O bond force constant variation, i.e., NO>HNO>HON>HON(TS). As shown in Table I, one imaginary frequency has been found, corresponding to the HNO bending mode, which indicates a first-order transition state.

A comparison of the [HON(TS)-HNO] energy differences (i.e., the barrier height from the HNO minimum), shows that there is qualitative agreement among the three levels of theory, with the DFT quantity being 4.4 kcal mol<sup>-1</sup> larger than the CCSD(T)/cc-pVQZ value. Again, the difference between the CCSD(T)/cc-pVTZ and CCSD(T)/cc-pVQZ barrier heights (0.5 kcal mol<sup>-1</sup>) shows that the CCSD(T)/cc-pVQZ value is very close to the CCSD(T) one-particle basis set limit.

The HON/HNO isomerization energy computed at the DFT level of theory (41.2 kcal mol<sup>-1</sup>) is in good agreement with an accurate value determined previously<sup>2</sup> at the CCSD(T)/*spdfg* level of theory (42.5 kcal mol<sup>-1</sup>). Since the heat of formation of HNO is now well known,<sup>13</sup> it is easily seen that HON is only weakly bound relative to the H+ON dissociated products, and therefore that HON(TS) is well above this dissociation channel. Thus the large

TABLE I. Structural and vibrational properties of HON(TS).

Parameters	DFT	CCSD(T)/ cc-pVTZ	CCSD(T)/ cc-pVQZ
$r_{\text{N-O}}$ (Å)	1.322	1.341	1.335
$r_{\text{H-O}}$	1.114	1.101	1.103
$\angle \text{HON}$ (deg.)	62.3	62.0	62.1
$E$ (au) <sup>a</sup>	-130.394 130	-130.182 330	-130.221 227
$\Delta E$ (kcal mol <sup>-1</sup> ) <sup>b</sup>	77.8	72.9	73.4
<b>Harmonic frequencies</b>			
$\omega_1$ (cm <sup>-1</sup> )	2759.9	2820.2	
$\omega_2$	2287.9i	2201.0i	
$\omega_3$	1257.6	1174.3	
$f_{\text{NO}}$ (aJ/Å <sup>2</sup> ) <sup>c,d</sup>	6.90	5.63	

<sup>a</sup>The electronic energy of HNO is -130.518 123, -130.298 456, and -130.338 133 a.u. by DFT, CCSD(T)/cc-pVTZ, and CCSD(T)/cc-pVQZ, respectively.

<sup>b</sup>The barrier height without ZPE corrections.

<sup>c</sup>The DFT value of  $f_{\text{NO}}$  is 16.79, 12.56, and 8.37 aJ/Å<sup>2</sup>, for NO, HNO, and HON, respectively.

<sup>d</sup>For completeness, the CCSD(T)/cc-pVTZ unique quadratic force constants are:  $f_{11}=2.521\ 25$ ,  $f_{21}=1.846\ 36$ ,  $f_{22}=5.630\ 39$ ,  $f_{31}=3.528\ 43$ ,  $f_{32}=-0.653\ 12$ , and  $f_{33}=-0.721\ 36$ , given in units consistent with energy in aJ, bond lengths in Å, and bond angle in radians. Here 1 and 2 label the OH and NO stretches and 3 refers to the HON angle.

HNO→HON barrier height obtained in the present work implies that the isomerization to HON should not occur to any appreciable degree even in rather severe combustion environments.

C.E.D. gratefully acknowledges support from NASA Contract No. NAS2-14031 to ELORET.

<sup>a</sup>Author to whom correspondence should be addressed

<sup>1</sup>T. J. Lee, J. Chem. Phys. **99**, 9783 (1993).

<sup>2</sup>T. J. Lee, Chem. Phys. Lett. **223**, 431 (1994).

<sup>3</sup>C. E. Dateo, T. J. Lee, and D. W. Schwenke, J. Chem. Phys. **101**, 5853 (1994).

<sup>4</sup>J. W. C. Johns and A. R. W. McKellar, J. Chem. Phys. **66**, 1217 (1977); J. W. C. Johns, A. R. W. McKellar, and E. Weinberger, Can. J. Phys. **61**, 1106 (1983).

<sup>5</sup>A. D. Becke, J. Chem. Phys. **98**, 5648 (1993); C. Lee, W. Yang, and R. G. Parr, Phys. Rev. B **37**, 785 (1988); P. J. Stephens, F. J. Devlin, C. F. Chabrowski, and M. J. Frisch, J. Phys. Chem. **98**, 11623 (1994).

<sup>6</sup>M. J. Frisch, G. W. Trucks, H. B. Schlegel, P. M. W. Gill, B. G. Johnson, M. A. Robb, J. R. Cheeseman, T. Keith, G. A. Petersson, J. A. Montgomery, K. Raghavachari, M. A. Al-Laham, V. G. Zakrzewski, J. V. Ortiz, J. B. Foresman, J. Cioslowski, B. B. Stefanov, A. Nanayakkara, M. Challacombe, C. Y. Peng, P. Y. Ayala, W. Chen, M. W. Wong, J. L. Andres, E. S. Replogle, R. Gomperts, R. L. Martin, D. J. Fox, J. S. Binkley, D. J. Defrees, J. Baker, J. P. Stewart, M. Head-Gordon, C. Gonzalez, and J. A. Pople, Gaussian 94 (Revision D.4), Gaussian, Inc., Pittsburgh PA, 1995.

<sup>7</sup>R. Krishnan, J. S. Binkley, R. Seeger, and J. A. Pople, J. Chem. Phys. **72**, 650 (1980); M. J. Frisch, J. A. Pople, and J. S. Binkley, *ibid.* **80**, 3265 (1984).

<sup>8</sup>T. H. Dunning, J. Chem. Phys. **90**, 1007, (1989).

<sup>9</sup>TITAN is a set of electronic structure programs written by T. J. Lee, A. P. Rendell, and J. E. Rice.

<sup>10</sup>R. Lindh, U. Ryu, and B. Liu, J. Chem. Phys. **95**, 5889 (1991).

<sup>11</sup>R. Guadagnini and G. C. Schatz, J. Chem. Phys. **102**, 774 (1995).

<sup>12</sup>T. J. Lee and P. R. Taylor, Int. J. Quantum Chem. Symp. **23**, 199 (1989); T. J. Lee and G. E. Scuseria in *Quantum Mechanical Electronic Structure Calculations with Chemical Accuracy*, edited by S. R. Langhoff (Kluwer Academic, Dordrecht, 1995), pp. 47–108.

<sup>13</sup>T. J. Lee and C. E. Dateo, J. Chem. Phys. **103**, 9110 (1995); R. N. Dixon, *ibid.* **104**, 6905 (1996).