

# Erratum: Nuclear Magnetic Resonance of the Aquated Proton. II. Chloroauric Acid Tetrahydrate. Phase Transitions and Molecular Motion

D. E. O'Reilly, E. M. Peterson, Carl Scheie, and Jack M. Williams

Citation: The Journal of Chemical Physics 56, 3734 (1972); doi: 10.1063/1.1677768

View online: http://dx.doi.org/10.1063/1.1677768

View Table of Contents: http://scitation.aip.org/content/aip/journal/jcp/56/7?ver=pdfcov

Published by the AIP Publishing

#### Articles you may be interested in

H 2 nuclear magnetic resonance study on the molecular motion in cyanoadamantane. II. Orientationally ordered and glassy crystalline phase

J. Chem. Phys. 126, 014701 (2007); 10.1063/1.2409695

Motions of diaquahydrogen ions and phase transitions in dodecatungstophosphoric acid hexahydrate studied by 1 H nuclear magnetic resonance technique

J. Chem. Phys. 113, 7453 (2000); 10.1063/1.1312334

Dynamics of ferroelectric phase transition in vinylidene fluoride/trifluoroethylene (VF2/F3E) copolymers.

II. Proton nuclear magnetic resonance study

J. Chem. Phys. 90, 6740 (1989); 10.1063/1.456293

Nuclear Magnetic Resonance of the Aquated Proton. II. Chloroauric Acid Tetrahydrate. Phase Transitions and Molecular Motion

J. Chem. Phys. 55, 5629 (1971); 10.1063/1.1675731

Nuclear Magnetic Resonance of the Aquated Proton: Hydronium Perchlorate

J. Chem. Phys. 54, 96 (1971); 10.1063/1.1674647



Our model calculations indicated that, for normal-direction kinetic isotope effects ( $k_{\rm light} > k_{\rm heavy}$ ), significant deviations of **r** from the 1.8–2.0 range may occur only when the individual <sup>14</sup>C and <sup>13</sup>C kinetic isotope effects are of unusually low magnitude and/or are associated with easily detectable temperature-dependence anomalies.<sup>4</sup> Although kinetic-isotope-effect temperature-dependence determinations have not been carried out for the oxaloacetic acid decarboxylation, the individual effects are probably too large to be associated with temperature-dependence anomalies. Thus, in this case also, either experimental error or factors not taken into account in the basic statistical-thermodynamic theory must be operative.

\* Supported in part by the U.S. Atomic Energy Commission under Contract AT(30-1)-3663.

† 1971-72 Chaim Weizmann Memorial Fellow, The Weizmann

Institute of Science, Rehovot, Israel.

† Present Address: College of Pharmaceutical Sciences, Columbia

University, New York, N.Y. 10023.

<sup>1</sup> J. Bigeleisen, J. Chem. Phys. 17, 675 (1949); J. Bigeleisen

and M. Wolfsberg, Advan. Chem. Phys. 1, 15 (1958).

<sup>2</sup> A. Fry, *Isotope Effects in Chemical Reactions*, edited by C. J. Collins and N. S. Bowman (Van Nostrand Reinhold, New York,

1970), Chap. 6, p. 369.

<sup>3</sup> A. Wood, Trans. Faraday Soc. 60, 1263 (1964); 62, 1231 (1966).

<sup>4</sup> P. C. Vogel and M. J. Stern, J. Chem. Phys. **54**, 779 (1971).

and B. E. Tabor for the communication which led to the discovery of this error.

### Erratum: Nuclear Magnetic Resonance of the Aquated Proton. II. Chloroauric Acid Tetrahydrate. Phase Transitions and Molecular Motion

[J. Chem. Phys. 55, 5629 (1971)]

D. E. O'REILLY, E. M. PETERSON, CARL SCHEIE, AND JACK M. WILLIAMS

Argonne National Laboratory, Argonne, Illinois 60439

The figures in this paper are given in the correct order but the figure captions are not in the correct order. The following table gives the correspondence between the correct figure numbers and the figure captions as they appeared in print.

Correct numbering 1 2 3 4 5 6 7 8

Printed caption 3 4 8 2 5 1 7 6

# Erratum: Corrections to the Fuoss-Onsager Theory of Electrolytes

[J. Chem. Phys. 53, 2173 (1970)]

T. J. MURPHY

University of Maryland, College Park, Maryland 20742

AND

E. G. D. COHEN

Rockefeller University, New York, New York 10021

In the first line of the expression for  $L_2$  in Eq. (20), the coefficient of the curly brackets should read  $-(1+q^2)/8q$  rather than  $-(1+q^2)/2q$ ; the quantity added to  $L_2$  in making this correction should also be added to the expression for  $\Lambda_2$  in Eq. (23). The comparison with experiment (Tables I and II and Figs. 3-10) is based on the correct formulas and hence is unaffected. The authors would like to thank E. Pitts

# Erratum: Flash Photolytic Production, Reactive Lifetime, and Collisional Quenching of $O_2(b \ ^1\Sigma_a^+, v'=0)$

[I. Chem. Phys. 52, 5502 (1970)]

S. V. FILSETH, A. ZIA, AND K. H. WELGE

Chemistry Department and Centre for Research in Experimental Space Science, York University, Toronto, Ontario, Canada

In Fig. 7 on p. 5506, the correct decay rates are exactly half those indicated on the ordinate. In Fig. 11 on p. 5508, the open circles represent the NH<sub>3</sub> data while the closed circles represent the H<sub>2</sub>O data. In Fig. 12 on p. 5509, the units of pressure should be millitorr. The deactivation rate coefficients which are recorded in Table I are correct as shown except for He and Ne which should read  $\sim 1.0 \times 10^{-17}$  cm<sup>3</sup> molecule<sup>-1</sup>·sec<sup>-1</sup>.