

## Sucrose Inversion in H<sub>2</sub>O – D<sub>2</sub>O

W. H. Hamill and V. K. La Mer

Citation: *The Journal of Chemical Physics* **4**, 294 (1936); doi: 10.1063/1.1749841

View online: <http://dx.doi.org/10.1063/1.1749841>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/4/4?ver=pdfcov>

Published by the AIP Publishing

---

### Articles you may be interested in

[Near ultraviolet photodissociation spectroscopy of Mn+\(H<sub>2</sub>O\) and Mn+\(D<sub>2</sub>O\)](#)

*J. Chem. Phys.* **141**, 204305 (2014); 10.1063/1.4901982

[Predictability of the inverse energy cascade in 2D turbulence](#)

*Phys. Fluids* **13**, 1060 (2001); 10.1063/1.1350877

[Radioluminescence of H<sub>2</sub>O and D<sub>2</sub>O ice](#)

*J. Chem. Phys.* **63**, 2690 (1975); 10.1063/1.431634

[Analysis of D<sub>2</sub>O/H<sub>2</sub>O by the Interferometer](#)

*Rev. Sci. Instrum.* **27**, 872 (1956); 10.1063/1.1715401

[The Mutarotation of Glucose in H<sub>2</sub>O–D<sub>2</sub>O Mixtures](#)

*J. Chem. Phys.* **4**, 144 (1936); 10.1063/1.1749803

---



### Sucrose Inversion in H<sub>2</sub>O–D<sub>2</sub>O

Kinetic studies in H<sub>2</sub>O–D<sub>2</sub>O on the mutarotation of glucose,<sup>1</sup> the decomposition of nitramide<sup>2</sup> and the hydrogen ion catalyzed inversion of sucrose,<sup>3,4</sup> for which the ratios  $k_{D_2O}/k_{H_2O}$  are about 1 : 3.8, 1 : 6, 2.1 : 1, respectively, show that the velocity constant  $k_{H_2O-D_2O}$  does not vary linearly with the fraction of heavy water,  $F_{D_2O} = \Delta s/0.1079$ . The negative departure from linearity, which all exhibit, increases in the above order. Since  $k_{H_2O-D_2O}$  probably depends only indirectly upon the medium it is important to consider other substances which may be present in isotopic modifications and kinetically significant.





FIG. 1.

Since the inversion of sucrose in H<sub>2</sub>O is catalyzed specifically by H<sup>+</sup> it seems likely that a simple relationship will exist between  $k_{H_2O-D_2O}$  and the fraction of heavy hydrogen ion  $F_{D^+} = (D^+)/((D^+) + (H^+))$ . In Table I the first two columns give the data<sup>3,4</sup> reported for this reaction and values of  $F_{D^+}$  in the third column are obtained from the equilibrium constant  $K_1 = 15.7^5$  for the exchange reaction



It will be seen from the graph that a plot of  $k_{H_2O-D_2O}/k_{H_2O}$  vs.  $F_{D^+}$  gives a nearly straight line (crosses), for an extrapolated value of 2.1 for  $k_{D_2O}/k_{H_2O}$ .

This is a specific application of a more general method which has already been applied to explain kinetic measure-

TABLE I.

$\Delta s$ 0.1079	$k_{H_2O-D_2O}$ $k_{H_2O}$	$F_{D^+}$	(H <sub>2</sub> O)	(HDO)	$\frac{(HDO)}{(H_2O)}$	$F_x$	$R_x$	$K_x$
0.000	1.00	0.000						
0.444	1.15	0.169	17.84	25.92	1.45	0.14	6.2	8.9
0.671	1.33	0.332	6.57	23.32	3.54	0.30	2.3	8.3
0.90	1.67	0.677	0.65	9.78	15.0	0.61	0.64	9.6
0.981	2.01	0.921	0.025	2.10	84	0.92	0.087	7.3
1.000	(2.10)	1.000						


ments on the mutarotation of glucose.<sup>1</sup> It may be applied to sucrose inversion without the necessity of identifying any substance, on the simple assumption that  $k_{H_2O-D_2O}/k_{H_2O}$  varies linearly with the fraction  $F_X = DX/(HX + DX)$  of some reactant or intermediate involved in the process of inversion. For each measured  $k_{H_2O-D_2O}/k_{H_2O}$  there is an experimental  $F_{D_2O}$  and a corresponding  $F_X$ , which is obtained on the basis of the assumed linear dependence

$$F_X = [(k_{H_2O-D_2O})/k_{H_2O} - 1]/1.1. \quad (2)$$

For any  $F_{D_2O}$  the corresponding  $R_w = (HDO)/(H_2O)$  is simply calculated, and  $R_X = (HX)/(DX) = 1 - F_X/F_X$ . Since  $R_x R_w = K_x = K_3$  for the reaction



Constancy of  $K_x$  values over the range H<sub>2</sub>O–D<sub>2</sub>O is a satisfactory test for the assumption made, as the last column of the table shows. Although it has been possible to identify  $K_x$  in the present instance with  $K_4$  for the equilibrium (4)



and in the previous example with the isotopic equilibrium constant for tetramethyl glucose, it can obviously be applied independently of any assumptions concerning the identity of HX and without foreknowledge of  $K_x$ . The method is not restricted to kinetic measurements.

W. H. HAMILL

Fordham University,

V. K. LA MER

Columbia University,  
March 14, 1936.

<sup>1</sup> Hamill and La Mer, J. Chem. Phys. 4, 144 (1936).

<sup>2</sup> La Mer and Greenspan, unpublished measurements.

<sup>3</sup> Moelwyn-Hughes and Bonhoeffer, Naturwissenschaften 22, 174 (1934).

<sup>4</sup> Gross, Suess and Steiner, ibid. 22, 662 (1934).

<sup>5</sup> La Mer and Korman, preliminary measurements by E.M.F., unpublished.

<sup>6</sup> From (1) and  $(HDO)^2/(D_2O)(H_2O) = 3.27$ .

### A New Comparison of the Viscosity of D<sub>2</sub>O with that of H<sub>2</sub>O

In an attempt to find a basis for comparison of the viscosity of D<sub>2</sub>O with that of H<sub>2</sub>O that might aid in explaining the observed difference,<sup>1</sup> it was noticed that at 15, 20, 25, 30 and 35°C, respectively, the viscosity of D<sub>2</sub>O is almost the same as that of H<sub>2</sub>O at a temperature 8.5° lower; or if  $\eta_{H_2O} = f(T)$ , then  $\eta_{D_2O} = f(T - B)$  where  $f$  is the same function in both cases, and  $B$  is approximately 8.5°. This would explain the observed decrease in the ratio  $\eta_{D_2O}/\eta_{H_2O}$  with increasing temperature.<sup>2</sup> Several other physical properties of the two waters show somewhat analogous variations, e.g., the melting point of D<sub>2</sub>O is 3.8°C,<sup>3</sup> and the temperature of maximum density for D<sub>2</sub>O is 11.6°C,<sup>4</sup> 7.6° higher than that of H<sub>2</sub>O.

The data are summarized in Table I. Column 1 gives the temperature, column 2 the viscosity of H<sub>2</sub>O in millipoises,<sup>5</sup> column 3 the ratio  $\eta_{D_2O}/\eta_{H_2O}$ ,<sup>2</sup> and column 4 the viscosity

TABLE I. The viscosity of  $D_2O$  compared with that of  $H_2O$ .

T (°C)	$\eta_1$ Vis- cosity <sup>1</sup> of $H_2O$ (milli- poises)	$\eta_2$ Vis- cosity of $D_2O$ (mp)	$\eta_1$ COM- PARE WITH PRE- VIOUS COL- UMN	$(\frac{1}{\eta_1})^{0.5}$ exp.	$(\frac{1}{\eta_1})^{0.5}$ calc.	$(\frac{1}{\eta_2})^{0.5}$ exp.	$(\frac{1}{\eta_2})^{0.5}$ calc.
-4	20.5			7.0	7.0		
0	17.93 <sub>8</sub>			7.47	7.52		
5	15.18 <sub>8</sub>	1.309	19.88	8.11	8.13	7.09	7.09
-3.5			20.1				
10	13.09 <sub>7</sub>	1.286	16.85	8.74	8.74	7.71	7.70
15	11.44 <sub>7</sub>	1.267	14.50	9.35	9.35	8.30	8.31
20	10.08 <sub>7</sub>	1.249	12.60	9.96	9.96	8.91	8.92
25	8.94 <sub>9</sub>	1.232	11.03	10.57	10.57	9.52	9.53
30	8.00 <sub>4</sub>	1.215	9.73	11.18	11.18	10.14	10.14
35	7.20 <sub>8</sub>	1.198	8.64	11.78	11.79	10.76	10.75
26.5			8.64 <sub>8</sub>				

<sup>1</sup> *International Critical Tables*.<sup>2</sup> Lewis and MacDonald, J. Am. Chem. Soc. **55**, 4730 (1933).<sup>3</sup>  $(1/\eta_1)^{0.5} = 0.122T + 7.52$ .<sup>4</sup>  $(1/\eta_2)^{0.5} = 0.122T + 6.48$ .

of  $D_2O$ . In columns 4 and 5 the viscosity of  $D_2O$  at several temperatures is compared with that of  $H_2O$  at a tempera-

ture  $8.5^\circ$  lower. The agreement of the viscosity of  $D_2O$  at temperatures between  $15^\circ$  and  $35^\circ$  with the corresponding  $H_2O$  viscosity is very satisfactory. At the two lowest temperatures,  $5^\circ$  and  $10^\circ$ , respectively,  $D_2O$  has the same viscosity as  $H_2O$  at a temperature  $8.2^\circ$  lower.

To facilitate comparisons,  $(1/\eta)^{0.5}$ , which varies almost linearly with temperature, was plotted as a function of temperature. The values of  $(1/\eta)^{0.5}$  for  $H_2O$  and  $D_2O$ , calculated from the experimental data are shown in columns 6 and 8, respectively. The values calculated from the equations  $(1/\eta_1)^{0.5} = 0.122T + 7.52$ , and  $(1/\eta_2)^{0.5} = 0.122(T - 8.5) + 7.52 = 0.122T + 6.48$  are given in columns 7 and 9. These equations are purely empirical and do not hold at temperatures much above  $35^\circ C$ , but do show the relations  $\eta_1 = f(T)$  and  $\eta_2 = f(T - 8.5)$  for the range  $5^\circ$  to  $35^\circ C$ .

WELDON N. BAKER

Allegheny College,  
Meadville, Pennsylvania,  
March 9, 1936.

<sup>1</sup> Most recent determination,  $\eta_{D_2O}/\eta_{H_2O} = 1.22996$  at  $25^\circ C$ , Jones and Fornwalt, J. Chem Phys. **4**, 30 (1936).

<sup>2</sup> Lewis and MacDonald, J. Am. Chem. Soc. **55**, 4730 (1933).

<sup>3</sup> La Mer and Baker, J. Am. Chem. Soc. **56**, 2641 (1934).

<sup>4</sup> Lewis and MacDonald, J. Am. Chem. Soc. **55**, 3057 (1933).

<sup>5</sup> *International Critical Tables*.