

Exchange Reactions of Protium and Deuterium

R. H. Crist and G. A. Dalin

Citation: The Journal of Chemical Physics 1, 677 (1933); doi: 10.1063/1.1749349

View online: http://dx.doi.org/10.1063/1.1749349

View Table of Contents: http://scitation.aip.org/content/aip/journal/jcp/1/9?ver=pdfcov

Published by the AIP Publishing

Articles you may be interested in

Exchange Reactions with Deuterium II. The Photochemical Exchange Between Deuterium and Hydrogen Chloride

J. Chem. Phys. 6, 345 (1938); 10.1063/1.1750267

Exchange Reactions with Deuterium I. Deuterium and Hydrogen Chloride

J. Chem. Phys. 4, 28 (1936); 10.1063/1.1749742

Mixtures of Deuterium and Protium Oxides as Ideal Solutions

J. Chem. Phys. 3, 197 (1935); 10.1063/1.1749631

The Exchange Reaction of Hydrogen and Deuterium Oxide

J. Chem. Phys. 2, 548 (1934); 10.1063/1.1749525

The Exchange Reaction of Hydrogen and Deuterium Oxide

J. Chem. Phys. 2, 442 (1934); 10.1063/1.1749506



LETTERS TO THE EDITOR

This section will accept reports of new work, provided these are terse and contain few figures, and especially few halftone cuts. The Editorial Board will not hold itself responsible for opinions expressed by the correspondents. Contributions to this section must reach the office of the Managing Editor

not later than the 15th of the month preceding that of the issue in which the letter is to appear. No proof will be sent to the authors. The usual publication charge (\$3.00 per page) will not be made and no reprints will be furnished free.

Exchange Reactions of Protium and Deuterium ¹

Investigations concerning the chemical nature of deuterium have been in progress for some time. The adequate analysis of small amounts of deuterium-rich hydrogen has proved of considerable difficulty, but the interest in the properties of deuterium justifies this preliminary notice.

A sample of water containing one percent deuterium² was circulated with ordinary hydrogen by means of a mercury jet pump through a quartz tube at 800° C for several days. The hydrogen was then freed from the original water by liquid air traps and phosphorus pentoxide. Photographs of the atomic discharge of this hydrogen and of ordinary hydrogen were made on the 21-foot grating. The discharge tube was swept out repeatedly with air and then run with several fresh samples of ordinary hydrogen. The deuterium line $H^2\beta$ was about one-fourth the intensity of the fourth order ghost from the protium $H^1\beta$. The discharge tube was then run with several samples

of the prepared hydrogen. The deuterium $H^2\beta$ of this hydrogen was of greater intensity than the fourth order ghost from $H^1\beta$. No effort was made to estimate the concentration of deuterium as experiments are in progress with more concentrated samples of water and better analytical methods. These experiments deal with the nature of the exchange reported here as well as the atomic metathesis of this and other suitable reactions.

R. H. CRIST G. A. DALIN

Department of Chemistry, Columbia University, New York, New York.

¹ These names have been proposed for the ordinary hydrogen atom and the isotope by Urey, Brickwedde and Murphy, J. Chem. Phys. 1, 512 (1933).

² This was kindly supplied by Professor H. C. Urey.