

Effect of Deuterium Substitution on Color

Samuel H. Maron and Victor K. LaMer

Citation: *The Journal of Chemical Physics* **6**, 299 (1938); doi: 10.1063/1.1750254

View online: <http://dx.doi.org/10.1063/1.1750254>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/6/5?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

[Effect of deuterium substitution on the triplet state dipole moments of substituted benzaldehydes](#)
J. Chem. Phys. **81**, 4237 (1984); 10.1063/1.447456

[Effect of Deuterium Substitution on ESR Linewidth Parameters](#)
J. Chem. Phys. **46**, 2445 (1967); 10.1063/1.1841059

[Effects of Temperature and Deuterium Substitution on the Electron Spin Resonance Spectra of Free Radicals](#)
J. Chem. Phys. **42**, 792 (1965); 10.1063/1.1696012

[Effect of Deuterium Substitution on the Lifetime of the Phosphorescent Triplet State of Naphthalene](#)
J. Chem. Phys. **33**, 302 (1960); 10.1063/1.1731110

[Effect of Deuterium Substitution on the Lifetime of the Phosphorescent Triplet State of Naphthalene](#)
J. Chem. Phys. **32**, 1261 (1960); 10.1063/1.1730887






FIG. 1. The transmission spectra of phenol at various temperatures. Cell thickness approximately 0.01 mm.

and the fact that bands at approximately this wave-length are found in the spectra of a large number of compounds containing OH constitute evidence that this band is connected with the hydroxyl group.

R. ROBERT BRATTAIN

Princeton University,
Princeton, N. J.,
April 12, 1938.

* In collaboration with Research Department, Calco Chemical Co. Inc.

¹ R. B. Barnes, R. R. Brattain and F. Seitz, *Phys. Rev.* **48**, 582 (1935).


² W. Gordy and A. H. Nielsen, *J. Chem. Phys.* **6**, 12 (1938).

Effect of Deuterium Substitution on Color*


Although it is well known that substitution of a deuterium for a hydrogen atom in a compound modifies the physical properties of the compound, still there is no case recorded in the literature where such a substitution changes the color of the compound. Such a color change has been observed by us with nitroethane under special conditions.

When an equivalent quantity of Ba(OD)_2 is added to about 0.02*N* proto-nitroethane in heavy water, a proton is

removed from the nitroethane to yield the ion $\text{CH}_3\text{CH}=\text{NO}_2^-$ according to the equation:




If to this solution be added now an equivalent quantity of D_2SO_4 , a deuteron is reintroduced into the nitroethane to yield:



Throughout these operations the solution remains perfectly colorless.

However, if an equivalent quantity of Ba(OD)_2 be again added, the solution turns *light yellow* very rapidly. In this case the reaction involved is:¹



The color can be discharged and brought back by repeated alternate additions of D_2SO_4 and Ba(OD)_2 .

These same operations with $\text{CH}_3\text{CH}_2\text{NO}_2$ in light water and with Ba(OH)_2 and H_2SO_4 yield no colored solutions.

It will be observed that the only difference between the products in Eq. (1), where there is no color, and Eq. (3), where the yellow color appears, is in the nitroethane ion. In the former the ion is $\text{CH}_3\text{CH}=\text{NO}_2^-$, while in the latter it is $\text{CH}_3\text{CD}=\text{NO}_2^-$, the difference being in the substitution of a deuterium for a hydrogen atom on the carbon α to the nitro group.

From these observations in both H_2O and D_2O it can be concluded that $\text{CH}_3\text{CH}_2\text{NO}_2$, $\text{CH}_3\text{CHDNO}_2$, $\text{CH}_3\text{CD}_2\text{NO}_2$ and $\text{CH}_3\text{CH}=\text{NO}_2^-$ are colorless, while the ion $\text{CH}_3\text{CD}=\text{NO}_2^-$ is colored a light yellow in solutions about 0.02*N* with respect to nitroethane. Preliminary absorption measurements indicate that absorption starts at 5000–5200Å and continues into the ultraviolet.

Because of the possible connection between the development of color and ease of proton and deuteron transfer in conjunction with acid-base catalysis and acid strength ultraviolet and visible absorption spectra are being investigated.

SAMUEL H. MARON
VICTOR K. LAMER

Department of Chemistry,
Columbia University,
New York, N. Y.,
March 29, 1938.

* Publication assisted by Ernest Kempton Adams Fund for Physical Research of Columbia University.

¹ Wynne-Jones, *J. Chem. Phys.* **2**, 381 (1934).