

Synaptic and Extrasynaptic Neurotransmitter Receptors in Glial Precursors' Quest for Identity

SHIBESHIH BELACHEW^{1,2} AND VITTORIO GALLO^{2*}

¹*Center for Cellular and Molecular Neurobiology, University of Liège, Liège, Belgium*

²*Center for Neuroscience Research, Children's Research Institute, Children's National Medical Center, Washington, DC*

KEY WORDS neural progenitors; channels; precursor cells; electrophysiological phenotype; development

ABSTRACT It is widely established that neurotransmitter receptors are expressed in non-neuronal cells, and particularly in neural progenitor cells in the postnatal central nervous system. The functional role of these receptors during development is unclear, but it needs to be revisited now that cells previously considered restricted to glial lineages have been shown to generate neurons. The present review integrates recent advances, to shed new light on how neurotransmitter receptors may, alternatively, serve as excitable mediators of neuron-glia and neuron-neuroblast interactions.

© 2004 Wiley-Liss, Inc.

INTRODUCTION

It is established that neurotransmitters are present and surround neural cells during brain maturation (Miranda-Contreras et al., 1998, 1999, 2000), and functional ligand-gated ion channels (i.e., ionotropic receptors) are expressed in neuronal precursor cells before the establishment of cortical and subcortical synapses (LoTurco et al., 1995; Ma and Barker, 1995; Serafini et al., 1995; Flint et al., 1998; Haydar et al., 2000). Many experimental reports indicate that the interaction between neurotransmitters and their receptors could play an essential role during CNS development, far before they become directly involved in synaptic transmission (Barker et al., 1998; Cameron et al., 1998a; Nguyen et al., 2001). Distinct neurotransmitter-activated ionotropic receptors appear to regulate a variety of processes at specific stages of neuronal development (LoTurco et al., 1995; Haydar et al., 2000; Ben Ari, 2002; Nguyen et al., 2003a). However, the precise role of neurotransmitter receptor-mediated signaling during neural differentiation *in vivo* remains largely unknown, and most loss-of-function studies have failed to disclose obvious developmental defects. Hence, there might be large redundancies between functions of distinct neurotransmitter receptor subunits, as surrogate

mechanisms compensate specific ionotropic receptor deficiencies during neural development.

We and others (Patneau et al., 1994; Steinhäuser and Gallo, 1996; Belachew et al., 1998a, 1998b, 2000; Gallo and Ghiani, 2000; Atluri et al., 2001; Rogers et al., 2001; Nguyen et al., 2001, 2002, 2003a) have recently demonstrated that type A γ -aminobutyric acid (GABA_A), AMPA/kainate, glycine, and nicotinic acetylcholine ionotropic receptors are also expressed in glial progenitors, particularly in oligodendrocyte progenitor cells (OPCs). Until very recently, extrasynaptic neurotransmitter receptors expressed in immature glial progenitor cells have been considered to be transducers of putative neurotransmitter-mediated neuron-glia interactions (Fields and Stevens-Graham, 2002). However, we and other investigators have now demonstrated that OPCs represent a subset of bipotential neuron-oligodendrocyte progenitor cells in specific locations of the postnatal brain (Belachew et al., 2003; Nunes et al.,

*Correspondence to: Vittorio Gallo, Center for Neuroscience Research, Children's Research Institute, Children's National Medical Center, Room 5345, 111 Michigan Ave, NW, Washington, DC 20010. E-mail: vgallo@cnmcresearch.org

Received 29 March 2004; Accepted 28 April 2004

DOI 10.1002/glia.20077

Published online 16 June 2004 in Wiley InterScience (www.interscience.wiley.com).

2003; Aguirre et al., 2004). It is possible that neurotransmitter-receptor interactions may regulate the development of these precursor cells. In the present review, we shall discuss the hypotheses that (1) neurotransmitter receptor expression by postnatal glial progenitors in neurogenic regions might modulate their neurogenic potential, and (2) these receptors represent instructive signaling pathways that may be involved in the regulation of specific stages of neuronal differentiation. Are neurotransmitter receptors constitutively expressed in postnatal glial progenitors that retain the ability to give rise to neurons in the appropriate extracellular environment? We will summarize the available evidence demonstrating neurotransmitter receptor expression in OPCs and in other glial precursors, to discuss whether these receptors might mediate developmental cues that regulate glial proliferation and differentiation, and/or that instruct terminal neuronal maturation.

EXPRESSION OF NEUROTRANSMITTER RECEPTORS AND VOLTAGE-GATED ION CHANNELS IN POSTNATAL NEURAL PRECURSOR CELLS

Current experimental evidence demonstrating that neurotransmitter receptors are expressed in neuron-committed and multipotent or glia-oriented progenitor cells is comprehensively summarized in Tables 1 and 2, respectively. These tables are restricted to ionotropic receptors studied in postnatal neural tissue.

Focus on a Newcomer: Glycine Receptors in Perinatal CNS Precursor Cells

Strychnine-sensitive glycine receptor chloride channels (GlyRs) are members of the nicotinic ligand-gated ion channel family (Le Novere and Changeux, 2001). They mediate postsynaptic inhibition by increasing the chloride permeability of the postsynaptic cell membrane in adult spinal cord, brainstem, and some higher brain regions (Betz, 1991; Kuhse et al., 1995; Schofield et al., 1996). GlyR is a pentameric transmembrane protein composed of glycosylated integral transmembrane α -(48-kDa) and β -subunits (58-kDa) (Schofield et al., 1996). Four α -subunit variants ($\alpha 1$, $\alpha 2$, $\alpha 3$, and $\alpha 4$) (Kuhse et al., 1990; Matzenbach et al., 1994) and one β -subunit variant (Schofield et al., 1996) have been described so far. In newborn rats, the GlyR is $\alpha 2$ -homomeric and differs from the adult stoichiometry, which is classically made of three $\alpha 1$ - and two β -subunits (Langosch et al., 1988). In contrast to the adult brain, where glycine is inhibitory, the activation of GlyR in immature neural cells causes excitation that results from a

depolarized chloride equilibrium potential in these cells during their early developmental period (Ben-Ari, 2002). GlyR-mediated depolarization of immature neurons induces a calcium entry through voltage-gated calcium channels (VGCC) (Reichling et al., 1994; Flint et al., 1998), which appears to modulate cell proliferation, migration, and differentiation in neural precursors (Ben Ari, 2001, 2002; Nguyen et al., 2001) and, later, maturation of fully differentiated neurons bearing inhibitory synapses (Legendre, 2001; Moss and Smart, 2001).

We recently demonstrated that GlyRs are expressed in three distinct types of early postnatal neural precursor cells *in vitro* (Belachew et al., 1998b, 2000; Nguyen et al., 2002, 2004). Functional GlyRs have been described in newborn rat astrocytes, oligodendrocytes, and OPCs in spinal cord slices (Pastor et al., 1995; Kirchhoff et al., 1996; Belachew et al., 1998a). During oligodendroglial lineage progression, GlyR expression ($\alpha 2$ - and β -subunits) culminates at the OPC stage and decreases thereafter (Belachew et al., 1998b). We demonstrated that glycine-induced chloride currents recorded from newborn rat cortical oligosphere-derived OPCs are mediated by GlyRs that seem to be not only pharmacologically, but also structurally distinct from the homologous neuronal receptor (Belachew et al., 1998b). Furthermore, glycine application on cultured OPCs induces a rise of $[Ca^{2+}]_i$ via the opening of VGCC. This effect is mediated by a depolarization related to the concurrent activation of GlyR and Na^+ -dependent glycine transporters GLYT1 and/or GLYT2 (Belachew et al., 2000).

Using the classical neurosphere culture model from early postnatal striatum (Reynolds et al., 1992; Reynolds and Weiss, 1992), we also demonstrated the expression of GlyR $\alpha 1$ -, $\alpha 2$ - and β genes, and could detect GlyR α subunit proteins in multipotent nestin-expressing stem/progenitor cells (Nguyen et al., 2002). Consistent with these findings, we observed that, in these multipotent progenitor cells, GlyR subunits forms functional chloride channels gated by glycine and blocked by the GlyR antagonists strychnine and picrotoxin (Nguyen et al., 2002).

Similarly, immunocytochemical analysis showed that GlyRs are expressed in PSA-NCAM⁺ neurogenic progenitors from perinatal rat striatum both *in vitro* and *in situ* (Nguyen et al., 2004). Glycine triggers inward strychnine-sensitive currents in most of these neurogenic striatal precursors (Nguyen et al., 2004). Finally, neurogenic PSA-NCAM⁺ progenitors from the perinatal striatum express GlyRs that display electrophysiological characteristics (Hill coefficient and EC₅₀ mean effective concentration) intermediate between nestin-expressing neural stem cells (Nguyen et al., 2002) and mature interneurons (Sergeeva, 1998; Sergeeva and Haas, 2001).

Consistent with the role of GlyR $\alpha 2$ subunit during retinal development (Young and Cepko, 2004), GlyRs and their agonists, in concert with other neurotransmitter receptors and voltage-gated ion channels, may

TABLE 1. *Ionotropic Neurotransmitter Receptors Expressed in Neuron-Committed Precursor Cells*

| Age | Receptor | Progenitor cell phenotype | Model | Species | Results and Effects of NT | References |
|----------------------|---------------------|---|----------------------------------|---------|--|---|
| GABA | | | | | | |
| P6–P8 | GABA _A | Cerebellar granule cell progenitors | In vitro cell culture | Rat | MAPK-dependent ↑ proliferation | Fiszman et al., 1999 |
| P6–P8 | GABA _A | Immature cerebellar granule cells | In vitro cell culture | Rat | Ca ²⁺ -dependent ↓ of neurite outgrowth | Borodinsky et al., 2003 |
| P0–P3 | GABA _A | Striatal PSA-NCAM ⁺ precursors | In vitro cell culture and slices | Rat | MAPK- and Ca ²⁺ -dependent ↓ of proliferation | Nguyen et al., 2003a |
| P13–P70 | GABA _A | anterior SVZ-RMS-olfactory bulb neuronal precursors | Ex vivo tissue slices | Mice | Neuronal progenitors of the SVZ/RMS/OB contain GABA and are depolarized by GABA-mediated activation of first extrasynaptic and then later synaptic GABA _A receptors | Belluzzi et al., 2003; Carleton et al., 2003; Gallo and Haydar, 2003; Wang et al., 2003 |
| P0–P1 | GABA _A | Anterior SVZ neuronal precursors | In vitro cell culture | Rat | SVZa neuronal precursor cells are GABAergic and express functional GABA _A receptors | Stewart et al., 2002 |
| Glutamate | | | | | | |
| P0–P3 | ND | Striatal PSA-NCAM ⁺ precursors | In vitro cell culture | Rat | Glutamate induces inward currents | Nguyen et al., 2003b |
| P0–adult | NMDA | Dentate gyrus progenitor cells | In vivo | Rat | Blockade of NMDA receptor increases cell death and [³ H]thymidine incorporation, particularly in granule cells | Cameron et al., 1995, Cameron et al., 1998b; Gould et al., 1994 |
| P7–P11 | NMDA | Immature cerebellar granule cells | In vitro and in vivo | Rat | Activation of NMDA receptor ↑ survival | Balazs et al., 1988a, 1988b; Monti et al., 2002; Monti and Contestabile, 2000 |
| P70 | AMPA | Tangentially migrating olfactory bulb neuroblasts | Ex vivo tissue slices | Mice | Expression of first extrasynaptic and later synaptic AMPA receptors | Carleton et al., 2003 |
| P70 | NMDA | Radially migrating olfactory bulb neuroblasts | Ex vivo tissue slices | Mice | Expression of extrasynaptic functional NMDA receptors | Carleton et al., 2003 |
| Glycine | | | | | | |
| P0–P3 | GlyR | Striatal PSA-NCAM ⁺ precursors | In vitro cell culture | Rat | Functional GlyR expressed in these neonatal neuronal precursor cells | Nguyen et al., 2004 |
| Acetylcholine | | | | | | |
| P0–P3 | ND | Striatal PSA-NCAM ⁺ precursors | In vitro cell culture | Rat | Acetylcholine induces inward currents | Nguyen et al., 2003b |
| P1–P3 | Nicotinic (α7) AChR | Olfactory bulb neuronal precursors | In vitro cell culture | Rat | Activation of nAChR ↑ neurite outgrowth | Coronas et al., 2000 |
| P5 | Nicotinic (α3) AChR | Cerebellar external granule layer (EGL) neuronal precursors | In vitro cell culture | Mouse | Activation of nAChR ↑ proliferation and survival in EGL neuroblasts | Opanashuk et al., 2001 |

also control the development of postnatal neural precursors, i.e. (1) cell cycle kinetics in different types of immature progenitors; (2) cell fate choice and differentiation from nestin-expressing stem cells, (3) neurogenesis from PSA-NCAM⁺ neuronal precursors, and (4) maturation of OPCs into myelinating cells.

OPCs Express Not Only Neurotransmitter Receptors, But Also Voltage-Gated Potassium Channels

Similar to the widespread array of neurotransmitter receptors that are displayed by CNS precursor cells, voltage-gated ion channels, and in particular voltage-

TABLE 2. *Iontropic Neurotransmitter Receptors Expressed in Multipotent and Glia-Oriented Precursor Cells*

| Age | Receptor | Progenitor cell phenotype | Model | Species | Results and effects of NT | References |
|----------------------|--------------------|--|-----------------------------------|---------------------|--|---|
| GABA | | | | | | |
| Adult | GABA _A | Astrocytes and astrocyte precursors from and hippocampus | In vitro tissue slices | Rat | GABA induces depolarizing chloride-mediated inward currents | MacVicar et al., 1989 |
| P0-P3 | GABA _A | Spinal cord and cerebral OPCs | In vitro cell culture and slices | Rat and mokerteruse | GABA induces depolarizing chloride-mediated inward currents and \uparrow $[Ca^{2+}]_i$ | Belachew et al., 1998a; Gilbert et al., 1984; Kettenmann et al., 1984b, Kirchhoff and Kettenmann, 1992, Pastor et al., 1995 |
| Glutamate | | | | | | |
| P1 | AMPA and kainate | Whole brain and hippocampus | In vitro cell culture and slices | Rat | Ca^{2+} -permeable functional AMPA/kainate receptor expression; their activation induces cell depolarization | Bowman and Kimeberg, 1984; Jabs et al., 1994; Kettenmann et al., 1984a; Porter and McCarthy, 1995; Sontheimer et al., 1988; Steinhauser et al., 1994 |
| Juvenile adult | AMPA and kainate | Hippocampus | In vitro tissue slices | Mouse | Glial inwardly rectifying K ⁺ channels (Kir) are rapidly blocked upon activation of AMPA/kainate-type glutamate receptors, most probably due a receptor-mediated influx of Na ⁺ , which plugs the channels from the intracellular side | Jabs et al., 1994; Robert and Magistretti, 1997; Schroder et al., 2002 |
| P1 | AMPA and kainate | Cortical astrocytes and mixed glial cells | In vitro cell culture and slices | Rat | Glutamate receptor-mediated regulation of immediate early gene expression (e.g., c-fos, c-jun, jun-B) | Mack et al., 1994; McNaughton and Hunt, 1992 |
| P1 | AMPA and kainate | Cortical and optic nerve OPCs | In vitro cell culture | Rat | Functional AMPA and kainate-specific subtypes of glutamate receptors are expressed by OPCs | Barres et al., 1990; Gallo et al., 1994a, 1994b; Jensen and Chiu 1993; Patneau et al., 1994 |
| P1 | AMPA and kainate | Cortical OPCs | In vitro cell culture and slices | Rat | AMPA/kainate receptor activation: (1) triggers Ca^{2+} -dependent early gene expression; (2) induces a blockade of voltage-dependent K ⁺ channels (Kv) that is mediated by depolarization and Na ⁺ entry; (3) inhibits proliferation at G1/S transition by regulation of cyclinE/cdk2 complex activity | Belachew et al., 2002; Chittajallu et al., 2002; Gallo et al., 1996; Gallo and Ghiani, 2000; Ghiani et al., 1999a, 1999b; Ghiani and Gallo, 2001; Knutson et al., 1997; Liu and Almazan, 1995; Pende et al., 1994, 1997 |
| P1 | AMPA | Cortical OPCs | In vitro cell culture | Rat | Overactivation of AMPA receptors causes apoptosis in oligodendrocyte progenitors through mechanisms involving Ca^{2+} influx, depletion of glutathione, and activation of JNK, calpain, and caspase-3 | Liu et al., 2002 |
| Glycine | | | | | | |
| P0-P3 | GlyR | Nestin-expressing, striatal stem cells | In vitro cell culture | Rat | Functional GlyR expressed in these stem cells | Nguyen et al., 2002 |
| P0-P3 | GlyR | Cortical OPCs | In vitro cell culture | Rat | Activation of functional GlyR induces depolarizing chloride-mediated inward currents and \uparrow $[Ca^{2+}]_i$ | Belachew et al., 1998, 2000 |
| Acetylcholine | | | | | | |
| P1 | Nicotinic (7) AChR | Hippocampal astrocytes | In vitro cell culture | Rat | Activation of nAChR \uparrow intracellular calcium concentration | Sharma and Vijayaraghavan, 2001 |
| P0-P3 | Nicotinic AChR | Cortical OPCs | In vitro cell culture | Rat | Functional nAChR expressed by cultured OPCs | Belachew et al., 1998a |
| P1 | Nicotinic AChR | OPCs from corpus callosum | Ex vivo and in vitro cell culture | Rat | Activation of nAChR \uparrow intracellular calcium concentration; expression of 3, 4, 5, 7, 2, and 4 subunits. | Rogers et al., 2001 |

dependent K^+ (Kv) channels, are expressed not only in neuron-committed, but also in multipotent and glia-oriented precursor cells. The prototypical Kv channels have been divided into four families (Kv1–Kv4), based on sequence homology to single gene orthologues in *Drosophila* (name given in italics): Kv1 (*Shaker*), Kv2 (*Shab*), Kv3 (*Shaw*), and Kv4 (*Shal*) (Trimmer and Rhodes, 2004). Importantly, there is a direct correlation between the expression of delayed outward-rectifying voltage-gated K^+ -currents (I_k) and the proliferative state of different types of cells (DeCoursey et al., 1984; Sobko et al., 1998). Furthermore, pharmacological block of such potassium channels interferes with cell proliferation (DeCoursey et al., 1984; Sontheimer et al., 1989; Barres et al., 1990; Berger et al., 1991; Kettenmann et al., 1991; Dubois and Rouzaille-Dubois, 1993; Wonderlin and Strobl, 1996; Knutson et al., 1997; MacFarlane and Sontheimer, 1997; Sobko et al., 1998). Proliferative OPCs express large I_k , whereas nondividing OPCs and mature oligodendrocytes display much smaller outward currents (Knutson et al., 1997). These large I_k currents occur in G1 phase of the OPC cell cycle and have been correlated to an RNA synthesis-dependent selective increase of expression of K^+ channel subunit proteins Kv1.3 and Kv1.5 (Gallo et al., 1996; Knutson et al., 1997; Ghiani et al., 1999b; Chittajallu et al., 2002).

The inhibition of outward K^+ channels by the blocker tetraethylammonium or by cell membrane depolarization with the Na^+ channel opener veratridine inhibits OPC proliferation (Knutson et al., 1997; Ghiani et al., 1999b). The use of subunit-specific pharmacological blockers first suggested that outward K^+ channels containing the Kv1.3 subunit play a critical role in cell cycle progression in proliferating OPCs (Chittajallu et al., 2002). We recently performed gain-of-function experiments providing evidence that overexpression of Kv1.3–1.4 increases OPC proliferation in the absence of mitogen, whereas Kv1.6 overexpression prevents mitogen-induced OPC cell cycle progression (Vautier et al., 2004). Kv1.3–1.6 subunit overexpression does not interfere with oligodendrocyte differentiation (Vautier et al., 2004). This study demonstrates that the activity of potassium channels comprising distinct Kv1 subunit proteins not only correlates with specific stages of oligodendrocyte lineage progression, but directly controls oligodendroglial proliferation in mitogenic or in growth factor-free conditions.

ESTABLISHED ROLE OF NEUROTRANSMITTER RECEPTORS IN POSTNATAL CNS PRECURSOR CELL BIOLOGY **AMPA/Kainate Receptors: A Matter of Cell Cycle or Cell Death**

In the postnatal and adult brain, glutamate is released from neurons (either synaptically or axonally) (Chiu and Kriegler, 1994; Bergles and Jahr, 1997) and

reaches effective concentrations that can activate different classes of glutamate receptors (GluRs) in glia. This event can cause (1) modulation of transmitter uptake, thereby affecting termination of synaptic transmission (Vernadakis, 1996; Bergles and Jahr, 1997); (2) modulation of K^+ conductance, with consequences on the extracellular ionic environment (Muller et al., 1992; Vernadakis, 1996); and (3) release of neuroactive substances from glia that can feedback and modulate synaptic transmission in neurons (Araque et al., 1999).

Glutamate receptor function is not limited to mature glial cells, because during late embryonic and early postnatal development, astrocytes and oligodendrocytes arise from CNS precursor cells of the subventricular zone that also express glutamate-activated channels (LoTurco et al., 1995; Yuan et al., 1998). Glutamate activates ionotropic GluRs in OPCs in purified cultures or in tissue slices, to inhibit their proliferation and prevent differentiation (Gallo et al., 1996; Yuan et al., 1998). Furthermore, glutamate inhibits SVZ progenitor proliferation in cortical slices (Haydar et al., 2000). The antiproliferative effects of glutamate are Ca^{2+} -independent and arise from an increase in intracellular Na^+ and subsequent block of outward K^+ currents (Gallo et al., 1996; Knutson et al., 1997). Agonists specific for AMPA/kainate receptors or blockers of outward K^+ channels cause a reversible G1 cell cycle arrest in OPCs. This is associated with accumulation of the cyclin-dependent kinase inhibitors p27^{Kip1} and p21^{CIP1}, selective decrease in cdk2 activity, and impairment of cyclin E-cdk2 complex formation (Ghiani et al., 1999a, 1999b; Ghiani and Gallo, 2001). Overexpression of cdk2 partially prevents G1 arrest induced by AMPA/kainate agonists or K^+ channel blockers (Belachew et al., 2002), indicating that regulation of cdk2 activity causally underlies cell cycle arrest. It still needs to be determined whether GluR activation interferes with cell cycle progression in CNS precursor cells from adult brain, and whether gray matter vs. white matter NG2⁺ precursor cells express distinct receptor and channel subtypes with area-specific functions (see below).

Several laboratories have also demonstrated that AMPA/kainate receptor activation triggers excitotoxic cell death in cells of the oligodendroglial lineage, although most of this work was performed using cultures of postmitotic O4⁺/GalC⁺ pre-oligodendrocytes (Oka et al., 1993; Yoshioka et al., 1995; Matute et al., 1997; Sanchez-Gomez and Matute, 1999; Alberdi et al., 2002; Sanchez-Gomez et al., 2003). In contrast, it appears that both proliferating OPCs and mature myelinating oligodendrocytes are not sensitive to excitotoxic insults (Gallo et al., 1996; Rosenberg et al., 2003).

The mechanism of AMPA/kainate receptor-mediated excitotoxic lesions in white matter oligodendroglial cells in vivo remains a subject of intense debate (Matute, 1998; McDonald et al., 1998; Follett et al., 2000; Li and Stys, 2000; Matute et al., 2001; Pitt et al., 2000; Smith et al., 2000; Tekkok and Goldberg, 2001), since surrounding microglia (Noda et al., 2000) and astro-

cytes (Jabs et al., 1994; Verkhratsky and Steinhauser, 2000; Schroder et al., 2002; Matthias et al., 2003) also express such receptors, and therefore may be directly or indirectly involved in the excitotoxic cascade. It is important to mention that the only *in situ* electrophysiological demonstration of functional AMPA/kainate receptor expression in adult cells identified as OPCs (based on NG2 expression; see below) was not provided in the white matter, but in the hippocampus (Bergles et al., 2000). It is also of interest to note that the demonstration of excitotoxicity in adult white matter oligodendroglial cells is not by itself evidence for the presence of AMPA/kainate receptors in these cells. This issue can be better defined in future experiments by recording astrocytes and oligodendroglial cells from the adult white matter of transgenic mice expressing reporter genes under cell-specific promoters (Zhuo et al., 1997; Belachew et al., 2001; Nolte et al., 2001; Mallon et al., 2002; Yuan et al., 2002). In particular, in the case of demyelinating disorders, this approach will reveal whether excitotoxic GluRs overactivation, which occurs primarily or secondarily to auto-immune activation, triggers direct oligodendroglial and axonal damages, and/or modulates astroglial/microglial components of the inflammatory reaction, which would then cause the final toxic effects (Pitt et al., 2000; Smith et al., 2000; Matute et al., 2001).

GABA_A Receptors

In the embryonic brain, the neurotransmitter γ -aminobutyric acid (GABA) partially prevents bFGF-induced proliferation of VZ progenitor cells (Antonopoulos et al., 1997). However, GABA was shown to increase the number of proliferating cells and to accelerate cell cycle kinetics in VZ progenitors analyzed in organotypic cultures under basal conditions (LoTurco et al., 1995; Haydar et al., 2000). In contrast, GABA decreases SVZ progenitor cell proliferation (Haydar et al., 2000). It appears, therefore, that neural progenitors in the SVZ display a differential physiological response to GABA, as compared with VZ progenitors, indicating that neural progenitors can rapidly modulate their sensitivity to the inhibitory neurotransmitter as development proceeds.

Consistent with these prior prenatal studies, GABA was found to inhibit cell proliferation in the postnatal SVZ. We recently analyzed the expression and function of GABA_A receptors (GABA_ARs) in SVZ/striatal neural precursor cells during postnatal development. We developed an isolation procedure that allowed us to purify in a buoyant density gradient proliferative neural precursor cells from early postnatal rat striatum and adjacent SVZ, based on their isopycnic properties. These cells expressed the polysialylated form of the neural cell adhesion molecule (PSA-NCAM) (Nguyen et al., 2003a). We investigated whether GABA might control the proliferation of these postnatal PSA-NCAM⁺ precursors and reported that (1) epidermal growth factor

(EGF)-responsive PSA-NCAM⁺ precursors generate neurospheres mostly committed to a neuronal fate; (2) postnatal striatal/SVZ PSA-NCAM⁺ precursors express functional GABA_ARs and the GABA-synthesizing enzymes glutamate decarboxylase (GAD)-65 and GAD-67 *in vitro* and *in situ*; (3) proliferation of PSA-NCAM⁺ precursors is inhibited by GABA produced by these cells under the control of EGF; and (4) GABA_AR-dependent inhibition of PSA-NCAM⁺ cell proliferation is mediated by a complex intracellular signaling involving an increase of intracellular calcium concentration by opening of voltage-gated calcium channels and inhibition of the mitogen-activated protein kinase (MAPK) pathway (Nguyen et al., 2003a). The demonstration that early postnatal PSA-NCAM⁺ striatal/SVZ neuronal precursors synthesized and released GABA suggests that an autocrine/paracrine GABA/GABA_AR-mediated mechanism might regulate EGF-driven proliferation of these progenitors.

Similarly, receptor activation by exogenous application of GABA_A agonists to perinatal OPCs in cultured organotypic cerebellar slices also inhibited proliferation of these precursor cells (Yuan et al., 1998). However, treatment of the slices with the GABA_A receptor antagonist bicuculline did not cause any effect (Yuan et al., 1998), suggesting that, at least in cerebellum, endogenous activation of GABA_A receptors is not a major mechanism of control of cell cycle progression in postnatal OPCs *in vivo*.

Altogether, these findings strongly indicate that GABA_A receptors may be directly involved in the regulation of proliferation and differentiation of perinatal precursor cells. Like early postnatal cells, also adult (P13–P70) neuronal progenitors of the SVZ/RMS/olfactory bulb pathway synthesize GABA and are depolarized by activation of first extrasynaptic and later synaptic GABA_A receptors (Carleton et al., 2003; Gallo and Haydar, 2003; Wang et al., 2003). Thus, GABA_A receptor-mediated signaling might be part of the complex array of extrinsic signals that regulate adult neurogenesis.

SYNAPTIC INPUTS ON NEURAL PROGENITORS: NEURON-GLIA OR NEURON-NEUROBLAST SIGNALING? NG2-Expressing Progenitor

In recent studies, Bergles and colleagues (Bergles et al., 2000; Lin and Bergles, 2003) have suggested that OPCs receive glutamatergic and GABAergic synaptic inputs from interneuronal collaterals in the CA1 region of the postnatal/adult hippocampus. The criteria used for identifying the cells analyzed electrophysiologically as OPCs was the immunohistochemical detection of the chondroitin sulfate proteoglycan NG2. NG2 has been widely used as an excellent marker of OPCs *in vivo* by many investigators (Levine et al., 2001). Although it cannot be disputed that NG2⁺ cells are capable of differentiating into myelinating oligodendrocytes in




Fig. 1. Schematic representation of white matter- and hippocampal-specific fate of NG2-expressing cells in the postnatal brain. In white matter regions, NG2⁺ cells generate myelinating oligodendrocytes through the formation of O4⁺ pre-oligodendrocytes and postmitotic O1⁺ oligodendrocytes. In the hippocampus, immature CNP-EGFP⁺ neurons express TOAD-64 (Cameron and McKay, 2001), are NG2⁺, and display high levels of EGFP, whereas differentiated NeuN⁺/CNP-EGFP⁺ neurons are NG2-negative and exhibit lower levels of EGFP fluorescence (Belachew et al., 2003). These findings indicated that NG2 proteoglycan expression and CNP gene promoter activity, as assessed by EGFP fluorescence intensity, are inversely

correlated with the progression of neuronal differentiation in vivo. Hence, altogether these results delineated a possible developmental link between a defined class of adult progenitor cells expressing NG2 chondroitin proteoglycan and the CNP gene, and newborn postnatal GABAergic hippocampal neurons. Left: we proposed that NG2⁺ cells can escape their restricted glial commitment in the hippocampus, and undergo the following developmental changes: NG2⁺/CNP-EGFP⁺⁺⁺ → TOAD-64⁺/NG2⁺/CNP-EGFP⁺⁺ → GAD-67⁺/NeuN⁺/NG2-negative/CNPEGFP⁺. It remains to be established whether this differentiation potential of NG2⁺ cells may be observed in other CNS location, particularly in the subventricular zone.

white matter regions, it is debatable whether this developmental program occurs for all CNS cells that express NG2, particularly in those brain regions that display relatively less oligodendrogenesis, but more neurogenesis in the postnatal and adult brain. We and others have now demonstrated that precursor cells which were thought to be restricted to an oligodendroglial fate, i.e., so-called OPCs, were able to generate neurons under certain conditions both in vitro and in vivo (Belachew et al., 2003; Kondo and Raff, 2000; Nunes et al., 2003) (Fig. 1). These precursor cells express NG2 or A2B5, and the 2',3'-cyclic nucleotide 3'-phosphodiesterase (CNP) gene (Belachew et al., 2003; Nunes et al., 2003). Furthermore, although neural stem cells are thought to possess an undifferentiated phenotype, cells with properties of the astroglial lineage, i.e., radial glia and astrocytes, act as neural stem cells in the embryonic and adult CNS across species (Doetsch et al., 1999; Malatesta et al., 2000; Noctor et al., 2001; Doetsch, 2003).

Importantly, these studies have also led to a redefinition of the neural stem cell nomenclature, to in-

clude glial progenitors as stem cells capable of generating neurons (Goldman, 2003). Based on these recent findings, it is premature to define functional synaptic connections onto all NG2-expressing cells in the hippocampus as directly involved in neuron-oligodendroglial signaling, until it is proved that all NG2⁺ cells that receive excitatory and inhibitory synapses in this particular region are restricted to an oligodendroglial fate. The synapses analyzed and characterized by Bergles and colleagues might, in fact, be synaptic contacts between mature resident interneurons and early postmitotic progenitors of adult GABAergic neurons. As previously described in the olfactory bulb (Carleton et al., 2003), this would also imply that the establishment of synaptic inputs occurs before progenitors for GABAergic neurons are able to propagate action potential spikes during the course of interneuron renewal in the hippocampus (Liu et al., 2003).

Unlike in the developing brain, where neurogenesis usually precedes the maturation of neural circuits, in

the adult newborn neurons must integrate into a pre-existing neuronal network.

During SVZ-RMS-OB (subventricular zone-rostromigratory stream-olfactory bulb) neurogenesis, it was shown that tangentially migrating neuroblasts in the RMS first express extrasynaptic GABA_A receptors and then AMPARs (Carleton et al., 2003). Again, this differs from the developing mammalian brain, where most often NMDARs are first detected, and functional AMPAR recruitment is induced by NMDAR activation (Durand et al., 1996). Adult-generated OB granule neurons receive synaptic inputs shortly after completing their radial migration, as they begin to develop their dendritic arbor, and GABAergic synaptic events appear before AMPAR-mediated excitatory postsynaptic potentials (Carleton et al., 2003). In sharp contrast with prenatal development, OB granule neurons developing in the adult brain receive both GABAergic and glutamatergic synaptic inputs before they acquire the property of propagating action potentials (Carleton et al., 2003). Unlike newborn OB granule neurons, however, in adult-generated OB periglomerular neurons the maturation of voltage-dependent sodium currents appears to precede the formation of synaptic contacts (Belluzzi et al., 2003). Thus, the sequential events that result in the functional integration of newly formed adult neurons appear to be age-specific. It remains to be elucidated whether GABA and glutamate, through the activation of extrasynaptic and synaptic GABA_A and AMPA receptors, are instructive signals required for subsequent OB neuronal maturation, including the emergence of voltage-gated channels.

In contrast to SVZ-RMS-OB neurogenesis, the migration pathway and the time course of electrophysiological maturation of adult-born hippocampal neurons have not yet been precisely defined. Since it is widely accepted that neurogenesis in the adult intact brain is restricted to the SVZ and the dentate gyrus, it could be speculated that the NG2⁺ cells analyzed by Bergles and colleagues in the adult CA1 may be immature neuronal precursors with the potential to become GABAergic interneurons (Gage, 2000; Lie et al., 2004). Recordings from NG2⁺ cells were obtained exclusively in the stratum radiatum region of CA1 in the hippocampus of young (P6–16) and adult (>P33) rats (Bergles et al., 2000; Lin and Bergles, 2003). To our knowledge, there is no evidence that neurogenesis occurs in this region of the normal hippocampus, a fact confirmed in a recent report (Liu et al., 2003). Furthermore, it is unlikely that synaptically integrated neuronal precursor cells could still be migratory. Thus, if CA1 NG2⁺ cells indeed represent neuronal precursor cells, they could be arrested in their differentiation process to become mature neurons, but still available to be recruited and terminally differentiate, as observed under pathological conditions such as ischemia (Nakatomi et al., 2002).

In conclusion, although it still needs to be established what percentage of NG2⁺ cells in the hippocampus give rise to neurons and oligodendrocytes, respec-

tively, a more detailed cellular and developmental characterization of the cell population studied is necessary to unequivocally conclude that they represent OPCs.

Recent data also emphasize that we can no longer rely on a single antigenic marker to identify a specific cell phenotype. In particular, there is now a crucial need for revisiting the definition of an OPC with region- and context-specific functional and antigenic criteria, with respect to a cell that would be restricted to a glial vs. a neuronal fate. The precise characterization of the developmental potential of NG2⁺ cells in distinct brain regions will go beyond issues of mere nomenclature, to define whether glutamate- and GABA-mediated signaling on these progenitors is an exclusive form of neuron-oligodendroglia communication or if a proportion is in fact representative of neuron to neuroblast signaling. Besides the white matter, where it is well defined that NG2⁺ cells do express AMPA/kainate receptors that are likely involved in neuron-oligodendroglial interactions (Yuan et al., 2002; Chittajallu et al., 2002), the physiological role of GluR-mediated signaling for NG2⁺ cells located in the olfactory bulb, hippocampus, striatum, and cerebral cortex is by far more elusive. Further studies that will analyze a variety of antigenic markers in combination with detailed electrophysiological analysis will elucidate the importance of synaptic and extrasynaptic GluRs in NG2⁺ cells located in gray matter regions of the perinatal and adult brain.

Astrocytes

The term neuroglia, or “nerve glue,” was coined in 1859 by Rudolph Virchow, who conceived it as an inactive connective tissue in the CNS. Many years later, we now fully recognize that glial cells play many active roles in brain function, metabolism and development. In the glial kingdom, astrocytes display sheet-like or lamellate processes enwrapping the brain surface, capillaries, neuronal cell bodies, dendrites, and synapses. The astrocytic processes contain a broad range of ion channels and neurotransmitter receptors that can trigger [Ca²⁺]_i elevation in response to synaptic activity (Araque et al., 2001; Haydon, 2001). Calcium signaling in astrocytes is thought to influence glial morphology, function and gene expression, and in turn to feed back onto neurons (Verkhratsky and Kettenmann, 1996). In particular, much attention has been paid to GluRs on astrocytes as possible mediators of activity-dependent neuron-glia interplay (Steinhauser and Gallo, 1996), in terms of both neuron-to-glia and glia-to-neuron signaling (Gallo and Ghiani, 2000; Haydon, 2001). However, until recently there was little experimental evidence that supported this idea or investigated possible mechanisms involved.

Cerebellar Bergmann glial cells ensheath the Purkinje cell dendrites and synapses, and express high levels of Ca²⁺-permeable glutamate receptors (GluRs) (Burna-

shev et al., 1992; Muller et al., 1992). The GluR2 subunit controls Ca^{2+} -permeability of AMPA receptors, i.e., receptors containing this subunit are impermeable to Ca^{2+} ions (Hollmann et al., 1991; Verdoorn et al., 1991). Suppression of Ca^{2+} permeability by virus-mediated GluR2 transfer in Bergmann glia has revealed that GluR-mediated Ca^{2+} signaling is essential to maintain both anatomical and functional connections between Bergmann glial cells and the glutamatergic synapses (Iino et al., 2001). The finding that a structural interaction between Bergmann glial processes and synapses on Purkinje cells is mediated by the glial Ca^{2+} -permeable GluRs further suggests that glia-synapse interaction could undergo structural modifications in response to synaptic activity (Gallo and Chittajallu, 2001; Watanabe, 2002). Astrocytes also express high-affinity glutamate transporters, and particularly GLAST and GLT-1, which play a major role in controlling the rapid clearance of glutamate released from neurons from the synaptic cleft (Chaudhry et al., 1995; Rothstein et al., 1996; Tanaka et al., 1997; Olie et al., 2001). This process is central in terminating the effects of released neurotransmitters, and contributes to the fine tuning of excitatory synaptic transmission in the mature brain.

Both astrocytic glutamate uptake and homeostasis of appropriate connections between Bergmann glia and Purkinje cell synapses that depend on GluR2-containing AMPA receptors represent unequivocal examples of neuron-glia signaling and interaction. In future studies, it will be important to assess similar functional and structural relationships between astrocytes and neighboring neurons or neuronal precursors in brain regions such as the hippocampus, where GFAP-expressing cells (defined as astrocytes) have been suggested to give rise to neurons (Doetsch, 2003; Seri et al., 2001). It is striking that, together with recent studies showing that astrocytes can regulate synapse formation and synaptic transmission (Mauch et al., 2001; Nagler et al., 2001; Ullian et al., 2001), other reports have demonstrated that hippocampal astrocytes can instruct neurogenesis from neural stem cells and/or from committed neuronal precursor cells in the adult brain (Song et al., 2002a,b). It will be important to establish to what extent neurotransmitter-mediated signaling may be involved in these specific developmental properties of astrocytes in neurogenic regions.

CONCLUSIONS

The studies reviewed in this article indicate that expression of ligand- and voltage-gated channels is not an exclusive property of mature neural cells fully integrated in synaptic circuits. The exciting finding that receptors and channels are found in neural progenitors is somewhat counterbalanced by the issue of defining the identity of these precursor cells and their fate, i.e., the type of differentiated progeny they can generate in distinct brain regions. The availability of new lineage

tracing approaches and transgenic mouse strains will greatly help in associating specific precursor cells to a defined electrophysiological phenotype, and to a particular progeny. Similar approaches, combined with gain- and loss-of-function studies will also contribute to clarifying the functional role of receptors and channels at specific stages of neural development.

ACKNOWLEDGMENTS

S.B. is a Research Associate of the Belgian National Fund for Scientific Research (F.N.R.S.).

REFERENCES

- Aguirre AA, Chittajallu R, Belachew S, Gallo V. 2004. NG2-expressing cells in the subventricular zone are Type C-like cells and contribute to interneuron generation in the postnatal hippocampus. *J Cell Biol* 165:575–589.
- Alberdi E, Sanchez-Gomez MV, Marino A, Matute C. 2002. Ca^{2+} influx through AMPA or kainate receptors alone is sufficient to initiate excitotoxicity in cultured oligodendrocytes. *Neurobiol Dis* 9:234–243.
- Antonopoulos J, Pappas IS, Parnavelas JG. 1997. Activation of the GABA_A receptor inhibits the proliferative effects of bFGF in cortical progenitor cells. *Eur J Neurosci* 9:291–298.
- Araque A, Parpura V, Sanzgiri RP, Haydon PG. 1999. Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci* 22: 208–215.
- Araque A, Carmignoto G, Haydon PG. 2001. Dynamic signaling between astrocytes and neurons. *Annu Rev Physiol* 63:795–813.
- Atluri P, Fleck MW, Shen Q, Mah SJ, Stadfelt D, Barnes W, Goderie SK, Temple S, Schneider AS. 2001. Functional nicotinic acetylcholine receptor expression in stem and progenitor cells of the early embryonic mouse cerebral cortex. *Dev Biol* 240:143–156.
- Balazs R, Hack N, Jorgensen OS. 1988a. Stimulation of the N-methyl-D-aspartate receptor has a trophic effect on differentiating cerebellar granule cells. *Neurosci Lett* 87:80–86.
- Balazs R, Jorgensen OS, Hack N. 1988b. N-methyl-D-aspartate promotes the survival of cerebellar granule cells in culture. *Neuroscience* 27:437–451.
- Barker JL, Behar T, Li YX, Liu QY, Ma W, Maric D, Maric I, Schaffner AE, Serafini R, Smith SV, Somogyi R, Vautrin JY, Wen XL, Xian H. 1998. GABAergic cells and signals in CNS development. *Perspect Dev Neurobiol* 5:305–322.
- Barres BA, Koroshetz WJ, Swartz KJ, Chun LL, Corey DP. 1990. Ion channel expression by white matter glia: the O-2A glial progenitor cell. *Neuron* 4:507–524.
- Belachew S, Malgrange B, Rigo JM, Rogister B, Coucke P, Mazy-Servais C, Moonen G. 1998a. Developmental regulation of neurotrophin-induced responses in cultured oligodendroglia. *NeuroReport* 9:973–980.
- Belachew S, Rogister B, Rigo JM, Malgrange B, Mazy-Servais C, Xhaufaire G, Coucke P, Moonen G. 1998b. Cultured oligodendrocyte progenitors derived from cerebral cortex express a glycine receptor which is pharmacologically distinct from the neuronal isoform. *Eur J Neurosci* 10:3556–3564.
- Belachew S, Malgrange B, Rigo JM, Rogister B, Leprince P, Hans G, Nguyen L, Moonen G. 2000. Glycine triggers an intracellular calcium influx in oligodendrocyte progenitor cells which is mediated by the activation of both the ionotropic glycine receptor and Na^{+} -dependent transporters. *Eur J Neurosci* 12:1924–1930.
- Belachew S, Yuan X, Gallo V. 2001. Unraveling oligodendrocyte origin and function by cell-specific transgenesis. *Dev Neurosci* 23:287–298.
- Belachew S, Aguirre AA, Wang H, Vautier F, Yuan X, Anderson S, Kirby M, Gallo V. 2002. Cyclin-dependent kinase-2 controls oligodendrocyte progenitor cell cycle progression and is downregulated in adult oligodendrocyte progenitors. *J Neurosci* 22:8553–8562.
- Belachew S, Chittajallu R, Aguirre AA, Yuan X, Kirby M, Anderson S, Gallo V. 2003. Postnatal NG2 proteoglycan-expressing progenitor cells are intrinsically multipotent and generate functional neurons. *J Cell Biol* 161:169–186.

- Belluzzi O, Benedusi M, Ackman J, LoTurco JJ. 2003. Electrophysiological differentiation of new neurons in the olfactory bulb. *J Neurosci* 23:10411–10418.
- Ben Ari Y. 2001. Developing networks play a similar melody. *Trends Neurosci* 24:353–360.
- Ben Ari Y. 2002. Excitatory actions of GABA during development: the nature of the nurture. *Nat Rev Neurosci* 3:728–739.
- Berger T, Schnitzer J, Kettenmann H. 1991. Developmental changes in the membrane current pattern, K^+ buffer capacity, and morphology of glial cells in the corpus callosum slice. *J Neurosci* 11:3008–3024.
- Bergles DE, Jahr CE. 1997. Synaptic activation of glutamate transporters in hippocampal astrocytes. *Neuron* 19:1297–1308.
- Bergles DE, Roberts JD, Somogyi P, Jahr CE. 2000. Glutamatergic synapses on oligodendrocyte precursor cells in the hippocampus. *Nature* 405:187–191.
- Betz H. 1991. Glycine receptors: heterogeneous and widespread in the mammalian brain. *Trends Neurosci* 14:458–461.
- Borodinsky LN, O'Leary D, Neale JH, Vicini S, Coso OA, Fiszman ML. 2003. GABA-induced neurite outgrowth of cerebellar granule cells is mediated by $GABA_A$ receptor activation, calcium influx and CaMKII and erk1/2 pathways. *J Neurochem* 84:1411–1420.
- Bowman CL, Kimelberg HK. 1984. Excitatory amino acids directly depolarize rat brain astrocytes in primary culture. *Nature* 311:656–659.
- Burnashev N, Khodorova A, Jonas P, Helm PJ, Wisden W, Monyer H, Seeburg PH, Sakmann B. 1992. Calcium-permeable AMPA-kainate receptors in fusiform cerebellar glial cells. *Science* 256:1566–1570.
- Cameron HA, McKay RD. 2001. Adult neurogenesis produces a large pool of new granule cells in the dentate gyrus. *J Comp Neurol* 435:406–417.
- Cameron HA, McEwen BS, Gould E. 1995. Regulation of adult neurogenesis by excitatory input and NMDA receptor activation in the dentate gyrus. *J Neurosci* 15:4687–4692.
- Cameron HA, Hazel TG, McKay RD. 1998a. Regulation of neurogenesis by growth factors and neurotransmitters. *J Neurobiol* 36:287–306.
- Cameron HA, Tanapat P, Gould E. 1998b. Adrenal steroids and N-methyl-D-aspartate receptor activation regulate neurogenesis in the dentate gyrus of adult rats through a common pathway. *Neuroscience* 82:349–354.
- Carleton A, Petreanu LT, Lansford R, Alvarez-Buylla A, Lledo PM. 2003. Becoming a new neuron in the adult olfactory bulb. *Nat Neurosci* 6:507–518.
- Chaudhry FA, Lehre KP, van Lookeren CM, Ottersen OP, Danbolt NC, Storm-Mathisen J. 1995. Glutamate transporters in glial plasma membranes: highly differentiated localizations revealed by quantitative ultrastructural immunocytochemistry. *Neuron* 15:711–720.
- Chittajallu R, Chen Y, Wang H, Yuan X, Ghiani CA, Heckman T, McBain CJ, Gallo V. 2002. Regulation of Kv1 subunit expression in oligodendrocyte progenitor cells and their role in G_i/S phase progression of the cell cycle. *Proc Natl Acad Sci USA* 99:2350–2355.
- Chiu SY, Kriegler S. 1994. Neurotransmitter-mediated signaling between axons and glial cells. *Glia* 11:191–200.
- Coronas V, Durand M, Chabot JG, Jourdan F, Quirion R. 2000. Acetylcholine induces neuritic outgrowth in rat primary olfactory bulb cultures. *Neuroscience* 98:213–219.
- DeCoursey TE, Chandy KG, Gupta S, Cahalan MD. 1984. Voltage-gated K^+ channels in human T lymphocytes: a role in mitogenesis? *Nature* 307:465–468.
- Doetsch F. 2003. The glial identity of neural stem cells. *Nat Neurosci* 6:1127–1134.
- Doetsch F, Caille I, Lim DA, Garcia-Verdugo JM, Alvarez-Buylla A. 1999. Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. *Cell* 97:703–716.
- Dubois JM, Rouzaire-Dubois B. 1993. Role of potassium channels in mitogenesis. *Prog Biophys Mol Biol* 59:1–21.
- Durand GM, Kovalchuk Y, Konnerth A. 1996. Long-term potentiation and functional synapse induction in developing hippocampus. *Nature* 381:71–75.
- Fields RD, Stevens-Graham B. 2002. New insights into neuron-glia communication. *Science* 298:556–562.
- Fiszman ML, Borodinsky LN, Neale JH. 1999. GABA induces proliferation of immature cerebellar granule cells grown in vitro. *Brain Res Dev Brain Res* 115:1–8.
- Flint AC, Liu X, Kriegstein AR. 1998. Nonsynaptic glycine receptor activation during early neocortical development. *Neuron* 20:43–53.
- Follett PL, Rosenberg PA, Volpe JJ, Jensen FE. 2000. NBQX attenuates excitotoxic injury in developing white matter. *J Neurosci* 20:9235–9241.
- Gage FH. 2000. Mammalian neural stem cells. *Science* 287:1433–1438.
- Gallo V, Chittajallu R. 2001. Unwrapping glial cells from the synapse: what lies inside? *Science* 292:872–873.
- Gallo V, Ghiani CA. 2000. Glutamate receptors in glia: new cells, new inputs and new functions. *Trends Pharmacol Sci* 21:252–258.
- Gallo V, Haydar T. 2003. GABA: exciting again in its own right. *J Physiol* 550:665.
- Gallo V, Patneau DK, Mayer ML, Vaccarino FM. 1994a. Excitatory amino acid receptors in glial progenitor cells: molecular and functional properties. *Glia* 11:94–101.
- Gallo V, Wright P, McKinnon RD. 1994b. Expression and regulation of a glutamate receptor subunit by bFGF in oligodendrocyte progenitors. *Glia* 10:149–153.
- Gallo V, Zhou JM, McBain CJ, Wright P, Knutson PL, Armstrong RC. 1996. Oligodendrocyte progenitor cell proliferation and lineage progression are regulated by glutamate receptor-mediated K^+ channel block. *J Neurosci* 16:2659–2670.
- Ghiani CA, Gallo V. 2001. Inhibition of cyclin E-cyclin-dependent kinase 2 complex formation and activity is associated with cell cycle arrest and withdrawal in oligodendrocyte progenitor cells. *J Neurosci* 21:1274–1282.
- Ghiani CA, Eisen AM, Yuan XQ, DePinho RA, McBain CJ, Gallo V. 1999a. Neurotransmitter receptor activation triggers $p27^{Kip1}$ and $p21^{CIP1}$ accumulation and G_1 cell cycle arrest in oligodendrocyte progenitors. *Development* 126:1077–1090.
- Ghiani CA, Yuan XQ, Eisen AM, Knutson PL, DePinho RA, McBain CJ, Gallo V. 1999b. Voltage-activated K^+ channels and membrane depolarization regulate accumulation of the cyclin-dependent kinase inhibitors $p27^{Kip1}$ and $p21^{CIP1}$ in glial progenitor cells. *J Neurosci* 19:5380–5392.
- Gilbert P, Kettenmann H, Schachner M. 1984. Gamma-aminobutyric acid directly depolarizes cultured oligodendrocytes. *J Neurosci* 4:561–569.
- Goldman S. 2003. Glia as neural progenitor cells. *Trends Neurosci* 26:590–596.
- Gould E, Cameron HA, McEwen BS. 1994. Blockade of NMDA receptors increases cell death and birth in the developing rat dentate gyrus. *J Comp Neurol* 340:551–565.
- Haydar TF, Wang F, Schwartz ML, Rakic P. 2000. Differential modulation of proliferation in the neocortical ventricular and subventricular zones. *J Neurosci* 20:5764–5774.
- Haydon PG. 2001. GLIA: listening and talking to the synapse. *Nat Rev Neurosci* 2:185–193.
- Hollmann M, Hartley M, Heinemann S. 1991. Ca^{2+} permeability of KA-AMPA-gated glutamate receptor channels depends on subunit composition. *Science* 252:851–853.
- Iino M, Goto K, Kakegawa W, Okado H, Sudo M, Ishiuchi S, Miwa A, Takayasu Y, Saito I, Tsuzuki K, Ozawa S. 2001. Glia-synapse interaction through Ca^{2+} -permeable AMPA receptors in Bergmann glia. *Science* 292:926–929.
- Jabs R, Kirchhoff F, Kettenmann H, Steinhäuser C. 1994. Kainate activates Ca^{2+} -permeable glutamate receptors and blocks voltage-gated K^+ currents in glial cells of mouse hippocampal slices. *Pflügers Arch* 426:310–319.
- Jensen AM, Chiu SY. 1993. Expression of glutamate receptor genes in white matter: developing and adult rat optic nerve. *J Neurosci* 13:1664–1675.
- Kettenmann H, Backus KH, Schachner M. 1984a. Aspartate, glutamate and gamma-aminobutyric acid depolarize cultured astrocytes. *Neurosci Lett* 52:25–29.
- Kettenmann H, Gilbert P, Schachner M. 1984b. Depolarization of cultured oligodendrocytes by glutamate and GABA. *Neurosci Lett* 47:271–276.
- Kettenmann H, Blankenfeld GV, Trotter J. 1991. Physiological properties of oligodendrocytes during development. *Ann NY Acad Sci* 633:64–77.
- Kirchhoff F, Kettenmann H. 1992. GABA triggers a $[Ca^{2+}]_i$ increase in murine precursor cells of the oligodendrocyte lineage. *Eur J Neurosci* 4:1049–1058.
- Kirchhoff F, Mulhardt C, Pastor A, Becker CM, Kettenmann H. 1996. Expression of glycine receptor subunits in glial cells of the rat spinal cord. *J Neurochem* 66:1383–1390.
- Knutson P, Ghiani CA, Zhou JM, Gallo V, McBain CJ. 1997. K^+ channel expression and cell proliferation are regulated by intracellular sodium and membrane depolarization in oligodendrocyte progenitor cells. *J Neurosci* 17:2669–2682.
- Kondo T, Raff M. 2000. Oligodendrocyte precursor cells reprogrammed to become multipotential CNS stem cells. *Science* 289:1754–1757.

- Kuhse J, Schmieden V, Betz H. 1990. Identification and functional expression of a novel ligand binding subunit of the inhibitory glycine receptor. *J Biol Chem* 265:22317–22320.
- Kuhse J, Betz H, Kirsch J. 1995. The inhibitory glycine receptor: architecture, synaptic localization and molecular pathology of a postsynaptic ion-channel complex. *Curr Opin Neurobiol* 5:318–323.
- Langosch D, Thomas L, Betz H. 1988. Conserved quaternary structure of ligand-gated ion channels: the postsynaptic glycine receptor is a pentamer. *Proc Natl Acad Sci USA* 85:7394–7398.
- Legendre P. 2001. The glycinergic inhibitory synapse. *Cell Mol Life Sci* 58:760–793.
- Le Novère N, Changeux JP. 2001. The ligand gated ion channel database: an example of a sequence database in neuroscience. *Philos Trans R Soc Lond B Biol Sci* 356:1121–1130.
- Levine JM, Reynolds R, Fawcett JW. 2001. The oligodendrocyte precursor cell in health and disease. *Trends Neurosci* 24:39–47.
- Li S, Stys PK. 2000. Mechanisms of ionotropic glutamate receptor-mediated excitotoxicity in isolated spinal cord white matter. *J Neurosci* 20:1190–1198.
- Lie DC, Song H, Colamarino SA, Ming GL, Gage FH. 2004. Neurogenesis in the adult brain: new strategies for central nervous system diseases. *Annu Rev Pharmacol Toxicol* 44:399–421.
- Lin SC, Bergles DE. 2003. Synaptic signaling between GABAergic interneurons and oligodendrocyte precursor cells in the hippocampus. *Nat Neurosci* 10:1038/nn1162.
- Liu HN, Almazan G. 1995. Glutamate induces c-fos proto-oncogene expression and inhibits proliferation in oligodendrocyte progenitors: receptor characterization. *Eur J Neurosci* 7:2355–2363.
- Liu HN, Giasson BI, Mushynski WE, Almazan G. 2002. AMPA receptor-mediated toxicity in oligodendrocyte progenitors involves free radical generation and activation of JNK, calpain and caspase 3. *J Neurochem* 82:398–409.
- Liu S, Wang J, Zhu D, Fu Y, Lukowiak K, Lu YM. 2003. Generation of functional inhibitory neurons in the adult rat hippocampus. *J Neurosci* 23:732–736.
- LoTurco JJ, Owens DF, Heath MJ, Davis MB, Kriegstein AR. 1995. GABA and glutamate depolarize cortical progenitor cells and inhibit DNA synthesis. *Neuron* 15:1287–1298.
- Ma W, Barker JL. 1995. Complementary expressions of transcripts encoding GAD67 and GABA_A receptor alpha 4, beta 1, and gamma 1 subunits in the proliferative zone of the embryonic rat central nervous system. *J Neurosci* 15:2547–2560.
- MacFarlane SN, Sontheimer H. 1997. Electrophysiological changes that accompany reactive gliosis in vitro. *J Neurosci* 17:7316–7329.
- Mack KJ, Kriegler S, Chang S, Chiu SY. 1994. Transcription factor expression is induced by axonal stimulation and glutamate in the glia of the developing optic nerve. *Brain Res Mol Brain Res* 23:73–80.
- MacVicar BA, Tse FW, Crichton SA, Kettenmann H. 1989. GABA-activated Cl[−] channels in astrocytes of hippocampal slices. *J Neurosci* 9:3577–3583.
- Malatesta P, Hartfuss E, Gotz M. 2000. Isolation of radial glial cells by fluorescent-activated cell sorting reveals a neuronal lineage. *Development* 127:5253–5263.
- Mallon BS, Shick HE, Kidd GJ, Macklin WB. 2002. Proteolipid promoter activity distinguishes two populations of NG2-positive cells throughout neonatal cortical development. *J Neurosci* 22:876–885.
- Matthias K, Kirchhoff F, Seifert G, Huttmann K, Matyash M, Kettenmann H, Steinhauser C. 2003. Segregated expression of AMPA-type glutamate receptors and glutamate transporters defines distinct astrocyte populations in the mouse hippocampus. *J Neurosci* 23:1750–1758.
- Matute C. 1998. Characteristics of acute and chronic kainate excitotoxic damage to the optic nerve. *Proc Natl Acad Sci USA* 95:10229–10234.
- Matute C, Sanchez-Gomez MV, Martinez-Millan L, Miledi R. 1997. Glutamate receptor-mediated toxicity in optic nerve oligodendrocytes. *Proc Natl Acad Sci USA* 94:8830–8835.
- Matute C, Alberdi E, Domercq M, Perez-Cerda F, Perez-Samartin A, Sanchez-Gomez MV. 2001. The link between excitotoxic oligodendroglial death and demyelinating diseases. *Trends Neurosci* 24:224–230.
- Matzenbach B, Maulet Y, Sefton L, Courtier B, Avner P, Guenet JL, Betz H. 1994. Structural analysis of mouse glycine receptor alpha subunit genes. Identification and chromosomal localization of a novel variant. *J Biol Chem* 269:2607–2612.
- Mauch DH, Nagler K, Schumacher S, Goritz C, Muller EC, Otto A, Pfrieger FW. 2001. CNS synaptogenesis promoted by glia-derived cholesterol. *Science* 294:1354–1357.
- McDonald JW, Althomsons SP, Hyrc KL, Choi DW, Goldberg MP. 1998. Oligodendrocytes from forebrain are highly vulnerable to AMPA/kainate receptor-mediated excitotoxicity. *Nat Med* 4:291–297.
- McNaughton LA, Hunt SP. 1992. Regulation of gene expression in astrocytes by excitatory amino acids. *Brain Res Mol Brain Res* 16:261–266.
- Miranda-Contreras L, Mendoza-Briceno RV, Palacios-Pru EL. 1998. Levels of monoamine and amino acid neurotransmitters in the developing male mouse hypothalamus and in histotypic hypothalamic cultures. *Int J Dev Neurosci* 16:403–412.
- Miranda-Contreras L, Benitez-Diaz PR, Mendoza-Briceno RV, Delgado-Saez MC, Palacios-Pru EL. 1999. Levels of amino acid neurotransmitters during mouse cerebellar neurogenesis and in histotypic cerebellar cultures. *Dev Neurosci* 21:147–158.
- Miranda-Contreras L, Ramirez-Martens LM, Benitez-Diaz PR, Pena-Contreras ZC, Mendoza-Briceno RV, Palacios-Pru EL. 2000. Levels of amino acid neurotransmitters during mouse olfactory bulb neurogenesis and in histotypic olfactory bulb cultures. *Int J Dev Neurosci* 18:83–91.
- Monti B, Contestabile A. 2000. Blockade of the NMDA receptor increases developmental apoptotic elimination of granule neurons and activates caspases in the rat cerebellum. *Eur J Neurosci* 12:3117–3123.
- Monti B, Marri L, Contestabile A. 2002. NMDA receptor-dependent CREB activation in survival of cerebellar granule cells during in vivo and in vitro development. *Eur J Neurosci* 16:1490–1498.
- Moss SJ, Smart TG. 2001. Constructing inhibitory synapses. *Nat Rev Neurosci* 2:240–250.
- Muller T, Moller T, Berger T, Schnitzer J, Kettenmann H. 1992. Calcium entry through kainate receptors and resulting potassium-channel blockade in Bergmann glial cells. *Science* 256:1563–1566.
- Nagler K, Mauch DH, Pfrieger FW. 2001. Glia-derived signals induce synapse formation in neurones of the rat central nervous system. *J Physiol* 533:665–679.
- Nakatomi H, Kuriu T, Okabe S, Yamamoto S, Hatano O, Kawahara N, Tamura A, Kirino T, Nakafuku M. 2002. Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors. *Cell* 110:429–441.
- Nguyen L, Rigo JM, Rocher V, Belachew S, Malgrange B, Rogister B, Leprince P, Moonen G. 2001. Neurotransmitters as early signals for central nervous system development. *Cell Tissue Res* 305:187–202.
- Nguyen L, Malgrange B, Belachew S, Rogister B, Rocher V, Moonen G, Rigo JM. 2002. Functional glycine receptors are expressed by postnatal nestin-positive neural stem/progenitor cells. *Eur J Neurosci* 15:1299–1305.
- Nguyen L, Malgrange B, Breuskin I, Bettendorff L, Moonen G, Belachew S, Rigo JM. 2003a. Autocrine/paracrine activation of the GABA_A receptor inhibits the proliferation of neurogenic polysialylated neural cell adhesion molecule-positive (PSA-NCAM⁺) precursor cells from postnatal striatum. *J Neurosci* 23:3278–3294.
- Nguyen L, Rigo JM, Malgrange B, Moonen G, Belachew S. 2003b. Untangling the functional potential of PSA-NCAM-expressing cells in CNS development and brain repair strategies. *Curr Med Chem* 10:2185–2196.
- Nguyen L, Malgrange B, Breuskin I, Lallemand F, Hans G, Moonen G, Belachew S, Rigo JM. 2004. Striatal PSA-NCAM⁺ precursor cells from the newborn rat express functional glycine receptors. *NeuroReport* 15:583–587.
- Noctor SC, Flint AC, Weissman TA, Dammerman RS, Kriegstein AR. 2001. Neurons derived from radial glial cells establish radial units in neocortex. *Nature* 409:714–720.
- Noda M, Nakanishi H, Nabekura J, Akaike N. 2000. AMPA-kainate subtypes of glutamate receptor in rat cerebral microglia. *J Neurosci* 20:251–258.
- Nolte C, Matyash M, Pivneva T, Schipke CG, Ohlemeyer C, Hanisch UK, Kirchhoff F, Kettenmann H. 2001. GFAP promoter-controlled EGFP-expressing transgenic mice: a tool to visualize astrocytes and astrogliosis in living brain tissue. *Glia* 33:72–86.
- Nunes MC, Roy NS, Keyoung HM, Goodman RR, McKhann G, Jiang L, Kang J, Nedergaard M, Goldman SA. 2003. Identification and isolation of multipotential neural progenitor cells from the subcortical white matter of the adult human brain. *Nat Med* 9:439–447.
- Oka A, Belliveau MJ, Rosenberg PA, Volpe JJ. 1993. Vulnerability of oligodendroglia to glutamate: pharmacology, mechanisms, and prevention. *J Neurosci* 13:1441–1453.
- Oliet SH, Piet R, Poulain DA. 2001. Control of glutamate clearance and synaptic efficacy by glial coverage of neurons. *Science* 292:923–926.
- Opanashuk LA, Pauly JR, Hauser KF. 2001. Effect of nicotine on cerebellar granule neuron development. *Eur J Neurosci* 13:48–56.
- Pastor A, Chvatal A, Sykova E, Kettenmann H. 1995. Glycine- and GABA-activated currents in identified glial cells of the developing rat spinal cord slice. *Eur J Neurosci* 7:1188–1198.

- Patneau DK, Wright PW, Winters C, Mayer ML, Gallo V. 1994. Glial cells of the oligodendrocyte lineage express both kainate- and AMPA-preferring subtypes of glutamate receptor. *Neuron* 12:357–371.
- Pende M, Holtzclaw LA, Curtis JL, Russell JT, Gallo V. 1994. Glutamate regulates intracellular calcium and gene expression in oligodendrocyte progenitors through the activation of DL-alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors. *Proc Natl Acad Sci USA* 91:3215–3219.
- Pende M, Fisher TL, Simpson PB, Russell JT, Blenis J, Gallo V. 1997. Neurotransmitter- and growth factor-induced cAMP response element binding protein phosphorylation in glial cell progenitors: role of calcium ions, protein kinase C, and mitogen-activated protein kinase/ribosomal S6 kinase pathway. *J Neurosci* 17:1291–1301.
- Pitt D, Werner P, Raine CS. 2000. Glutamate excitotoxicity in a model of multiple sclerosis. *Nat Med* 6:67–70.
- Porter JT, McCarthy KD. 1995. GFAP-positive hippocampal astrocytes in situ respond to glutamatergic neuroleptoligands with increases in $[Ca^{2+}]_i$. *Glia* 13:101–112.
- Reichling DB, Kyzioz A, Wang J, MacDermott AB. 1994. Mechanisms of GABA and glycine depolarization-induced calcium transients in rat dorsal horn neurons. *J Physiol* 476:411–421.
- Reynolds BA, Weiss S. 1992. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science* 255:1707–1710.
- Reynolds BA, Tetzlaff W, Weiss S. 1992. A multipotent EGF-responsive striatal embryonic progenitor cell produces neurons and astrocytes. *J Neurosci* 12:4565–4574.
- Robert A, Magistretti PJ. 1997. AMPA/kainate receptor activation blocks K^+ currents via internal Na^+ increase in mouse cultured stellate astrocytes. *Glia* 20:38–50.
- Rogers SW, Gregori NZ, Carlson N, Gahring LC, Noble M. 2001. Neuronal nicotinic acetylcholine receptor expression by O2A/oligodendrocyte progenitor cells. *Glia* 33:306–313.
- Rosenberg PA, Dai W, Gan XD, Ali S, Fu J, Back SA, Sanchez RM, Segal MM, Follett PL, Jensen FE, Volpe JJ. 2003. Mature myelin basic protein-expressing oligodendrocytes are insensitive to kainate toxicity. *J Neurosci Res* 71:237–245.
- Rothstein JD, Dykes-Hoberg M, Pardo CA, Bristol LA, Jin L, Kuncl RW, Kanai Y, Hediger MA, Wang Y, Schielke JP, Welty DF. 1996. Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* 16:675–686.
- Sanchez-Gomez MV, Matute C. 1999. AMPA and kainate receptors each mediate excitotoxicity in oligodendroglial cultures. *Neurobiol Dis* 6:475–485.
- Sanchez-Gomez MV, Alberdi E, Ibarretxe G, Torre I, Matute C. 2003. Caspase-dependent and caspase-independent oligodendrocyte death mediated by AMPA and kainate receptors. *J Neurosci* 23:9519–9528.
- Schofield PR, Lynch JW, Rajendra S, Pierce KD, Handford CA, Barry PH. 1996. Molecular and genetic insights into ligand binding and signal transduction at the inhibitory glycine receptor. *Cold Spring Harb Symp Quant Biol* 61:333–342.
- Schroder W, Seifert G, Huttmann K, Hinterkeuser S, Steinhauser C. 2002. AMPA receptor-mediated modulation of inward rectifier K^+ channels in astrocytes of mouse hippocampus. *Mol Cell Neurosci* 19:447–458.
- Serafini R, Valev AY, Barker JL, Poulter MO. 1995. Depolarizing GABA-activated Cl^- channels in embryonic rat spinal and olfactory bulb cells. *J Physiol* 488(Pt 2):371–386.
- Sergeeva OA. 1998. Comparison of glycine- and GABA-evoked currents in two types of neurons isolated from the rat striatum. *Neurosci Lett* 243:9–12.
- Sergeeva OA, Haas HL. 2001. Expression and function of glycine receptors in striatal cholinergic interneurons from rat and mouse. *Neuroscience* 104:1043–1055.
- Seri B, Garcia-Verdugo JM, McEwen BS, Alvarez-Buylla A. 2001. Astrocytes give rise to new neurons in the adult mammalian hippocampus. *J Neurosci* 21:7153–7160.
- Sharma G, Vijayaraghavan S. 2001. Nicotinic cholinergic signaling in hippocampal astrocytes involves calcium-induced calcium release from intracellular stores. *Proc Natl Acad Sci USA* 98:4148–4153.
- Smith T, Groom A, Zhu B, Turski L. 2000. Autoimmune encephalomyelitis ameliorated by AMPA antagonists. *Nat Med* 6:62–66.
- Sobko A, Peretz A, Shirihai O, Etkin S, Cherepanova V, Dagan D, Attali B. 1998. Heteromultimeric delayed-rectifier K^+ channels in Schwann cells: developmental expression and role in cell proliferation. *J Neurosci* 18:10398–10408.
- Song H, Stevens CF, Gage FH. 2002a. Astroglia induce neurogenesis from adult neural stem cells. *Nature* 417:39–44.
- Song HJ, Stevens CF, Gage FH. 2002b. Neural stem cells from adult hippocampus develop essential properties of functional CNS neurons. *Nat Neurosci* 5:438–445.
- Sontheimer H, Kettenmann H, Backus KH, Schachner M. 1988. Glutamate opens Na^+/K^+ channels in cultured astrocytes. *Glia* 1:328–336.
- Sontheimer H, Trotter J, Schachner M, Kettenmann H. 1989. Channel expression correlates with differentiation stage during the development of oligodendrocytes from their precursor cells in culture. *Neuron* 2:1135–1145.
- Steinhauser C, Gallo V. 1996. News on glutamate receptors in glial cells. *Trends Neurosci* 19:339–345.
- Steinhauser C, Jabs R, Kettenmann H. 1994. Properties of GABA and glutamate responses in identified glial cells of the mouse hippocampal slice. *Hippocampus* 4:19–35.
- Stewart RR, Hoge GJ, Zigova T, Luskin MB. 2002. Neural progenitor cells of the neonatal rat anterior subventricular zone express functional GABA_A receptors. *J Neurobiol* 50:305–322.
- Tanaka K, Watase K, Manabe T, Yamada K, Watanabe M, Takahashi K, Iwama H, Nishikawa T, Ichihara N, Kikuchi T, Okuyama S, Kawashima N, Hori S, Takimoto M, Wada K. 1997. Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 276:1699–1702.
- Tekkuk SB, Goldberg MP. 2001. AMPA/kainate receptor activation mediates hypoxic oligodendrocyte death and axonal injury in cerebral white matter. *J Neurosci* 21:4237–4248.
- Trimmer JS, Rhodes KJ. 2004. Localization of voltage-gated ion channels in mammalian brain. *Annu Rev Physiol* 66:477–519.
- Ullian EM, Sapperstein SK, Christopherson KS, Barres BA. 2001. Control of synapse number by glia. *Science* 291:657–661.
- Vautier F, Belachew S, Chittajallu R, Gallo V. 2004. *Shaker*-type potassium channel subunits Kv1.3–1.4 and Kv1.6 differentially control oligodendrocyte progenitor cell cycle progression. *Glia* (in press).
- Verdoorn TA, Burnashev N, Monyer H, Seeburg PH, Sakmann B. 1991. Structural determinants of ion flow through recombinant glutamate receptor channels. *Science* 252:1715–1718.
- Verkhratsky A, Kettenmann H. 1996. Calcium signalling in glial cells. *Trends Neurosci* 19:346–352.
- Verkhratsky A, Steinhauser C. 2000. Ion channels in glial cells. *Brain Res Brain Res Rev* 32:380–412.
- Vernadakis A. 1996. Glia-neuron intercommunications and synaptic plasticity. *Prog Neurobiol* 49:185–214.
- Wang DD, Krueger DD, Bordey A. 2003. GABA depolarizes neuronal progenitors of the postnatal subventricular zone via GABA_A receptor activation. *J Physiol* 550:785–800.
- Watanabe M. 2002. Glial processes are glued to synapses via Ca^{2+} -permeable glutamate receptors. *Trends Neurosci* 25:5–6.
- Wonderlin WF, Strobl JS. 1996. Potassium channels, proliferation and G1 progression. *J Membr Biol* 154:91–107.
- Yoshioka A, Hardy M, Younkin DP, Grinspan JB, Stern JL, Pleasure D. 1995. Alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA) receptors mediate excitotoxicity in the oligodendroglial lineage. *J Neurochem* 64:2442–2448.
- Young TL, Cepko CL. 2004. A role for ligand-gated ion channels in rod photoreceptor development. *Neuron* 41:867–879.
- Yuan X, Eisen AM, McBain CJ, Gallo V. 1998. A role for glutamate and its receptors in the regulation of oligodendrocyte development in cerebellar tissue slices. *Development* 125:2901–2914.
- Yuan X, Chittajallu R, Belachew S, Anderson S, McBain CJ, Gallo V. 2002. Expression of the green fluorescent protein in the oligodendrocyte lineage: a transgenic mouse for developmental and physiological studies. *J Neurosci Res* 70:529–545.
- Zhuo L, Sun B, Zhang CL, Fine A, Chiu SY, Messing A. 1997. Live astrocytes visualized by green fluorescent protein in transgenic mice. *Dev Biol* 187:36–42.