



Survey

Regulation and dysregulation of tumor necrosis factor receptor-1

Leen Puimège^{a,b}, Claude Libert^{a,b}, Filip Van Hauwermeiren^{a,b,*}^a Inflammation Research Center, VIB, Ghent, Belgium^b Department of Biomedical Molecular Biology, Ghent University, Ghent, Belgium

ARTICLE INFO

Article history:
Available online 24 March 2014

Keywords:
TNFR1
Gene regulation
Inflammation

ABSTRACT

TNF is an essential regulator of the immune system. Dysregulation of TNF plays a role in the pathology of many auto-immune diseases. TNF-blocking agents have proven successful in the treatment of such diseases. Development of novel, safer or more effective drugs requires a deeper understanding of the regulation of the pro-inflammatory activities of TNF and its receptors. The ubiquitously expressed TNFR1 is responsible for most TNF effects, while TNFR2 has a limited expression pattern and performs immune-regulatory functions. Despite extensive knowledge of TNFR1 signaling, the regulation of TNFR1 expression, its modifications, localization and processing are less clear and the data are scattered. Here we review the current knowledge of TNFR1 regulation and discuss the impact this has on the host.

© 2014 Elsevier Ltd. All rights reserved.

Contents

1. TNFR1, the major receptor of TNF	286
2. The biological significance of TNFR1	287
3. Transcriptional and posttranscriptional regulation of TNFR1	289
3.1. Transcriptional regulation of TNFR1 expression	289
3.1.1. Promoter analysis	289
3.1.2. Factors affecting TNFR1 expression	289
3.1.3. Transcription factor prediction	290
3.2. Post-transcriptional regulation of TNFR1 expression	290
3.2.1. RNA binding proteins	290
3.2.2. microRNAs	290
3.3. Post-translational regulation of TNFR1 expression	290
3.3.1. Storage	290
3.3.2. Shedding	291
3.3.3. Exocytosis	292
3.3.4. Ligand-binding and lipid rafts	292
3.3.5. Endocytosis	292
4. Dysregulation of TNFR1 expression	292
4.1. TNFR1 genetic variants	292
4.1.1. Missense mutations	292
4.1.2. Alternative splicing	293
4.2. TNFR1 in human diseases	293
4.3. Viral interference in TNFR1 expression and function	295
5. Conclusion	296
Acknowledgements	296
References	297

* Corresponding author at: VIB – Ghent University, FSVM Building, Technologiepark 927, B-9052 Zwijnaarde, Belgium. Tel.: +32 93313700; fax: +32 93313609.
E-mail addresses: filip.vanhauwermeiren@VIB-UGent.be, fvhauwer@gmail.com (F. Van Hauwermeiren).

1. TNFR1, the major receptor of TNF

TNF is a type II transmembrane glycoprotein consisting of three monomers with a typical β -jellyroll structure. Each subunit consists of two packed β -sheets of five antiparallel β -strands with three additional β -strands in the N-terminal part (Fig. 1). The first 76 amino acids form a highly conserved hydrophobic sequence that anchors the precursor polypeptide in the membrane. This immature protein (transmembrane pro-TNF) has a molecular mass of 26 kDa and is proteolytically cleaved, mainly by the metalloprotease TNF α converting enzyme (TACE or ADAM17), to a 17-kDa active unit [1]. Also other proteases, such as ADAM10 [2], MMP7 [3] and MMP13 [4], have been shown to cut pro-TNF and generate soluble TNF. Soluble TNF is a homotrimer with a molecular mass of 52 kDa. The TNF protein structure and its interaction with TNFR1 have been described in great detail from high resolution crystals. The homotrimer looks like a triangular cone or bell in which the three subunits are arranged edge to face [5]. The receptor binding sites of TNF are located in the lower half of the triangular cone, in the groove between two subunits [6].

TNF binds with high affinity to two type I transmembrane receptors: TNFR1 (CD120a), which is activated by both soluble TNF (sTNF) and transmembrane TNF (tmTNF), and TNFR2 (CD120b), which is activated mainly by tmTNF. Most of the biological activities of TNF are initiated by binding to TNFR1 [7].

Mouse **TNFR1** has a length of 454 amino acids (AA), of which the 21 N-terminal AAs are a signaling peptide, followed by an extracellular domain (ECD) of 191 AA, a helical transmembrane domain (TMD) of 23 AA, and an intracellular domain (ICD) of 219 AA (Fig. 2). The extracellular regions of TNFR1 and TNFR2 are structurally highly homologous. The N-terminal ECD contains two extracellular topological domains (AA 22–43 and AA 197–212) and four cysteine rich domains (CRD) at AA 44–82, AA 83–125, AA 126–166 and AA 167–196, each of which contains six cysteines. Transmembrane TNFR1 is also a substrate of TACE. The major TNFR1 cleavage site is the spacer region close to the

transmembrane domain between N202 and V203 [8]. There is no significant homology in the intracellular region between TNFR1 and TNFR2, indicating that these receptors activate distinct signaling pathways. TNFR1 contains a cytoplasmic death domain (DD) which is a homophilic protein–protein interaction region of 86 AA (356–441) required for TNF-induced apoptosis, and an N-SMASE activation domain (NSD) spanning an 11-AA motif N-terminal to the DD.

The pre-ligand assembly domain (PLAD) is a homophilic protein–protein interaction motif located in CRD1 and plays an important role in both TNFR1 and TNFR2 signaling pathways by assisting in the assembly of the receptor complex required for TNF binding. It has been proposed that PLAD-mediated homomultimer formation stabilizes CRD2 in a conformation necessary for high affinity ligand binding [9]. In autoimmune diseases, this PLAD region might serve as a target to prevent TNF signaling [10].

Binding of TNF to TNFR1 results in trimerization of the pre-existing receptor complexes and clustering of the intracellular death domains. Subsequently, the adapter molecule TRADD binds by interaction of death domains (DD) of TRADD and TNFR1. TRADD acts as a platform adapter that can recruit TNFR associated factor 2 or 5 (TRAF2/5), cellular inhibitor of apoptosis 1 and 2 (cIAP1/2) and receptor interacting protein 1 (RIP1) to form the membrane-bound **complex I** [11]. This allows cIAP to K63-ubiquitinate RIP1 and TRAF2/5, leading to activation of the inhibitor of κ B (I- κ B) kinase complex (IKK) [12]. Additionally, linear ubiquitination of IKK γ or NEMO by the LUBAC complex stabilizes the IKK complex [13]. Phosphorylation of I- κ B by IKK ensures I- κ B K48-ubiquitination and degradation by the proteasome and consequent activation of NF- κ B. Activation of AP-1 involves a phosphorylation cascade mediated by the mitogen activated protein (MAP) kinases. These kinases are responsible for activating c-Jun N-terminal kinases JNK1, 2 and 3 and p38, leading to activation and nuclear translocation of c-Fos and c-Jun [11]. Hence, complex I stimulates pathways leading to activation of NF- κ B and AP-1 and induction of pro-inflammatory and anti-apoptotic genes. One such anti-apoptotic gene encodes c-FLICE inhibitory protein (c-FLIP), which




Fig. 1. Crystal structure of TNF (PDB 1TNF in green) and binding to its receptor TNFR1 (PDB 1EXT in blue). Top view is on the left and side view on the right [171,172].




Fig. 2. Representation of TNFR1 (UniProtKB P25118). For detailed description see text.

is a specific inhibitor of caspase 8 (an essential molecule in the apoptotic pathway) [11]. However, prolonged JNK activation can also induce pro-apoptotic proteins such as Itch, which can mediate c-FLIP degradation [14]. However, the duration of NF- κ B activation is limited by several feedback mechanisms, such as the induction of I- κ B, CYLD and A20. CYLD, a protease that specifically cleaves K63-ubiquitin chains, de-ubiquitinates TRAF2, thereby inhibiting the recruitment of TAB/TAK and activation of IKK [15], and A20 deactivates RIP1 by removing the K63-ubiquitin chain and adding a K48-ubiquitin chain [16]. Sustained MAPK activation is prevented by MAPK phosphatases such as MKP-1. However, TNF-induced reactive oxygen species (ROS) can disturb some of the feedback systems, for example by oxidizing MKP-1 [17].

Upon endocytosis of complex I, TRADD dissociates from TNFR1 and associates with Fas-associated protein with death domain (FADD) to form the intracellular located **complex II**. De-ubiquitination of RIP1 leads to recruitment and autocatalytic cleavage of pro-caspase 8, thereby initiating apoptosis [18]. However, in conditions where caspase 8 is inhibited, RIP3 can be recruited and RIP1 and RIP3 become phosphorylated by auto phosphorylation or cross phosphorylation, leading to necroptosis, a regulated form of necrosis. Necroptosis causes rapid plasma membrane permeabilization with release of reactive oxygen species (ROS) and exposure of damage-associated molecular patterns (DAMPs), which provide strong stimulation of the immune system [19].

2. The biological significance of TNFR1

TNF signals through two distinct receptors, TNFR1 and TNFR2. These receptors initiate diverse important effects, including

proliferation, differentiation, migration, inflammation and cell death [20]. The pro-inflammatory and pathogen-clearing activities of TNF are mediated mainly through activation of TNFR1, which is a strong activator of NF- κ B, while TNFR2 may be more responsible for suppression of inflammation [21].

Several groups generated **TNFR1 deficient mice** by gene targeting [22–24]. Studies on these mice have contributed a lot to understanding the role of this receptor in the biological activities of TNF. It is clear that TNF has a key role in immunity and immunomodulation as well as pro-inflammatory and antitumor activities. Resistance against bacterial infection, e.g. *Listeria monocytogenes*, is mediated by TNFR1: mice deficient in TNFR1 are extremely sensitive to *L. monocytogenes* and other Gram-positive bacteria, including *Yersinia enterocolitica* [25,26], *Pseudomonas aeruginosa* [27], *Legionella pneumophila* [28] and *Burkholderia pseudomallei* [29], and also to other pathogens, such as viruses, e.g. influenza virus [30] and vaccinia virus [31]. However, injection of TNF in mice leads to systemic inflammation, which, depending on the dose, can be lethal. Studies on TNFR1 knockout (KO) mice have shown that lethality, which involves inflammation [32] and necroptosis [19], is mediated entirely by TNFR1. Chronically overexpressed TNF leads to arthritis and inflammatory bowel diseases, both of which depend on TNFR1 [33]. TNF-induced antitumor effects are also mediated entirely by TNFR1 on host-derived neo-vascular endothelial cells, which grow into the tumor and are important for sustaining tumor growth and survival [34].

Anti-TNF therapy is effective for treatment of autoimmune diseases such as rheumatoid arthritis and inflammatory bowel diseases, indicating that the TNF signaling pathway is an appropriate target [35,36]. Nevertheless, long-term TNF blockade




Fig. 3. TNFR1 pathway. Binding of TNF to TNFR1 triggers recruitment of TRADD, TRAF2, CIAP1/2 and RIP1, forming complex I. Activation of the IKK complex, leads to proteolytic degradation of I-κB proteins and subsequent activation of NF-κB which is then translocated to the nucleus to activate transcription of target genes. Upon endocytosis of complex I, TRADD dissociates from TNFR1 and associates with FADD and caspase 8 to form intracellular complex II. Caspase 8 becomes activated and initiates apoptosis. For detailed description, see text.

can cause substantial side-effects, such as opportunistic infections [37] and development of additional autoimmune diseases, including lupus, type 1 diabetes, uveitis, multiple sclerosis and psoriasis, as well as lymphoma and leukemia [38,39].

Interestingly, TNFR1 +/- mice, which express 50% TNFR1 on cells, were also found to be completely resistant to TNF-induced lethal inflammation over a huge dose range [40]. These data, together with others, such as those showing that mice expressing higher levels of cell-bound TNFR1 develop spontaneous inflammation [41], suggest that TNFR1 is a potential drug target, and that minor TNFR1 regulation might have substantial physiological and pathological effects. Furthermore, in humans, mutations affecting TNFR1 have been linked with the development of TNFR1-associated periodic syndromes (TRAPS), which are characterized by recurrent fever attacks and localized inflammation [42]. Hence, selective inhibition of TNFR1 signaling might be a better alternative for the treatment of TNF/TNFR1-mediated autoimmune diseases because it would reduce the pro-inflammatory actions of TNFR1 without impairing the immunosuppressive properties of TNFR2 [43,44]. A variety of compounds have already been generated, including monoclonal antibodies and derivatives, antagonistic TNF variants (R1AntTNF), RNAi and antisense oligo's (ASOs), which are all able to inhibit TNF-mediated NF-κB gene expression and apoptosis. Many of these inhibitors have already proven to block TNF-induced lethality, allowing safe anti-tumor

therapy [40,45], while others have shown to be protective in mouse models of several TNF-induced chronic disease, such as CIA, EAE, RA, VILI, CCL4-, conA- and TNF/GalN-hepatitis [46–53].

TNF receptors form homotrimers upon activation by TNF but no heterotrimers are assembled and the TNFR1/TNFR2 protein ratio has been found to be important for the TNF response. TNF affects NF-κB activation predominantly through TNFR1, whereas TNFR2 activates transcription poorly [54]. Nevertheless, TNFR2 stimulation can result in competition for TRAF2 and cIAPs and thereby inhibit TNFR1-induced NF-κB transcription. Reduction of NF-κB activation promotes apoptosis in certain cell types due to diminished production of anti-apoptotic factors [55]. Hence, constant TNFR1 expression coupled to changeable TNFR2 levels alters the TNFR1/TNFR2 ratio and controls the response of the cell to TNF stimulation [56]. On the other hand, the kinetics of TNF binding to TNFR2 suggests a mechanism by which TNFR2 might increase the apparent rate of TNF binding to TNFR1. TNFR2 has a higher affinity and longer TNF-binding half-life than TNFR1. By a so-called ligand-passing mechanism TNFR2 is thought to associate with TNF, increasing its concentration near TNFR1 receptors, and making TNF available for activating TNFR1 [57]. Clearly, the outcome of TNFR1 and TNFR2 signaling is complex and may also depend on the cell type and the activation status of the cell.

As an important step to completely understand the biology of TNF and its receptors and to develop specific TNFR1 inhibiting

Fig. 4. Promoter of mouse *Tnfrsf1a* (NM011609). The 5'UTR is shown in red and the CDS in green. Transcriptional regulatory sequences are marked in green and the GC rich elements in yellow.

tools, a comprehensive overview of the available knowledge about regulation of TNFR1 on all levels is provided in this review.

3. Transcriptional and posttranscriptional regulation of TNFR1

3.1. Transcriptional regulation of TNFR1 expression

Despite the critical role that TNFR1 plays in TNF-mediated signaling, little is known about the regulation of its promoter. It has been suggested that the TNFR1 promoter is constitutively active, like the promoters of “housekeeping” genes, at low levels and in all nucleated cell types [56,58]. TNFR2, on the other hand, is inducible and expressed exclusively by immune cells, endothelial cells and some neuronal populations [59].

3.1.1. Promoter analysis

The 5' flanking region of the TNFR1-coding gene (*TNFRSF1A* in humans and *Tnfrsf1a* in mice) was scanned for the presence of sequence motifs that have been associated with regulation of gene transcription [60]. According to the UCSC genome browser, the transcription start site (TSS) is located 107 bp upstream of the putative TSS that was described by Takao and Jacob. A putative **TATA box** (TTAAATT), the core promoter sequence, is now located between +63 and +69 downstream of the TSS, which presumes that it may not be a true TATA box. Two **GC-rich elements** are present between -44 and +3 and between +5 and +42. These elements have been shown to possess enhancer activity in many eukaryotic genes [60].

The TNFR1 promoter contains a functionally important **binding site** for CCAAT/enhancer binding protein (**C/EBP**), which contributes to the constitutive activity of the promoter. C/EBP transcription factors play essential roles in regulating different cellular processes, including differentiation, energy metabolism, and inflammation. Both C/EBP α and C/EBP β bind to a sequence located between +5 and +12 [60,61]. There are two copies of the consensus **binding site for AP-1**, a transcription factor that regulates gene expression in response to different stimuli, including cytokines, growth factors, stress, and bacterial and viral infections [60]. Also four potential binding sites for the **AP-2** family of transcription factors (AP-2 α , AP-2 β , AP-2 γ , AP-2 δ and AP-2 ϵ) have been found. The general functions of this family are stimulation of proliferation and suppression of differentiation during embryonic development [62]. An **NF- κ B binding site** in the promoter between -489 and

–498 activates the expression of TNFR1. This element is active and important, since in mammary-specific β -lactoglobulin Cre⁺/*Ilk2*^{fl} mice, NF- κ B DNA-binding activity is diminished by 50%, leading to reduced expression of TNFR1 mRNA levels and reduction of TNF-induced apoptosis [63]. A consensus sequence related to the **IFN γ activated site (GAS)** of signal transducer and activator of transcription (**STAT-1**) factors has been found between –235 and –243 of the mouse TNFR1 promoter. In oligodendrocytes, IFN γ indeed induces TNFR1 transcription via activation and binding of STAT-1 homodimers to the GAS site in the TNFR1 promoter [64,65] (Fig. 4 and Table 1).

3.1.2. Factors affecting TNFR1 expression

Gene transcription is controlled by dynamic acetylation and deacetylation of histone proteins (and other proteins), which alter chromatin structure and affect transcription factor access to the DNA.

HDAC5 (histone deacetylase 5) overexpression has been shown to inhibit tumor cell growth and induce spontaneous apoptosis by altering gene expression, including a fourfold up-regulation of TNFR1 [66]. Transcriptional up-regulation of TNFR1 by **NF- κ B** activation has been observed and been shown to influence the contact between T-cells and monocytes [67]. Moreover, **IL18** and

Table 1

Transcriptional regulatory sequences in the promoter of *Tnfrsf1a*. Lower case indicates deviations from the consensus (Y=C/T, R=A/G, M=A/C, K=G/T, W=A/T, S=G/C).

Transcriptional element	Consensus sequence	Actual sequence	Position
TATA box?	TATAWAW	TtAAAtT	+63/+69
C/EBP binding site	RTTGCGYAA	cTTGCaac	+5/+12
GC rich element			-44/+3
AP-1 binding site	TGASTCA	TGAAtTct	+5/+41
		aGAGTCA	-941/-947
AP-2 binding site	GCCNNNGGC	cCCAACGGa	-467/-473
		cCCCAcCgt	-843/-850
		aCCcAGGca	-527/-535
		cCCCAGGct	-436/-444
		cCCCCGcca	-11/-19
NF- κ B binding site	GGGACTTTCC	GGGACTTgtC	+17/+25
IL-6 responsive element	CTGGGA	CTGGGA	-489/-498
cAMP responsive element	TGACGTCA	TGAgGTCA	-669/-674
IFN γ activated site	TTCNNNGAA	TTCCGTGAA	-853/-860
			-235/-243

Table 3
miRNA target prediction for Tnfrsf1a. 1/green indicates predicted, 0/red indicates not predicted.

	microCosm	miRanda	miRGen	NBmiRTar	TargetScan	miRDB	DIANA-microT	microInspector	miRWalk	PICTAR	PITA	RNA22	RNAhybrid	sum
mmu-mir-29a/b1	1	1	1	0	1	0	1	0	1	1	1	0	0	8
mmu-mir-29b2/c	1	1	1	0	1	0	1	0	1	1	1	0	0	8
mmu-mir-125a	1	0	1	1	1	0	1	1	1	0	1	0	0	8
mmu-mir-680-1	1	1	1	0	1	0	1	1	1	0	1	0	0	8
mmu-mir-351	1	0	1	0	1	0	1	1	1	0	1	0	0	7
mmu-mir-763	1	1	1	0	1	0	1	0	1	0	1	0	0	7
mmu-mir-125b	1	0	1	0	1	0	1	0	1	0	1	0	0	6
mmu-mir-149	0	0	1	0	1	0	1	1	1	0	1	0	0	6
mmu-mir-511	1	1	0	0	1	0	0	1	1	0	1	0	0	6

proteins may play an important role in controlling the cell's response to TNF [81]. The Golgi pool of TNFR1 is believed to serve to replenish cell surface TNFR1 receptors (Fig. 5).

3.3.2. Shedding

TNFR1 can be released from the cell surface by a proteolytic process called ectodomain shedding (Fig. 5). Like TNF, TNFR1 and TNFR2 are cleaved by TNF- α converting enzyme (TACE) [1]. Proteolytic cleavage of TNFR1 decreases the surface expression. Moreover, the generation of antagonistic soluble receptors by shedding can regulate TNF bioactivity by preventing its binding to

membrane receptors [82]. TACE may therefore exert either pro- or anti-inflammatory effects, depending on whether it acts on an effector cell (e.g. releasing TNF from macrophages) or target cell (e.g. releasing TNFR1 from endothelial cell) [83]. However, low levels of sTNFR may stabilize the activity of TNF and provide a reservoir of TNF [84]. In humans, mutations affecting the shedding of TNFR1 have been linked with the development of TNFR1-associated periodic syndromes (TRAPS), which are characterized by recurrent fever attacks and localized inflammation. Similarly, knock-in mice expressing a mutant non-sheddable TNFR1 are very sensitive to TNF-induced inflammation and develop several




Fig. 5. Compartmentalization of TNFR1 signaling. Black arrows indicate transport of TNFR1, 1 = translation and storage, 2 = membrane expression, 3 = shedding, 4 = exocytosis, 5 = endocytosis. Yellow arrows indicate signaling pathways. For detailed description, see text.

auto-immune-like diseases, such as spontaneous hepatitis, arthritis, and autoimmune encephalomyelitis [41].

Basal expression of TACE is observed in all vasculature cell types, including endothelial cells, vascular smooth muscle cells, fibroblasts, and leukocytes. TACE-mediated shedding is enhanced by stimulation with pro-inflammatory cytokines (TNF, IFN γ), TLR ligands (LPS), growth factors (PDGF, VEGF), GPCR ligands (thrombin) or oxidative stress (ROS), and thus involved in many regulatory pathways. Transcriptional regulation, maturation by furin, trafficking from storage pools, intracellular phosphorylation, changes in cellular distribution within membrane lipid rafts, interaction with adapter molecules (tetraspanins) and conformational changes of the protease all lead to enhanced substrate cleavage by TACE [85]. TACE is expressed in intracellular compartments and on the cell surface. Removal of its pro-domain by a furin protease probably occurs within the Golgi apparatus [86]. Distribution of TACE and its substrates within lipid rafts affects shedding because mature TACE is associated with cholesterol-rich lipid rafts, and depletion of rafts induces shedding of TNF, TNFR1 and TNFR2 [87,88]. MAP kinases, oxidative stress and other TACE regulating pathways have been studied in great detail [89–93].

The mechanism of TNFR1 shedding might also involve interactions with regulatory ectoproteins. A direct relationship exists between the level of the type II integral membrane protein **ARTS-1** (aminopeptidase regulator of TNFR1 shedding) and the degree of TNFR1 shedding. ARTS-1 overexpression increases TNFR1 shedding and decreases membrane-associated TNFR1, while expression of antisense ARTS-1 mRNA decreases membrane-associated ARTS-1 and TNFR1 shedding but increases membrane-associated TNFR1. ARTS-1 neither bound to TNFR2 nor altered its shedding, indicating its specificity for TNFR1. Findings suggest that ARTS-1 does not possess TNFR1 sheddase activity, which indicates that it is a multifunctional ectoprotein capable of binding to and promoting TNFR1 shedding [94].

3.3.3. Exocytosis

TNFR1 shedding is a consequence of proteolytic cleavage of the 28-kDa ectodomain by TACE. But in human vascular epithelial cells (HUVECs) and bronchoalveolar lavage (BAL) fluid, the predominant form of soluble TNFR1 is the full-length 55-kDa protein (eTNFR1) associated with **exosome-like vesicles** (Fig. 5). Exosomes are small membrane-enclosed vesicles of 30–100 nm released from the cell by exocytosis. These intracellular TNFR1-containing vesicles enable constitutive release of full length TNFR1 into the extracellular compartment. This is an alternative pathway for generating soluble cytokine receptors independently of proteolytic cleavage of the receptor ectodomain [95]. It has been reported that histamine induces redistribution of TNFR1 from the Golgi to vesicles and subsequently into the medium as soluble receptors, with a consequent decrease in cell surface TNFR1 [82]. The precise significance of such remarkable full-length TNFR1 shedding is still unclear.

3.3.4. Ligand-binding and lipid rafts

Upon binding of TNF, TNFR1 translocates to cholesterol- and sphingolipid-enriched membrane microdomains called **lipid rafts**, where the associations with RIP1 and the adaptor proteins TRADD and TRAF2 occur and form the signaling complex I (Fig. 3). This complex triggers pathways leading to induction of pro-inflammatory and anti-apoptotic proteins. Activation of p42, a member of the MAP kinase family, leads to TNFR1 phosphorylation at a consensus MAPK site in the cytoplasmic domain, which alters its subcellular localization, resulting in changes in its signaling properties [96]. Moreover, in lipid rafts TNFR1 and RIP1 become ubiquitinated, which leads to their degradation via the proteasome

pathway [97]. Disruption of lipid rafts, by depleting cholesterol, not only abolishes ubiquitination but also totally blocks TNF-induced NF- κ B activation, leading to a switch to apoptosis induction which indicates that the translocation of NF- κ B to lipid rafts is essential for its activation [98]. So, the subcellular localization of TNFR1 determines its available amount and signaling outcome.

3.3.5. Endocytosis

Formation of complex I is transient because most of TRADD, RIP1 and TRAF2 dissociate from TNFR1 within an hour, when TNFR1 starts to undergo endocytosis (Fig. 5). Treatment of cells with TNF led to maximal down-modulation of the TNF receptors within 30 min. Ligand-induced down-modulation of TNF receptors is caused by TNFR1 internalization [99]. The liberated DD of TRADD now binds to FADD, resulting in caspase 8 recruitment and complex II formation. Complex II initiates apoptosis, provided that NF- κ B signaling has terminated [98]. Endocytosis or TNFR1 internalization and intracellular trafficking plays an important role in selection of the signaling pathway: either internalization-independent (pro-inflammatory complex I) or internalization-dependent (pro-apoptotic complex II). Apoptosis is totally blocked by preventing internalization with monodansylcadaverine (MDC), an inhibitor of transglutaminase, a membrane-bound enzyme that actively participates in internalization of various receptor systems [100]. The endosome should therefore be recognized as a signaling organelle involved in selectively transmitting death signals from TNFR1 [101]. Internalization proceeds through the classical receptor-mediated endocytosis pathway, i.e. via clathrin-coated vesicles and endosomes that fuse and enter multivesicular bodies (MVB) before accumulating in lysosomes [102]. After apoptosis, TNFR1 protein can be eliminated in a caspase-dependent manner. *In vitro*, the cytoplasmic tail of hTNFR1 is susceptible to cleavage by the downstream executioner caspase 7, the only caspase capable of cleaving TNFR1. Identification of the cleavage site revealed an EXE motif instead of the classic EXD motif. Homologous sequence alignments showed that the EXE motif is conserved in rat and pig but not in mouse and cow, which have 3–4 missing amino acids at this site [103]. TNF thus causes the down-modulation of its own receptor by internalization which might function as a negative feedback mechanism as it will desensitize cells for further TNF stimulation until the membrane TNFR levels have recovered.

4. Dysregulation of TNFR1 expression

4.1. TNFR1 genetic variants

4.1.1. Missense mutations

TNF receptor-associated periodic syndrome (TRAPS) is caused by missense mutations in the TNFRSF1A gene and characterized by periodic high fevers, rashes, abdominal pain, chest pain, conjunctivitis, arthralgia and myalgia. More than 100 mutations have been identified in TNFR1, all of them in the extracellular domain. (<http://fmf.igh.cnrs.fr/ISSAID/infevers>) [104]. TRAPS mutations occur predominantly in CRD1 and CRD2 and many of them involve intramolecular disulfide bonds. Others occur at residues predicted to have an effect on the secondary structure or at residues involved in hydrogen bonds between loops of the receptor. The absence of large deletions suggests that synthesis of the mutant protein is important for pathogenesis. It has been hypothesized that TRAPS pathology is driven by defective receptor shedding resulting in reduced serum levels of soluble TNFR1. This hypothesis was based on observations that cells of some TRAPS patient are resistant to PMA-induced shedding and that serum from TRAPS patients contained reduced levels of

circulating TNFR1. However, the shedding defect was not always present and varied depending on the studied cell types [105]. Molecular modeling indicates that nine TNFR1 mutants are unable to bind TNF (H22Y, C30S, C30R, C33G, C34S, T50M, C52F, C88R, R92P), whereas the R92Q and P46L mutants can bind TNF. Due to misfolding, they accumulate in the endoplasmic reticulum (ER) instead of localizing to the cell surface, and are most likely degraded by the proteasome [106]. But accumulation of mutant TNFR1 in the ER may also trigger the ER stress response, which can directly or indirectly lead to inflammation or to reduced TNF-induced apoptosis. Interestingly, neutrophils and dermal fibroblasts from TRAPS patients with several different mutations have reduced apoptosis but produce the proinflammatory cytokines IL-6 and IL-8 normally when exposed to TNF [107,108]. Failure of activated cells to undergo apoptosis in TRAPS could lead to accumulation of pro-inflammatory cytokines. However, TRAPS mutations might not all act by the same mechanism [109,110]. Glucocorticoids are effective in decreasing the severity and duration of the fever attacks, although their efficacy fades with time [111]. The soluble TNF receptor Etanercept, the IL-1 β receptor antagonist Anakinra and the IL-6 receptor antagonist Tocilizumab are effective in some patients. However, since no controlled clinical trials have been performed for this rare syndrome, there is currently no single recommended treatment [112].

Two rare single amino acid mutations in *TNFRSF1A* have been identified as low-penetrance risk factors for TRAPS. The R92Q substitution is carried by ~2% of North American and Irish populations, and the P46L mutation is present in 9% of African populations. TRAPS patients with these polymorphisms have a milder syndrome with almost no incidence of amyloidosis [42]. But the R92Q mutation has also been linked with other diseases associated with inflammation, such as **rheumatoid arthritis** and **atherosclerosis** [113,114]. Clinical observations have identified some **multiple sclerosis** (MS) patients carrying the R92Q mutation and exhibiting additional TRAPS symptoms. The co-existence of MS and TRAPS could be mediated by this mutation. The R92Q mutation acts like a genetic risk factor for MS and other inflammatory diseases, including TRAPS. Nevertheless, this mutation does not appear to be a severity marker, modifying neither the progression of MS nor its response to therapy. But an alteration in TNF/TNFR1 signaling may increase pro-inflammatory signals [115].

Single nucleotide polymorphisms (SNPs) in the *TNFRSF1A* gene may also influence the innate immune response against **invasive pulmonary aspergillosis** (IPA), an increasingly common opportunistic fungal infection that usually occurs in immune compromised patients [116]. In this infection, TNF acting through TNFR1 plays a pivotal role in immune regulation and host immune responses. Three SNPs were genotyped in 275 individuals (52% immunocompromised hematological patients with high-risk of developing IPA and 48% healthy controls): A383C and G609T in the 5' UTR, and A36G in exon 1. The last two are associated with IPA susceptibility. The role of *TNFRSF1A* SNPs is also supported by significantly lower TNFR1 mRNA levels in IPA compared to IPA-resistant patients and by a strong correlation between the -609 SNP and TNFR1 expression levels. TNFR1 polymorphisms may influence the risk of IPA disease and might be useful for risk analysis [117]. The A36G mutation is also significantly associated with lower hemoglobin levels, causing iron deficiency anemia in patients with early or established rheumatoid arthritis. The frequency of anemia is higher in GG homozygous patients [118,119].

The pathologies induced by mutations in TNFR1 further emphasize the important role of TNFR1 in autoimmune diseases and immunity. Furthermore, they indicate that TNFR1 targeting might be a better alternative to current TNF inhibition therapies

since Etanercept and Infliximab fail in certain TRAPS family with T50M, C30S and R92Q mutations [120,121].

4.1.2. Alternative splicing

Two disease-associated isoforms of TNFR1 mRNA produced by alternative splicing were recently described.

The **TNFR1- Δ 6** splicing pattern is caused by a variation (rs1800693) in the exon 6/intron 6 border region that alters exon 6 splicing. The frame shift caused by skipping of exon 6 results in a protein lacking the intracellular and transmembrane domains and part of the extracellular domain. This TNFR1 splice form is associated with multiple sclerosis (MS) but not with other autoimmune conditions such as rheumatoid arthritis, psoriasis or Crohn's disease. The MS risk allele induces the expression of a soluble TNFR1 form that can block TNF. Importantly, TNF-blocking drugs can promote onset or exacerbation of MS, but they are very effective for autoimmune diseases that are not associated with rs1800693. This indicates that the clinical experience with these drugs corroborates the disease association of rs1800693, and that the MS-associated TNFR1 variant mimics the effect of TNF-blocking drugs [122].

The **TNFR1- Δ 2** splice variant is regulated by three variations affecting the phenotype of TRAPS. They occur in the promoter, exon 1 and intron 4 (rs4149570, rs767455 and rs1800692, respectively) of the *TNFRSF1A* gene. Exon 2 alternative splicing increased with the T-A-T haplotype at rs4149570–rs767455–rs1800692 as compared with the G-G-C haplotype and transcriptional activity increased with the T-T haplotype compared with the G-C haplotype, suggesting that regulation of TNFR1- Δ 2 expression may occur via a coupling mechanism between transcription and splicing. Whereas TNFR1 is ubiquitously expressed, TNFR1- Δ 2 is expressed tissue-specifically in human PBLs, brain, heart, kidney, skeletal muscle, small intestine and spinal cord, but not in liver or lung [123].

4.2. TNFR1 in human diseases

Because TNF is a key immune system modulator and has broad effects, excessive signaling can cause significant damage. A delicate balance exists between beneficial immune stimulation and pathogenesis. Several autoimmune and inflammatory diseases have been associated with the effects of dysregulated TNF activation [7]. It is not always evident whether abnormal levels of TNF, TNFR1 or signaling components are the cause or consequence of a given pathology. Therefore, understanding the regulation of TNF and its receptors is essential for elucidating how TNF can either prevent or induce various diseases. Here (Table 4), we provide an overview of misregulations in TNF/TNFR levels as described in several human diseases.

Patients with **systemic lupus erythematosus** (SLE), a multi-organ inflammatory autoimmune disease, have altered expression of TNF-related signaling molecules, suggesting that imbalance of TNF signaling favors cellular activation rather than apoptosis. SLE patients have increased levels of TNFR1, TNFR2 and TRAF2 and decreased levels of RIP1 on various naive and memory B-cell and T-cell subsets as compared to controls. However, the levels of these molecules are not correlated with their RNA expression or with serum TNF levels in peripheral whole blood [124] (Table 4).

Up-regulation of the intercellular adhesion molecule (ICAM)-1 on human conjunctival epithelial cells (HCECs) is an important feature of **ocular allergic inflammation**. TNF produced by human conjunctival mast cells up-regulates the expression of TNFR1 on HCECs, resulting in a stronger TNF-mediated response, including up-regulation of ICAM-1. So, up-regulation of TNFR1 expression results in enhanced ICAM-1 expression in response to TNF

Table 4

The regulation of TNFR1 in several TNFR1-mediated diseases. For detailed description, see text.

Disease	Biomarker	Tissue	Effect	Ref.
Systemic lupus erythematosus	TNFR1 protein up	B-cells, T-cells	Cellular activation	[124]
Ocular allergic inflammation	TNFR1 protein up	HCECs	I-CAM expression	[125]
Aging	TNFR1 protein up	Lymphocytes	Apoptosis	[126]
Acute myocardial infarct	TNFR1 protein up	Cardiomyocytes, ECs	Angiogenesis down	[127,128]
Ischemic cerebrovascular diseases	sTNFR1 protein up	Serum	Proliferation	[129,130]
Lung epithelium injury	sTNFR1 protein up	BALF	Neutralize TNF	[131]
Lupus nephritis	TNFR1 protein up	Urine	?	[132]
Behcet's disease	sTNFR1 protein up	Serum, synovial fluid	?	[133–136]
Huntington's Disease	TNFR1 mRNA down	Skeletal muscle	?	[137–139]
Hepatocellular carcinoma	sTNFR1 protein up	Serum, ascetic fluid	?	[140,141]
	TNFR1 mRNA down	Liver		[142]

stimulation, demonstrating that targeting TNFR1 expression may be an effective treatment for ocular inflammation [125].

Aging is characterized by increased susceptibility of T-cells to TNF-induced apoptosis due to increased expression of TNFR1 and TRADD and decreased expression of TNFR2 and TRAF-2. Moreover, there is an increased activation of caspase 8 and caspase 3, confirming that increased TNF-induced apoptosis may play a role in T-cell deficiency associated with human aging [126].

Development of arterial diseases is associated with age-related impairment of angiogenesis. Because TNFR1 mediates the cytotoxic effects of TNF, whereas TNFR2 is mostly involved in the protective effects, TNF signaling via its receptors has diverse effects on neovascularization, repair and regeneration in adult tissue after **acute myocardial infarction** (AMI). Intact signaling through both TNFR1 and TNFR2 assures sufficient NF- κ B activation, followed by transcriptional activation of *VEGFA*, *FGFB* and other pro-angiogenic genes. But age-associated decrease of TNFR2 levels, coupled with post-ischemic increase in systemic TNF, favors apoptosis in adult cardiomyocytes and endothelial cells due to reduced NF- κ B

activation, leading to reduced angiogenesis. Decreased TNFR2 expression in adult tissue also stimulates pro-apoptotic signaling through TNFR1 by the release of vacant TRADD [127]. But signaling via both TNFR1 and TNFR2 is necessary to prevent reperfusion injury after AMI during late preconditioning. So, total blocking of TNF is not advised in AMI patients. Modulation of TNFR1 and/or TNFR2 expression at different stages of AMI may have important implications for prevention of myocardial injury and enhancement of myocardial repair and regeneration [128].

Furthermore, it has been shown that patients with **acute ischemic cerebrovascular diseases** also have higher plasma levels of sTNFR1 [129]. Animal studies have shown that TNFR1 is associated with decreased neuronal proliferation after stroke and that deletion of TNFR1 enhances neuroblast formation and recovery. These results provide evidence that TNFR1 is a negative regulator of stroke-induced progenitor proliferation; the proliferative response after stroke might be promoted by blocking TNFR1 signaling [130].

Ozone (O₃) is an air pollutant that causes **lung epithelium injury** leading to inflammation. The inflammatory response to

Table 5

Viral mechanisms of escape from the host response. For detailed description, see text.

Virus	Viral protein	Interference	Mechanism	Ref.
Epstein–Barr virus	BZLF1	TNFR1 transcription	Reduction of TNFR1 promoter activity	[144]
Hepatitis C virus	?	TNFR1 transcription	Inhibition of TNFR1 mRNA expression	
Herpes simplex virus	UL41	TNFR1 translation	Degradation of TNFR1 mRNA and inhibition of apoptosis	[145]
Cytomegalovirus	Immediate early (IE) viral products	TNFR1 Golgi-trafficking	Down-regulation of TNFR1 surface expression	[146,147]
Poliovirus	3A	TNFR1 Golgi-trafficking	Down-regulation of TNFR1 surface expression	[148]
Respiratory syncytial virus	?	TNFR1 shedding	IL10 production and increased production of soluble TNF receptor	[149,150]
Adenovirus	E3-14.7K	TNFR1 internalization	Inhibition of apoptosis and increased NF- κ B signaling	[151]
Myxoma virus	Secreted M-T2 (pseudo sTNFR)	TNFR1 signaling	Inhibition of TNF-induced cytotoxicity in T-cells	[152,153]
Myxoma virus	Intracellular M-T2	TNFR1 signaling	Inhibition of apoptosis (T-cells)	[153]
Orthopoxviruses	CrmB,C,D,E (pseudo sTNFR)	TNFR1 signaling	Blocking TNF function	[154]
Epstein–Barr virus	LMP-1 (pseudo TNFR)	TNFR1 signaling	Ligand independent NF- κ B activation (infected B-cells)	[155,156]
Human herpes virus 6B	U20	TNFR1 signaling	Inhibition of TNFR1 signaling and apoptosis	[157]
Herpesvirus saimiri	STP (pseudo TNFR)	TNFR1 signaling	Ligand independent NF- κ B activation via association with TRAFs (infected T-cells)	[158]
Human papilloma virus 16	E6	TNFR1 signaling	TNFR1 binding with inhibition of TRADD binding and DISC formation leading to reduced TNF-induced apoptosis	[159]
Respiratory syncytial virus	RSV-G protein (pseudo TNFR1)	TNFR1 signaling	Inhibition of TNF-induced apoptosis by I- κ B proteolysis	[160]
Hepatitis C virus	HCV core protein	TNFR1 signaling	TNFR1 DD binding with inhibition of TRADD binding, facilitating FADD binding and leading to increased TNF-induced apoptosis (HepG2, Hela)	[161,162]
Hepatitis C virus	HCV core protein	?	Increasing NF- κ B nuclear retention and DNA binding and increasing I- κ B degradation leading to reduced apoptosis	[163,164]
Hepatitis C virus	HCV core protein	?	Inhibition of TNF-induced apoptosis (MCF7)	
Tanapox virus	Glycopeptide	?	Inhibition of TNF-induced NF- κ B activation leading to reduced cell adhesion molecules	[167]
Parvovirus	?	?	Activation of caspase 3 and down-regulation of c-myc leading to increased apoptosis	[168,169]

acute ozone exposure includes the production of numerous cytokines and chemokines, resulting in influx of neutrophils. Genetic linkage studies on both mice and humans have shown that a locus encompassing the *TNF* gene plays a role in responses to O₃. In mice, age-related differences in the inflammatory response to acute O₃ exposure vary with TNFR1 expression. In fact, it has been shown that sTNFR1 increases during disease in the BAL fluid, and that this sTNFR1 neutralizes TNF in the lung and so protects against O₃-induced inflammation [131].

Other diseases may also be associated with differences in TNFR1 levels, however the effects are less clear (Table 4). In mice and patients with **lupus nephritis**, an inflammation of the kidney caused by SLE, TNFR1 is found to be enriched in the urine especially at the peak of disease. Whether the increased urinary TNFR1 levels reflect increased shedding and are relevant to disease progression remains unclear [132]. Patients with active **Behcet's disease** (BD), a chronic, multi-organ immunoinflammatory vasculitis that often presents with mucous membrane ulcers, ocular lesions and arthritis, have increased levels of plasma sTNFR1 and sTNFR2, especially when arthritis is present [133]. TNF appears to be important in initiating Behcet's disease, as indicated by the effects of blocking TNF. Infliximab is a promising treatment for uveitis associated with the disease and Etanercept is useful for patients with mainly skin and mucosal symptoms [134–136]. Since the level of TNFR1 correlates well with disease activity index scores of these diseases, the increase may be useful for diagnosis of Behcet's and lupus.

In **Huntington's disease** (HD), a neurodegenerative disorder caused by expansion of a glutamine-encoding CAG repeat which affects muscle coordination and results in cognitive regression and psychiatric problems, the opposite is observed. Mouse models representing the earlier stages of human HD (*Hdh*^{CAG(150)} knock-in mice) or the later stages of the rare, but more severe, juvenile form of human HD (R6/2 transgenic mice) [137,138] and muscle biopsies from HD patients, showed decreased *Tnfrsf1a* gene expression [139]. Studies with TNFR1 deficient mice could be very useful to confirm these observations.

Hepatocellular carcinoma (HCC) is the fifth most common neoplasia in the world and the first cause of death by cancer in some regions. sTNFR1 levels in the serum and ascites fluid of patients with HCC are significantly higher than in controls, and they correlate positively with total bilirubin and alpha fetoprotein in the peripheral blood. This reflects an abnormal immune status of HCC patients and can help to predict the progression of the tumor [140]. Moreover, the disruption of death receptor-dependent cell signaling is linked to poor survival in patients with HCC [141]. On the other hand, several genetic alterations of the TNF-TNFR superfamily in HCC were detected by sequencing HCC DNA samples. In particular, the TNFR1 promoter –329G/T polymorphism was strongly associated with primary HCC, where the T allele resulted in the repression of TNFR1 expression. Therefore, these results suggest that the TNFR1 329G/T polymorphism may play an important role in the development of HCC [142].

4.3. Viral interference in TNFR1 expression and function

TNF acting via TNFR1 is considered an important anti-viral agent, often acting synergistically with IFN γ [143]. However, growing evidence has shown that both DNA and RNA viruses can interfere with the TNFR1 pathway and thereby escape the host immune response. In infected cells, viral proteins can either affect TNFR1 availability by acting upon transcription, translation, trafficking or shedding of TNFR1, or modulate TNFR1 activity by acting on internalization or signaling of TNFR1 (see Table 5).

Epstein–Barr virus (EBV) immediate-early protein BZLF1 prevents TNF-induced activation of target genes and TNF-induced

apoptosis by down-regulation of TNFR1 during the EBV lytic replication cycle. Thus, EBV has developed a mechanism for evading TNF-induced antiviral effects during lytic reactivation or primary infection [144]. The **hepatitis C virus** (HCV) core protein is besides a component of viral nucleocapsids also a multifunctional protein influencing multiple cellular processes. In HCV core protein-activated Hep191 cells, transcriptional profiling identified decreased expression of TNFR1. Since RT-PCR confirmed that TNFR1 is down-regulated and that TNF-induced DNA fragmentation is suppressed in these cells, expression of HCV core protein at physiological levels might inhibit apoptotic cell death of HCV-infected cells [144].

Herpes simplex virus 1 (HSV) uses many strategies to inactivate host functions that are harmful to its replication and dissemination, including taking advantage of the short half-life of TNFR1. Steady-state levels of TNFR1 require continuous renewal by translation of its mRNA, and the HSV viral protein UL41 prevents this constant replenishment by degrading the TNFR1 mRNA [145].

Cytomegalovirus (CMV) is known to target the cell cycle, cellular transcription and immunoregulation to optimize the cellular environment for viral DNA replication. CMV infection also prevents external signaling to the cell by reducing the cell surface expression of TNFR1. Viral early gene products may be responsible for interfering with TNFR1 trafficking through the Golgi apparatus to the cell surface. So, upon infection CMV isolates the cell from host-mediated signals, forcing it to respond only to virus-specific signals [146,147]. **Poliovirus** also triggers host defensive reactions by activating intrinsic (intracellular) and extrinsic (receptor-mediated) apoptotic pathways. Poliovirus nonstructural protein 2A is an inhibitor of cellular translation enhancing the sensitivity to TNFR1-induced apoptosis. On the other hand, poliovirus nonstructural protein 3A neutralizes the pro-apoptotic activity of 2A by eliminating TNFR1 from the cell surface. Consequently, poliovirus infection dramatically decreases TNF receptor abundance on the surfaces of infected cells as early as four hours post-infection. Poliovirus-mediated resistance to TNF is caused by protein 3A interfering with protein trafficking through the endoplasmic reticulum and Golgi: the effect of protein 3A on TNF signaling can be imitated by brefeldin A [148].

Alveolar macrophages and respiratory epithelial cells infected with **respiratory syncytial virus** (RSV) suppress the production of early inflammatory cytokines such as TNF by producing IL10, resulting in an ineffective response to the virus [149]. Moreover, the soluble form of TNFR1, but not TNFR2, was secreted from these cells in a time- and RSV dose-dependent way. As the secretion of soluble TNFR1 blocks TNF responses, increased shedding might be another counteraction against the immune response [150].

The **adenovirus** protein E3-14.7K inhibits TNF-induced apoptosis by preventing TNF-induced TNFR1 internalization, which results in inhibition of the DISC formation. In contrast, E3-14.7K did not affect TNF-induced NF- κ B activation via recruitment of RIP-1 and TRAF-2. Inhibition of endocytosis by E3-14.7K is due to a failure in the coordinated temporal and spatial assembly of essential components of the endocytic machinery, such as Rab5 and dynamin2, at the site of the activated TNFR1. This is another mechanism by which Adenoviruses escape the host immune response [151].

Many poxviruses encode several immunomodulatory proteins, such as homologs of host cytokine receptors, also referred to as viroreceptors. These receptors mimic host function by binding to host cytokines, allowing the virus to circumvent the immune defense. The T2 protein of **myxoma virus** (M-T2) is a pseudo TNF receptor that has two distinct activities. The secreted dimeric M-T2 glycoprotein binds TNF with high affinity, inhibiting direct TNF-mediated cytotoxicity of infected cells and other secondary immune

responses dependent on TNF. However, intracellular M-T2 prevents myxoma-infected T-cells from undergoing apoptosis, supporting viral replication [152,153]. The cytokine response modifier (Crm)-like pseudo TNF receptors, including CrmB, CrmC, CrmD and CrmE and a putative fifth member from cowpox virus that closely resembles CD30, have been identified in several **orthopoxviruses**. These viral TNF receptors resemble secreted versions of the extracellular domains of their counterpart cellular receptors and form functional oligomers that bind and block TNF [154].

The latent membrane protein 1 (LMP1) of **Epstein–Barr virus** (EBV) contributes to the immortalizing activity of EBV in primary human B lymphocytes [155]. LMP1 is targeted to the plasma membrane of infected cells as a constitutive pseudo TNF receptor that activates NF- κ B through two independent domains in its cytoplasmic tail. One site is similar to TNFR2 interacting with TRAF1 and TRAF2 and the second site is similar to TNFR1 associating with TRADD. As LMP1 acts independently of the ligand, it replaces the T-cell-derived activation to maintain unlimited B-cell proliferation. LMP1-mediated signaling through the TRAF system plays a role in the pathogenesis of EBV-infected lymphomas that emerge in immunosuppressed patients [156]. Furthermore, the activity of the TNFR1 promoter is dramatically decreased by the EBV protein BZLF1, helping the virus to oppose the anti-viral effects of TNF [144]. Infection by **human herpesvirus 6B** (HHV-6B) blocks caspase 3 and 8 activation and I- κ B phosphorylation, indicating inhibition of both the inflammatory and apoptotic signaling pathways. The viral pseudo TNF receptor U20 was shown to localize to the cell membrane, and siRNA knockdown of U20 showed that the protein is necessary for HHV-6B-mediated inhibition of TNFR signaling during infection [157]. The STP pseudo TNF receptors of the **herpesvirus saimiri** (HVS) are stably associated with TRAF1, 2, or 3. Mutational analysis revealed that STP-C488 induces NF- κ B activation that correlates with its ability to associate with TRAFs. Thus, TRAF/STP association might be involved in immortalization of T lymphocytes following HVS infection [158]. High-risk strains of **human papillomavirus** (HPV) such as HPV16 cause human cervical carcinoma. The E6 protein of HPV16 mediates the rapid degradation of p53. But transfection of HPV16 E6 into the TNF-sensitive LM cell line protects the cells from TNF-induced apoptosis independently of p53. Caspase 3 and 8 activation is significantly reduced in E6-expressing cells, indicating that E6 acts early in the TNF apoptotic pathway. In fact, E6 binds directly to TNFR1. E6 requires the same TNFR1 C-terminal part for binding as does TRADD, and TNFR1/TRADD interactions are decreased in the presence of E6. HPV E6 binding to TNFR1 interferes with formation of the DISC and thus with transduction of pro-apoptotic signals. HPV, like several other viruses, can evade the TNF-mediated host immune response [159]. Moreover, the central conserved region of the attachment protein G of **respiratory syncytial virus** (RSV) shows structural homology with the fourth subdomain of TNFR1. Although the functions of both protein domains are unknown, the structural similarity of the two protein domains suggests that the cysteine noose of RSV-G may interfere with the antiviral and apoptotic effect mediated by TNF [160].

The **hepatitis C virus** (HCV) core protein binds to the cytoplasmic domain of TNFR1, namely the death domain (DD) thereby inhibiting TRADD binding but facilitating FADD recruitment to TNFR1 [161,162]. By sensitizing cells to TNF-mediated apoptosis, the HCV core protein may provide a selective advantage for HCV replication by enabling evasion of host antiviral defense mechanisms. In contrast, HCV core protein was shown to inhibit TNF-induced apoptosis via NF- κ B activation. The expression of the

core protein enhances nuclear retention and DNA binding activity of NF- κ B and TNF-triggered I- κ B degradation. This ability of HCV core protein to inhibit TNF-mediated apoptotic signaling may contribute to the chronically activated, persistent state of HCV-infected cells [163,164]. The controversy among these reports may be attributed to differences in cell types, conditions of core protein expression, and protocols of TNF stimulation [165,166].

Tanapoxvirus (TPV)-infected cells secrete an early 38-kDa glycopeptide that selectively inhibits TNF-induced NF- κ B activation and transcriptional activation of cell-adhesion molecules such as E-selectin, ICAM-1 and VCAM-1 on the surface of endothelial cells [167]. **Parvovirus** H-1 infection leads to activation of caspase 3, leading to morphologic changes characteristic of apoptosis and resembling the effects of TNF treatment. This effect is also observed when U937 cells are infected with a recombinant H-1 virus that expresses the nonstructural proteins but in which the capsid genes are replaced by a reporter gene, indicating that the induction of apoptosis can be assigned to the cytotoxic nonstructural proteins in this system. Furthermore, the c-Myc protein, which is over-expressed in the monocytoid cell line U937, is rapidly down-regulated during parvovirus infection, consistent with a possible role of c-Myc in mediating the apoptotic cell death induced by H-1 virus infection. Interestingly several clones derived from the U937 cell line and selected for their resistance to H-1 virus fail to decrease c-Myc expression upon treatment with differentiation agents, and they also resist the induction of cell death after TNF treatment [168,169].

5. Conclusion

TNFR1 mediates the signaling of most cellular effects of TNF. It is involved in numerous physiological and pathological functions at different levels and in different cell types. However, despite extensive knowledge of its signaling pathway, the precise TNF/TNFR1 activation mechanism remains unclear [170]. Moreover, there is still little knowledge about the transcriptional and translational regulation of TNFR1 expression. While the general assumption is that TNFR1 is constitutively expressed and regulated as a housekeeping gene, it is clearly sensitive to many stimuli. Predictions of several transcription factor binding sites and microRNA target sequences suggest that there are many unknown regulators. TNFR1 availability and localization are regulated by diverse mechanisms such as shedding, exosome formation, and internalization. Evidence for the importance of these and other mechanisms of TNFR1 control can be found in the many human diseases that are associated with TNFR1 regulation, levels and activity [42].

Since some of the side effects of the current TNF therapy might be absent with TNFR1 inhibition therapy, further increase of our knowledge on TNFR1 regulation might allow the development of new therapeutic strategies that more effectively target this interesting receptor and the pathogenic TNF signaling.

Conflict of interest

The authors have no conflicting financial interests.

Acknowledgements

We thank Dr. Amin Bredan for editing the manuscript. This research was supported by the Fund for Scientific Research, Flanders, the Belgian Interuniversity Attraction Poles program (IAP VI/18), the agency for Innovation by Science and Technology (IWT), VIB, and the Belgian Association Against Cancer.

References

- [1] Black RA, Rauch CT, Kozlosky CJ, Peschon JJ, Slack JL, Wolfson MF, et al. A metalloproteinase disintegrin that releases tumour-necrosis factor- α from cells. *Nature* 1997;385:729–33.
- [2] Hikita A, Tanaka N, Yamane S, Ikeda Y, Furukawa H, Tohma S, et al. Involvement of a disintegrin and metalloproteinase 10 and 17 in shedding of tumor necrosis factor- α . *Biochem Cell Biol* 2009;87:581–93.
- [3] Haro H, Crawford HC, Fingleton B, Shinomiya K, Spengler DM, Matrisian LM. Matrix metalloproteinase-7-dependent release of tumor necrosis factor- α in a model of herniated disc resorption. *J Clin Invest* 2000;105:143–50.
- [4] Becker-Pauls C, Rose-John S. TNF α cleavage beyond TACE/ADAM17: matrix metalloproteinase 13 is a potential therapeutic target in sepsis and colitis. *EMBO Mol Med* 2013;5(July (7)):902–4. <http://dx.doi.org/10.1002/emmm.201302899> [Epub 2013 Jun 11].
- [5] Fiers W. Tumor necrosis factor. Characterization at the molecular, cellular and in vivo level. *FEBS Lett* 1991;285:199–212.
- [6] Idriss HT, Naismith JH. TNF α and the TNF receptor superfamily: structure-function relationship(s). *Microsc Res Tech* 2000;50:184–95.
- [7] Chen G, Goeddel DV. TNF-R1 signaling: a beautiful pathway. *Science* 2002;296:1634–5.
- [8] Gullberg U, Lantz M, Lindvall L, Olsson I, Himmler A. Involvement of an Asn/Val cleavage site in the production of a soluble form of a human tumor necrosis factor (TNF) receptor. Site-directed mutagenesis of a putative cleavage site in the p55 TNF receptor chain. *Eur J Cell Biol* 1992;58:307–12.
- [9] Chan FK, Chun HJ, Zheng L, Siegel RM, Bui KL, Lenardo MJ. A domain in TNF receptors that mediates ligand-independent receptor assembly and signaling. *Science* 2000;288:2351–4.
- [10] Chan FK. The pre-ligand binding assembly domain: a potential target of inhibition of tumour necrosis factor receptor function. *Ann Rheum Dis* 2000;59(Suppl. 1):i50–3.
- [11] Ashkenazi A, Dixit VM. Death receptors: signaling and modulation. *Science* 1998;281:1305–8.
- [12] Chen ZJ. Ubiquitin signalling in the NF- κ B pathway. *Nat Cell Biol* 2005;7:758–65.
- [13] Tokunaga F, Iwai K. LUBAC, a novel ubiquitin ligase for linear ubiquitination, is crucial for inflammation and immune responses. *Microbes Infect* 2012;14:563–72.
- [14] Chang L, Kamata H, Solinas G, Luo JL, Maeda S, Venuprasad K, et al. The E3 ubiquitin ligase itch couples JNK activation to TNF α -induced cell death by inducing c-FLIP(L) turnover. *Cell* 2006;124:601–13.
- [15] Trompouki E, Hatzivassiliou E, Tschirritzis T, Farmer H, Ashworth A, Mosialos G. CYLD is a deubiquitinating enzyme that negatively regulates NF- κ B activation by TNFR family members. *Nature* 2003;424:793–6.
- [16] Wertz IE, O'Rourke KM, Zhou H, Eby M, Aravind L, Seshagiri S, et al. Deubiquitination and ubiquitin ligase domains of A20 downregulate NF- κ B signalling. *Nature* 2004;430:694–9.
- [17] Clark J, Vagenas P, Panesar M, Cope AP. What does tumour necrosis factor excess do to the immune system long term? *Ann Rheum Dis* 2005;64(Suppl. 4):iv70–6.
- [18] Favaloro B, Allocati N, Graziano V, Di Ilio C, De Laurenzi V. Role of apoptosis in disease. *Aging (Albany NY)* 2012;4:330–49.
- [19] Vandenabeele P, Galluzzi L, Vanden Berghe T, Kroemer G. Molecular mechanisms of necroptosis: an ordered cellular explosion. *Nat Rev Mol Cell Biol* 2010;11:700–14.
- [20] Baud V, Karin M. Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol* 2001;11:372–7.
- [21] Kollias G. TNF pathophysiology in murine models of chronic inflammation and autoimmunity. *Semin Arthritis Rheum* 2005;34:3–6.
- [22] Rothe J, Lesslauer W, Lotscher H, Lang Y, Koebele P, Kontgen F, et al. Mice lacking the tumour necrosis factor receptor 1 are resistant to TNF-mediated toxicity but highly susceptible to infection by *Listeria monocytogenes*. *Nature* 1993;364:798–802.
- [23] Pfeffer K, Matsuyama T, Kundig TM, Wakeham A, Kishihara K, Shahinian A, et al. Mice deficient for the 55 kd tumor necrosis factor receptor are resistant to endotoxic shock, yet succumb to *L. monocytogenes* infection. *Cell* 1993;73:457–67.
- [24] Peschon JJ, Torrance DS, Stocking KL, Glaccum MB, Otten C, Willis CR, et al. TNF receptor-deficient mice reveal divergent roles for p55 and p75 in several models of inflammation. *J Immunol* 1998;160:943–52.
- [25] Elicabe RJ, Arias JL, Rabinovich GA, Di Genaro MS. TNFRp55 modulates IL-6 and nitric oxide responses following Yersinia lipopolysaccharide stimulation in peritoneal macrophages. *Immunobiology* 2011;216:1322–30.
- [26] Elicabe RJ, Cargnelutti E, Serer MI, Stege PW, Valdez SR, Toscano MA, et al. Lack of TNFR p55 results in heightened expression of IFN- γ and IL-17 during the development of reactive arthritis. *J Immunol* 2010;185:4485–95.
- [27] Choi S, Park YS, Koga T, Treloar A, Kim KC. TNF- α is a key regulator of MUC1, an anti-inflammatory molecule, during airway *Pseudomonas aeruginosa* infection. *Am J Respir Cell Mol Biol* 2011;44:255–60.
- [28] Fujita M, Ikegame S, Harada E, Ouchi H, Inoshima I, Watanabe K, et al. TNF receptor 1 and 2 contribute in different ways to resistance to *Legionella pneumophila*-induced mortality in mice. *Cytokine* 2008;44:298–303.
- [29] Barnes JL, Williams NL, Ketheesan N. Susceptibility to *Burkholderia pseudomallei* is associated with host immune responses involving tumor necrosis factor receptor-1 (TNFR1) and TNF receptor-2 (TNFR2). *FEMS Immunol Med Microbiol* 2008;52:379–88.
- [30] Belisle SE, Tisoncik JR, Korth MJ, Carter VS, Proll SC, Swayne DE, et al. Genomic profiling of tumor necrosis factor α (TNF- α) receptor and interleukin-1 receptor knockout mice reveals a link between TNF- α signaling and increased severity of 1918 pandemic influenza virus infection. *J Virol* 2010;84:12576–88.
- [31] Tian T, Dubin K, Jin Q, Qureshi A, King SL, Liu L, et al. Disruption of TNF- α /TNFR1 function in resident skin cells impairs host immune response against cutaneous vaccinia virus infection. *J Invest Dermatol* 2012;132:1425–34.
- [32] Kettelhut IC, Fiers W, Goldberg AL. The toxic effects of tumor necrosis factor in vivo and their prevention by cyclooxygenase inhibitors. *Proc Natl Acad Sci U S A* 1987;84:4273–7.
- [33] Kontoyiannis D, Pasparakis M, Pizarro TT, Cominelli F, Kollias G. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity* 1999;10:387–98.
- [34] Stoelcker B, Ruhland B, Hehlhans T, Bluethmann H, Luther T, Mannel DN. Tumor necrosis factor induces tumor necrosis via tumor necrosis factor receptor type 1-expressing endothelial cells of the tumor vasculature. *Am J Pathol* 2000;156:1171–6.
- [35] Breedveld FC, Weisman MH, Kavanaugh AF, Cohen SB, Pavelka K, van Vollenhoven R, et al. The PREMIER study: a multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;54:26–37.
- [36] Klareskog L, van der Heijde D, de Jager JP, Gough A, Kalden J, Malaise M, et al. Therapeutic effect of the combination of etanercept and methotrexate compared with each treatment alone in patients with rheumatoid arthritis: double-blind randomised controlled trial. *Lancet* 2004;363:675–81.
- [37] Keystone EC. Does anti-tumor necrosis factor- α therapy affect risk of serious infection and cancer in patients with rheumatoid arthritis?: a review of longterm data. *J Rheumatol* 2011;38:1552–62.
- [38] de Gannes GC, Ghoreishi M, Pope J, Russell A, Bell D, Adams S, et al. Psoriasis and pustular dermatitis triggered by TNF- α inhibitors in patients with rheumatoid conditions. *Arch Dermatol* 2007;143:223–31.
- [39] Cleynen I, Vermeire S. Paradoxical inflammation induced by anti-TNF agents in patients with IBD. *Nat Rev Gastroenterol Hepatol* 2012;9:496–503.
- [40] Van Hauwermeiren F, Armaka M, Karagianni N, Kranidioti K, Vandenbroucke RE, Loges S, et al. Safe TNF-based anti-tumor therapy following p55TNFR reduction in intestinal epithelium. *J Clin Invest* 2013;123:2590–603.
- [41] Xanthoulea S, Pasparakis M, Kousteni S, Brakebusch C, Wallach D, Bauer J, et al. Tumor necrosis factor (TNF) receptor shedding controls thresholds of innate immune activation that balance opposing TNF functions in infectious and inflammatory diseases. *J Exp Med* 2004;200:367–76.
- [42] Kimberley FC, Lobito AA, Siegel RM, Sreter GR. Falling into TRAPS – receptor misfolding in the TNF receptor 1-associated periodic fever syndrome. *Arthritis Res Ther* 2007;9:217.
- [43] Van Hauwermeiren F, Vandenbroucke RE, Libert C. Treatment of TNF mediated diseases by selective inhibition of soluble TNF or TNFR1. *Cytokine Growth Factor Rev* 2011;22:311–9.
- [44] Zettlitz KA, Lorenz V, Landauer K, Munkel S, Herrmann A, Scheurich P, et al. ATROSAB, a humanized antagonistic anti-tumor necrosis factor receptor one-specific antibody. *MAbs* 2010;2:639–47.
- [45] Van Hauwermeiren F, Vandenbroucke RE, Grine L, Puimège L, Van Wouterghem E, Zhang H, et al. Antisense oligonucleotides against TNFR1 prevent toxicity of TNF/IFN γ treatment in mouse tumor models. *Int J Cancer* 2013. <http://dx.doi.org/10.1002/ijc.28704> [Epub ahead of print].
- [46] Kudo H, Takahara T, Yata Y, Kawai K, Zhang W, Sugiyama T. Lipopolysaccharide triggered TNF- α -induced hepatocyte apoptosis in a murine non-alcoholic steatohepatitis model. *J Hepatol* 2009;51:168–75.
- [47] Bertok S, Wilson MR, Morley PJ, de Wildt R, Bayliffe A, Takata M. Selective inhibition of intra-alveolar p55 TNF receptor attenuates ventilator-induced lung injury. *Thorax* 2012;67:244–51.
- [48] Shibata H, Yoshioka Y, Ohkawa A, Minowa K, Mukai Y, Abe Y, et al. Creation and X-ray structure analysis of the tumor necrosis factor receptor-1-selective mutant of a tumor necrosis factor- α antagonist. *J Biol Chem* 2008;283:998–1007.
- [49] Nomura T, Abe Y, Kamada H, Shibata H, Kayamuro H, Inoue M, et al. Therapeutic effect of PEGylated TNFR1-selective antagonistic mutant TNF in experimental autoimmune encephalomyelitis mice. *J Control Release* 2011;149:8–14.
- [50] Arntz OJ, Geurts J, Veenbergen S, Bennink MB, van den Brand BT, Abdollahi-Roodsaz S, et al. A crucial role for tumor necrosis factor receptor 1 in synovial lining cells and the reticuloendothelial system in mediating experimental arthritis. *Arthritis Res Ther* 2010;12:R61.
- [51] Shibata H, Yoshioka Y, Abe Y, Ohkawa A, Nomura T, Minowa K, et al. The treatment of established murine collagen-induced arthritis with a TNFR1-selective antagonistic mutant TNF. *Biomaterials* 2009;30:6638–47.
- [52] Shibata H, Yoshioka Y, Ohkawa A, Abe Y, Nomura T, Mukai Y, et al. The therapeutic effect of TNFR1-selective antagonistic mutant TNF- α in murine hepatitis models. *Cytokine* 2008;44:229–33.

- [53] Huang XW, Yang J, Dragovic AF, Zhang H, Lawrence TS, Zhang M. Antisense oligonucleotide inhibition of tumor necrosis factor receptor 1 protects the liver from radiation-induced apoptosis. *Clin Cancer Res* 2006;12:2849–55.
- [54] McFarlane SM, Pashmi G, Connell MC, Littlejohn AF, Tucker SJ, Vandenabeele P, et al. Differential activation of nuclear factor-kappaB by tumour necrosis factor receptor subtypes. TNFR1 predominates whereas TNFR2 activates transcription poorly. *FEBS Lett* 2002;515:119–26.
- [55] Fotin-Mleczek M, Henkler F, Samel D, Reichwein M, Hausser A, Parmryd I, et al. Apoptotic crosstalk of TNF receptors: TNF-R2-induces depletion of TRAF2 and IAP proteins and accelerates TNF-R1-dependent activation of caspase-8. *J Cell Sci* 2002;115:2757–70.
- [56] MacEwan DJ. TNF receptor subtype signalling: differences and cellular consequences. *Cell Signal* 2002;14:477–92.
- [57] Tartaglia LA, Pennica D, Goeddel DV. Ligand passing: the 75-kDa tumor necrosis factor (TNF) receptor recruits TNF for signaling by the 55-kDa TNF receptor. *J Biol Chem* 1993;268:18542–48.
- [58] Vandenabeele P, Declercq W, Beyaert R, Fiers W. Two tumour necrosis factor receptors: structure and function. *Trends Cell Biol* 1995;5:392–9.
- [59] McCoy MK, Tansey MG. TNF signaling inhibition in the CNS: implications for normal brain function and neurodegenerative disease. *J Neuroinflammation* 2008;5:45.
- [60] Takao S, Jacob CO. Mouse tumor necrosis factor receptor type I: genomic structure, polymorphism, and identification of regulatory regions. *Int Immunol* 1993;5:775–82.
- [61] Bristol JA, Morrison TE, Kenney SC. CCAAT/enhancer binding proteins alpha and beta regulate the tumor necrosis factor receptor 1 gene promoter. *Mol Immunol* 2009;46:2706–13.
- [62] Eckert D, Buhl S, Weber S, Jager R, Schorle H. The AP-2 family of transcription factors. *Genome Biol* 2005;6:246.
- [63] Baxter FO, Came PJ, Abell K, Kedjouar B, Huth M, Rajewsky K, et al. IKKbeta/2 induces TWEAK and apoptosis in mammary epithelial cells. *Development* 2006;133:3485–94.
- [64] Dopp JM, Mackenzie-Graham A, Otero GC, Merrill JE. Differential expression, cytokine modulation, and specific functions of type-1 and type-2 tumor necrosis factor receptors in rat glia. *J Neuroimmunol* 1997;75:104–12.
- [65] Agresti C, Bernardo A, Del Russo N, Marziali G, Battistini A, Aloisi F, et al. Synergistic stimulation of MHC class I and IRF-1 gene expression by IFN-gamma and TNF-alpha in oligodendrocytes. *Eur J Neurosci* 1998;10:2975–83.
- [66] Huang Y, Tan M, Gosink M, Wang KK, Sun Y. Histone deacetylase 5 is not a p53 target gene, but its overexpression inhibits tumor cell growth and induces apoptosis. *Cancer Res* 2002;62:2913–22.
- [67] Parry SL, Sebbag M, Feldmann M, Brennan FM. Contact with T cells modulates monocyte IL-10 production: role of T cell membrane TNF-alpha. *J Immunol* 1997;158:3673–81.
- [68] Dai SM, Matsuno H, Nakamura H, Nishioka K, Yudoh K. Interleukin-18 enhances monocyte tumor necrosis factor alpha and interleukin-1beta production induced by direct contact with T lymphocytes: implications in rheumatoid arthritis. *Arthritis Rheum* 2004;50:432–43.
- [69] Mathiasen IS, Hansen CM, Foghsgaard L, Jaattela M. Sensitization to TNF-induced apoptosis by 1,25-dihydroxy vitamin D(3) involves up-regulation of the TNF receptor 1 and cathepsin B. *Int J Cancer* 2001;93:224–31.
- [70] Yogesha SD, Khapli SM, Wani MR. Interleukin-3 and granulocyte-macrophage colony-stimulating factor inhibits tumor necrosis factor (TNF)-alpha-induced osteoclast differentiation by down-regulation of expression of TNF receptors 1 and 2. *J Biol Chem* 2005;280:11759–69.
- [71] Joyce DA, Gibbons DP, Green P, Steer JH, Feldmann M, Brennan FM. Two inhibitors of pro-inflammatory cytokine release, interleukin-10 and interleukin-4, have contrasting effects on release of soluble p75 tumor necrosis factor receptor by cultured monocytes. *Eur J Immunol* 1994;24:2699–705.
- [72] Williams L, Bradley L, Smith A, Foxwell B. Signal transducer and activator of transcription 3 is the dominant mediator of the anti-inflammatory effects of IL-10 in human macrophages. *J Immunol* 2004;172:567–76.
- [73] Broos S, Soete A, Hooghe B, Moran R, van Roy F, De Bleser P. PhysBinder: Improving the prediction of transcription factor binding sites by flexible inclusion of biophysical properties. *Nucleic Acids Res* 2013;41:W531–4.
- [74] Adeli K. Translational control mechanisms in metabolic regulation: critical role of RNA binding proteins, microRNAs, and cytoplasmic RNA granules. *Am J Physiol Endocrinol Metab* 2011;301:E1051–64.
- [75] Winzen R, Kafert S, Preiss B, Mylius-Spencker HA, Resch K, Holtmann H. Interaction between the mRNA of the 55-kDa tumor necrosis factor receptor and cellular proteins. Possible involvement in post-transcriptional regulation of receptor expression. *J Biol Chem* 1996;271:13461–67.
- [76] Kafert S, Winzen R, Loos A, Bollig F, Resch K, Holtmann H. Protein binding regions of the mRNAs for the 55 kDa tumor necrosis factor receptor and the glucose transporter 1: sequence homology and competition for cellular proteins. *FEBS Lett* 1998;421:2–6.
- [77] Hutvagner G, Zamore PD. A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 2002;297:2056–60.
- [78] Bartel DP, Chen CZ. Micromanagers of gene expression: the potentially widespread influence of metazoan microRNAs. *Nat Rev Genet* 2004;5:396–400.
- [79] Valencia-Sanchez MA, Liu J, Hannon GJ, Parker R. Control of translation and mRNA degradation by miRNAs and siRNAs. *Genes Dev* 2006;20:515–24.
- [80] Gaeta ML, Johnson DR, Kluger MS, Pober JS. The death domain of tumor necrosis factor receptor 1 is necessary but not sufficient for Golgi retention of the receptor and mediates receptor desensitization. *Lab Invest* 2000;80:1185–94.
- [81] Storey H, Stewart A, Vandenabeele P, Luzio JP. The p55 tumour necrosis factor receptor TNFR1 contains a trans-Golgi network localization signal in the C-terminal region of its cytoplasmic tail. *Biochem J* 2002;366:15–22.
- [82] Wang J, Al-Lamki RS, Zhang H, Kirkiles-Smith N, Gaeta ML, Thiru S, et al. Histamine antagonizes tumor necrosis factor (TNF) signaling by stimulating TNF receptor shedding from the cell surface and Golgi storage pool. *J Biol Chem* 2003;278:21751–60.
- [83] Bradley JR. TNF-mediated inflammatory disease. *J Pathol* 2008;214:149–60.
- [84] Aderka D, Engelmann H, Maor Y, Brakebusch C, Wallach D. Stabilization of the bioactivity of tumor necrosis factor by its soluble receptors. *J Exp Med* 1992;175:323–9.
- [85] Dreyer Mueller D, Pruessmeyer J, Groth E, Ludwig A. The role of ADAM-mediated shedding in vascular biology. *Eur J Cell Biol* 2012;91:472–85.
- [86] Gonzales PE, Solomon A, Miller AB, Leesnitzer MA, Sagi I, Milla ME. Inhibition of the tumor necrosis factor-alpha-converting enzyme by its pro domain. *J Biol Chem* 2004;279:31638–45.
- [87] Tellier E, Canault M, Rebsomen L, Bonardo B, Juhan-Vague I, Nalbone G, et al. The shedding activity of ADAM17 is sequestered in lipid rafts. *Exp Cell Res* 2006;312:3969–80.
- [88] Tellier E, Canault M, Poggi M, Bonardo B, Nicolay A, Alessi MC, et al. HDLs activate ADAM17-dependent shedding. *J Cell Physiol* 2008;214:687–93.
- [89] Soond SM, Everson B, Riches DW, Murphy G. ERK-mediated phosphorylation of Thr735 in TNFalpha-converting enzyme and its potential role in TACE protein trafficking. *J Cell Sci* 2005;118:2371–80.
- [90] Chen YJ, Chang LS. Arecoline-induced death of human leukemia K562 cells is associated with surface up-modulation of TNFR2. *J Cell Physiol* 2012;227:2240–51.
- [91] Reddy P, Slack JL, Davis R, Cerretti DP, Kozlosky CJ, Blanton RA, et al. Functional analysis of the domain structure of tumor necrosis factor-alpha converting enzyme. *J Biol Chem* 2000;275:14608–14.
- [92] Willems SH, Tape CJ, Stanley PL, Taylor NA, Mills IG, Neal DE, et al. Thiol isomerases negatively regulate the cellular shedding activity of ADAM17. *Biochem J* 2010;428:439–50.
- [93] Wang Y, Herrera AH, Li Y, Belani KK, Walcheck B. Regulation of mature ADAM17 by redox agents for L-selectin shedding. *J Immunol* 2009;182:2449–57.
- [94] Cui X, Hawari F, Alsaaty S, Lawrence M, Combs CA, Geng W, et al. Identification of ARTS-1 as a novel TNFR1-binding protein that promotes TNFR1 ectodomain shedding. *J Clin Invest* 2002;110:515–26.
- [95] Hawari FI, Rouhani FN, Cui X, Yu ZX, Buckley C, Kaler M, et al. Release of full-length 55-kDa TNF receptor 1 in exosome-like vesicles: a mechanism for generation of soluble cytokine receptors. *Proc Natl Acad Sci U S A* 2004;101:1297–302.
- [96] Cottin V, Van Linden A, Riches DW. Phosphorylation of tumor necrosis factor receptor CD120a (p55) by p42(mapk/erk2) induces changes in its subcellular localization. *J Biol Chem* 1999;274:32975–87.
- [97] Legler DF, Micheau O, Doucey MA, Tschopp J, Bron C. Recruitment of TNF receptor 1 to lipid rafts is essential for TNFalpha-mediated NF-kappaB activation. *Immunity* 2003;18:655–64.
- [98] Micheau O, Tschopp J. Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. *Cell* 2003;114:181–90.
- [99] Higuchi M, Aggarwal BB. TNF induces internalization of the p60 receptor and shedding of the p80 receptor. *J Immunol* 1994;152:3550–8.
- [100] Schutze S, Machleidt T, Adam D, Schwandner R, Wiegmann K, Kruse ML, et al. Inhibition of receptor internalization by monodansylcadaverine selectively blocks p55 tumor necrosis factor receptor death domain signaling. *J Biol Chem* 1999;274:10203–12.
- [101] Tchikov V, Bertsch U, Fritsch J, Edelmann B, Schutze S. Subcellular compartmentalization of TNF receptor-1 and CD95 signaling pathways. *Eur J Cell Biol* 2011;90:467–75.
- [102] Mosselmans R, Hepburn A, Dumont JE, Fiers W, Galand P. Endocytic pathway of recombinant murine tumor necrosis factor in L-929 cells. *J Immunol* 1988;141:3096–100.
- [103] Ethell DW, Bossy-Wetzel E, Bredesen DE. Caspase 7 can cleave tumor necrosis factor receptor-I (p60) at a non-consensus motif, in vitro. *Biochim Biophys Acta* 2001;1541:231–8.
- [104] Touitou I, Lesage S, McDermott M, Cuisset L, Hoffman H, Dode C, et al. Infefers: an evolving mutation database for auto-inflammatory syndromes. *Hum Mutat* 2004;24:194–8.
- [105] Huggins ML, Radford PM, McIntosh RS, Bainbridge SE, Dickinson P, Draper-Morgan KA, et al. Shedding of mutant tumor necrosis factor receptor superfamily 1A associated with tumor necrosis factor receptor-associated periodic fever syndrome: differences between cell types. *Arthritis Rheum* 2004;50:2651–9.
- [106] Lobito AA, Kimberley FC, Muppidi JR, Komarow H, Jackson AJ, Hull KM, et al. Abnormal disulfide-linked oligomerization results in ER retention and altered signaling by TNFR1 mutants in TNFR1-associated periodic fever syndrome (TRAPS). *Blood* 2006;108:1320–7.
- [107] D'Ossualdo A, Ferlito F, Prigione I, Obici L, Meini A, Zulian F, et al. Neutrophils from patients with TNFRSF1A mutations display resistance to tumor necrosis factor-induced apoptosis: pathogenetic and clinical implications. *Arthritis Rheum* 2006;54:998–1008.
- [108] Siebert S, Amos N, Fielding CA, Wang EC, Aksentijevich I, Williams BD, et al. Reduced tumor necrosis factor signaling in primary human fibroblasts containing a tumor necrosis factor receptor superfamily 1A mutant. *Arthritis Rheum* 2005;52:1287–92.

- [109] Tsukamoto H, Ueda N, Horiuchi T. Progress in classification and treatment for TNF receptor-associated periodic syndrome. *Nihon Rinsho Meneki Gakkai Kaishi* 2011;34:361–8.
- [110] Bulua AC, Mogul DB, Aksentijevich I, Singh H, He DY, Muenz LR, et al. Efficacy of etanercept in the tumor necrosis factor receptor-associated periodic syndrome: a prospective, open-label, dose-escalation study. *Arthritis Rheum* 2012;64:908–13.
- [111] McDermott MF, Aksentijevich I, Galon J, McDermott EM, Ogunkolade BW, Centola M, et al. Germline mutations in the extracellular domains of the 55 kDa TNF receptor, TNFR1, define a family of dominantly inherited auto-inflammatory syndromes. *Cell* 1999;97:133–44.
- [112] Offer M, Lachmann HJ, Bybee A, Rowczenio DM, Goodman HJ, Hawkins PN. Phenotype and response to treatment of 22 patients with autoinflammatory disease associated with mutations in TNFRSF1A. FMF and beyond. In: Fourth international conference on systemic inflammatory disorders; 2005.
- [113] Poirier O, Nicaud V, Gariépy J, Courbon D, Elbaz A, Morrison C, et al. Polymorphism R92Q of the tumour necrosis factor receptor 1 gene is associated with myocardial infarction and carotid intima-media thickness – the ECTIM, AXA, EVA and GENIC Studies. *Eur J Hum Genet* 2004;12:213–9.
- [114] Amoura Z, Dode C, Hue S, Caillat-Zucman S, Braham S, Delpech M, et al. Association of the R92Q TNFRSF1A mutation and extracranial deep vein thrombosis in patients with Behcet's disease. *Arthritis Rheum* 2005;52:608–11.
- [115] Caminero A, Comabella M, Montalban X. Role of tumour necrosis factor (TNF)-alpha and TNFRSF1A R92Q mutation in the pathogenesis of TNF receptor-associated periodic syndrome and multiple sclerosis. *Clin Exp Immunol* 2011;166:338–45.
- [116] Maschmeyer G, Haas A, Cornely OA. Invasive aspergillosis: epidemiology, diagnosis and management in immunocompromised patients. *Drugs* 2007;67:1567–601.
- [117] Sainz J, Salas-Alvarado I, Lopez-Fernandez E, Olmedo C, Comino A, Garcia F, et al. TNFR1 mRNA expression level and TNFR1 gene polymorphisms are predictive markers for susceptibility to develop invasive pulmonary aspergillosis. *Int J Immunopathol Pharmacol* 2010;23:423–36.
- [118] Glossop JR, Dawes PT, Hassell AB, Matthey DL. Anemia in rheumatoid arthritis: association with polymorphism in the tumor necrosis factor receptor I and II genes. *J Rheumatol* 2005;32:1673–8.
- [119] Dieude P, Osorio J, Petit-Teixeira E, Moreno S, Garnier S, Cailleau-Moindraut S, et al. A TNFR1 genotype with a protective role in familial rheumatoid arthritis. *Arthritis Rheum* 2004;50:413–9.
- [120] Jacobelli S, Andre M, Alexandra JF, Dode C, Papo T. Failure of anti-TNF therapy in TNF Receptor 1-Associated Periodic Syndrome (TRAPS). *Rheumatology (Oxford)* 2007;46:1211–2.
- [121] Nedjai B, Quillinan N, Coughlan RJ, Church L, McDermott MF, Hitman GA, et al. Lessons from anti-TNF biologics: infliximab failure in a TRAPS family with the T50M mutation in TNFRSF1A. *Adv Exp Med Biol* 2011;691:409–19.
- [122] Gregory AP, Dendrou CA, Attfield KE, Haghighi A, Xifara DK, Butter F, et al. TNF receptor 1 genetic risk mirrors outcome of anti-TNF therapy in multiple sclerosis. *Nature* 2011;488:508–11.
- [123] Rittore C, Sanchez E, Soler S, Barat-Houari M, Albers M, Obici L, et al. Identification of a new exon 2-skipped TNFR1 transcript: regulation by three functional polymorphisms of the TNFR-associated periodic syndrome (TRAPS) gene. *Ann Rheum Dis* 2014;73(1):290–7. <http://dx.doi.org/10.1136/annrheumdis-2012-203023> [Epub 2013 Mar 16].
- [124] Zhu LJ, Landolt-Marticorena C, Li T, Yang X, Yu XQ, Gladman DD, et al. Altered expression of TNF-alpha signaling pathway proteins in systemic lupus erythematosus. *J Rheumatol* 2010;37:1658–66.
- [125] Cook EB, Stahl JL, Graziano FM, Barney NP. Regulation of the receptor for TNFalpha, TNFR1, in human conjunctival epithelial cells. *Invest Ophthalmol Vis Sci* 2008;49:3992–8.
- [126] Aggarwal S, Gollapudi S, Gupta S. Increased TNF-alpha-induced apoptosis in lymphocytes from aged humans: changes in TNF-alpha receptor expression and activation of caspases. *J Immunol* 1999;162:2154–61.
- [127] Bao C, Guo J, Lin G, Hu M, Hu Z. TNFR gene-modified mesenchymal stem cells attenuate inflammation and cardiac dysfunction following MI. *Scand Cardiovasc J* 2008;42:56–62.
- [128] Kishore R, Tkebuchava T, Sasi SP, Silver M, Gilbert HY, Yoon YS, et al. Tumor necrosis factor-alpha signaling via TNFR1/p55 is deleterious whereas TNFR2/p75 signaling is protective in adult infarct myocardium. *Adv Exp Med Biol* 2011;691:433–48.
- [129] Elneihoum AM, Falke P, Axelsson L, Lundberg E, Lindgarde F, Ohlsson K. Leukocyte activation detected by increased plasma levels of inflammatory mediators in patients with ischemic cerebrovascular diseases. *Stroke* 1996;27:1734–8.
- [130] Iosif RE, Ahlenius H, Ekdahl CT, Darsalia V, Thored P, Jovinge S, et al. Suppression of stroke-induced progenitor proliferation in adult subventricular zone by tumor necrosis factor receptor 1. *J Cereb Blood Flow Metab* 2008;28:1574–87.
- [131] Shore SA, Williams ES, Chen L, Benedito LA, Kasahara DI, Zhu M. Impact of aging on pulmonary responses to acute ozone exposure in mice: role of TNFR1. *Inhal Toxicol* 2011;23:878–88.
- [132] Wu T, Xie C, Wang HW, Zhou XJ, Schwartz N, Calixto S, et al. Elevated urinary VCAM-1, P-selectin, soluble TNF receptor-1, and CXC chemokine ligand 16 in multiple murine lupus strains and human lupus nephritis. *J Immunol* 2007;179:7166–75.
- [133] Turan B, Pfister K, Diener PA, Hell M, Moller B, Boyvat A, et al. Soluble tumour necrosis factor receptors sTNFR1 and sTNFR2 are produced at sites of inflammation and are markers of arthritis activity in Behcet's disease. *Scand J Rheumatol* 2008;37:135–41.
- [134] Sfrikakis PP, Theodossiadis PG, Katsiari CG, Kaklamanis P, Markomichelakis NN. Effect of infliximab on sight-threatening panuveitis in Behcet's disease. *Lancet* 2001;358:295–6.
- [135] Sfrikakis PP. Behcet's disease: a new target for anti-tumour necrosis factor treatment. *Ann Rheum Dis* 2002;61(Suppl. 2):ii51–3.
- [136] Melikoglu M, Fresko I, Mat C, Ozyazgan Y, Gogus F, Yurdakul S, et al. Short-term trial of etanercept in Behcet's disease: a double blind, placebo controlled study. *J Rheumatol* 2005;32:98–105.
- [137] Menalled LB, Chesselet MF. Mouse models of Huntington's disease. *Trends Pharmacol Sci* 2002;23:32–9.
- [138] Lin CH, Tallaksen-Greene S, Chien WM, Cearley JA, Jackson WS, Crouse AB, et al. Neurological abnormalities in a knock-in mouse model of Huntington's disease. *Hum Mol Genet* 2001;10:137–44.
- [139] Strand AD, Aragaki AK, Shaw D, Bird T, Holton J, Turner C, et al. Gene expression in Huntington's disease skeletal muscle: a potential biomarker. *Hum Mol Genet* 2005;14:1863–76.
- [140] Gao L, Bai L, Nan QZ, Yang XS, Chen K, Wen HP, et al. Detection of soluble tumor necrosis factor-p55 levels in the serum and ascitic fluid of patients with hepatocellular carcinoma. *Di Yi Jun Yi Da Xue Xue Bao* 2002;22:641–3.
- [141] Muntane J. Targeting cell death and survival receptors in hepatocellular carcinoma. *Anticancer Agents Med Chem* 2011;11:576–84.
- [142] Kim S, Moon SM, Kim YS, Kim JJ, Ryu HJ, Kim YJ, et al. TNFR1 promoter –329G/T polymorphism results in allele-specific repression of TNFR1 expression. *Biochem Biophys Res Commun* 2008;368:395–401.
- [143] Reis LF, Ho Lee T, Vilcek J. Tumor necrosis factor acts synergistically with autocrine interferon-beta and increases interferon-beta mRNA levels in human fibroblasts. *J Biol Chem* 1989;264:16351–54.
- [144] Morrison TE, Mauser A, Klingelutz A, Kenney SC. Epstein-Barr virus immediate-early protein BZLF1 inhibits tumor necrosis factor alpha-induced signaling and apoptosis by downregulating tumor necrosis factor receptor 1. *J Virol* 2004;78:544–9.
- [145] Liang L, Roizman B. Herpes simplex virus 1 precludes replenishment of the short-lived receptor of tumor necrosis factor alpha by virion host shutoff-dependent degradation of its mRNA. *J Virol* 2006;80:7756–9.
- [146] Baillie J, Sahlender DA, Sinclair JH. Human cytomegalovirus infection inhibits tumor necrosis factor alpha (TNF-alpha) signaling by targeting the 55-kilodalton TNF-alpha receptor. *J Virol* 2003;77:7007–16.
- [147] Popkin DL, Virgin HW 4th. Murine cytomegalovirus infection inhibits tumor necrosis factor alpha responses in primary macrophages. *J Virol* 2003;77:10125–30.
- [148] Nezanov N, Kondratova A, Chumakov KM, Angres B, Zhumabayeva B, Agol VI, et al. Poliovirus protein 3A inhibits tumor necrosis factor (TNF)-induced apoptosis by eliminating the TNF receptor from the cell surface. *J Virol* 2001;75:10409–20.
- [149] Panuska JR, Merolla R, Rebert NA, Hoffmann SP, Tsivits P, Cirino NM, et al. Respiratory syncytial virus induces interleukin-10 by human alveolar macrophages. Suppression of early cytokine production and implications for incomplete immunity. *J Clin Invest* 1995;96:2445–53.
- [150] Arnold R, Humbert B, Werchau H, Gallati H, König W. Interleukin-8, interleukin-6, and soluble tumour necrosis factor receptor type I release from a human pulmonary epithelial cell line (A549) exposed to respiratory syncytial virus. *Immunology* 1994;82:126–33.
- [151] Schneider-Brachert W, Tchikov V, Merkel O, Jakob M, Hallas C, Kruse ML, et al. Inhibition of TNF receptor 1 internalization by adenovirus 14.7K as a novel immune escape mechanism. *J Clin Invest* 2006;116:2901–13.
- [152] Schreiber M, Rajarathnam K, McFadden G. Myxoma virus T2 protein, a tumor necrosis factor (TNF) receptor homolog, is secreted as a monomer and dimer that each bind rabbit TNFalpha, but the dimer is a more potent TNF inhibitor. *J Biol Chem* 1996;271:13333–41.
- [153] Schreiber M, Sedger L, McFadden G. Distinct domains of M-T2, the myxoma virus tumor necrosis factor (TNF) receptor homolog, mediate extracellular TNF binding and intracellular apoptosis inhibition. *J Virol* 1997;71:2171–81.
- [154] Johnston JB, McFadden G. Poxvirus immunomodulatory strategies: current perspectives. *J Virol* 2003;77:6093–100.
- [155] Izumi KM, Kaye KM, Kieff ED. The Epstein-Barr virus LMP1 amino acid sequence that engages tumor necrosis factor receptor associated factors is critical for primary B lymphocyte growth transformation. *Proc Natl Acad Sci U S A* 1997;94:1447–52.
- [156] Liebowitz D. Epstein-Barr virus and a cellular signaling pathway in lymphomas from immunosuppressed patients. *N Engl J Med* 1998;338:1413–21.
- [157] Kofod-Olsen E, Ross-Hansen K, Schleimann MH, Jensen DK, Moller JM, Bundgaard B, et al. U20 is responsible for human herpesvirus 6B inhibition of tumor necrosis factor receptor-dependent signaling and apoptosis. *J Virol* 2012;86:11483–92.
- [158] Lee H, Choi JK, Li M, Kaye K, Kieff E, Jung JU. Role of cellular tumor necrosis factor receptor-associated factors in NF-kappaB activation and lymphocyte transformation by herpesvirus Saimiri STP. *J Virol* 1999;73:3913–9.

- [159] Filippova M, Song H, Connolly JL, Dermody TS, Duerksen-Hughes PJ. The human papillomavirus 16 E6 protein binds to tumor necrosis factor (TNF) R1 and protects cells from TNF-induced apoptosis. *J Biol Chem* 2002;277:21730–39.
- [160] Langedijk JP, de Groot BL, Berendsen HJ, van Oirschot JT. Structural homology of the central conserved region of the attachment protein G of respiratory syncytial virus with the fourth subdomain of 55-kDa tumor necrosis factor receptor. *Virology* 1998;243:293–302.
- [161] Zhu N, Khoshnan A, Schneider R, Matsumoto M, Dennert G, Ware C, et al. Hepatitis C virus core protein binds to the cytoplasmic domain of tumor necrosis factor (TNF) receptor 1 and enhances TNF-induced apoptosis. *J Virol* 1998;72:3691–7.
- [162] Zhu N, Ware CF, Lai MM. Hepatitis C virus core protein enhances FADD-mediated apoptosis and suppresses TRADD signaling of tumor necrosis factor receptor. *Virology* 2001;283:178–87.
- [163] Marusawa H, Hijikata M, Chiba T, Shimotohno K. Hepatitis C virus core protein inhibits Fas- and tumor necrosis factor alpha-mediated apoptosis via NF-kappaB activation. *J Virol* 1999;73:4713–20.
- [164] You LR, Chen CM, Lee YH. Hepatitis C virus core protein enhances NF-kappaB signal pathway triggering by lymphotoxin-beta receptor ligand and tumor necrosis factor alpha. *J Virol* 1999;73:1672–81.
- [165] Nelson DR, Lim HL, Marousis CG, Fang JW, Davis GL, Shen L, et al. Activation of tumor necrosis factor-alpha system in chronic hepatitis C virus infection. *Dig Dis Sci* 1997;42:2487–94.
- [166] Kallinowski B, Haseroth K, Marinos G, Hanck C, Stremmel W, Theilmann L, et al. Induction of tumour necrosis factor (TNF) receptor type p55 and p75 in patients with chronic hepatitis C virus (HCV) infection. *Clin Exp Immunol* 1998;111:269–77.
- [167] Paulose M, Bennett BL, Manning AM, Essani K. Selective inhibition of TNF-alpha induced cell adhesion molecule gene expression by tanapox virus. *Microb Pathog* 1998;25:33–41.
- [168] Lopez-Guerrero JA, Rayet B, Tuynder M, Rommelaere J, Dinsart C. Constitutive activation of U937 promonocytic cell clones selected for their resistance to parvovirus H-1 infection. *Blood* 1997;89:1642–53.
- [169] Rayet B, Lopez-Guerrero JA, Rommelaere J, Dinsart C. Induction of programmed cell death by parvovirus H-1 in U937 cells: connection with the tumor necrosis factor alpha signalling pathway. *J Virol* 1998;72:8893–903.
- [170] Chan FK. Three is better than one: pre-ligand receptor assembly in the regulation of TNF receptor signaling. *Cytokine* 2007;37:101–7.
- [171] Eck MJ, Sprang SR. The structure of tumor necrosis factor-alpha at 2.6 Å resolution Implications for receptor binding. *J Biol Chem* 1989;264:17595–605.
- [172] Naismith JH, Devine TQ, Kohno T, Sprang SR. Structures of the extracellular domain of the type I tumor necrosis factor receptor. *Structure* 1996;4:1251–62.



Leen Puimège graduated as a master in biotechnology in 2009 and started as a PhD student in the lab of Prof. Claude Libert at Ghent University. She recently finished her PhD on the involvement of microRNAs in the regulation of TNFR1 in the inbred mouse strain Spret/Ei and the generation of a new TNFR1 blocking tools.



Claude Libert obtained his PhD in molecular biology in 1993 in the lab of Walter Fiers. After a postdoc in the IRBM in Rome, Italy, he became a group leader with VIB in 1997 and a professor at Ghent University in 2003. His main interest is the elucidation of molecular mechanism of complex acute inflammatory reactions, such as sepsis, and the identification of new players. His approach is a mouse molecular genetic approach and his aim is to define novel therapeutic interventions. Currently, his group consists of 14 researchers.



Filip Van Hauwermeiren graduated as a master in biomedical sciences in 2006 and obtained his PhD in 2011 in the lab of Prof. Claude Libert. He is mainly interested in the molecular mechanisms, responsible for TNF induced toxicity and pathology. Currently, he works as a FWO-funded postdoc in the lab of Prof. Mohamed Lamkanfi where he is investigating the role of NLRs in infections and inflammatory diseases.