

CHILDHOOD TRAUMA IN OBSESSIVE-COMPULSIVE DISORDER, TRICHOTILLOMANIA, AND CONTROLS

Christine Lochner, M.A.,* Pieter L. du Toit, M.A., Nompumelelo Zungu-Dirwayi, B.A., Adele Marais, M.A., Jeanine van Kradenburg, B.Curr., Soraya Seedat, M.Med., Dana J.H. Niehaus, M.Med., and Dan J. Stein, M.D., Ph.D.

There is relatively little data on the link between childhood trauma and obsessive-compulsive/putative obsessive-compulsive spectrum disorders. The revised Childhood Trauma Questionnaire (CTQ), which assesses physical, emotional, and sexual abuse as well as physical and emotional neglect, was administered to female patients with obsessive-compulsive disorder (OCD; $n = 74$; age: 36.1 ± 16.3), TTM ($n = 36$; age: 31.8 ± 12.3), and a group of normal controls ($n = 31$; age: 21.5 ± 1.0). The findings showed a significantly greater severity of childhood trauma in general, and emotional neglect specifically, in the patient groups compared to the controls. Although various factors may play a role in the etiology of both OCD and trichotillomania (TTM), this study is consistent with some evidence from previous studies suggesting that childhood trauma may play a role in the development of these disorders. Depression and Anxiety 15:66–68, 2002. © 2002 Wiley-Liss, Inc.

Key words: *obsessive-compulsive disorder; trichotillomania; childhood trauma*

INTRODUCTION

In recent years, there has been growing acceptance that early adverse experiences, including childhood physical and sexual abuse, may be associated with or contribute to psychiatric morbidity in adulthood [Bryer et al., 1987; Keaney and Farley, 1996; Kessler et al., 1997; Mulder et al., 1998; Stein et al., 1996; Walker et al., 1992]. An increased rate of childhood trauma has been reported in previous studies of anxiety disorder patients [e.g., Stein et al., 1996]. More specifically, panic disorder, agoraphobia, obsessive-compulsive disorder (OCD), and social anxiety disorder have been linked with a history of physical and sexual abuse during childhood [Burnam et al., 1988; David et al., 1995; Hofmann et al., 2001; Murrey et al., 1993; Saunders et al., 1992; Stein et al., 1996].

Although the etiology of OCD remains incompletely understood, there is clear evidence of a familial component [Pato et al., 2001; Pauls and Alsobrook, 1999], and a growing appreciation that specific neuro-anatomic, neurochemical, and neuroimmunological variables may mediate the pathogenesis of this disorder [e.g., Baxter, 1994; Leonard and Swedo, 2001; Micallef and Blin, 2001]. Freud's early contention that obsessional neurosis was a manifestation of premature sexual experience [Freud, 1909], has received little empirical study [Stein and Stone, 1997]. There is some anecdotal evidence of a link between severe trauma and onset of OCD [de Silva and Marks, 1999].

In recent studies, it has been suggested that several

disorders, including trichotillomania (TTM), lie on a spectrum of OCD disorders characterized by overlapping phenomenology and psychobiology [McElroy et al., 1994; Stein and Hollander, 1993]. While this hypothesis has received some support from neurobiological studies [Stein, 2000], little work has been done comparing different stressors (such as childhood trauma) across these disorders. Therefore, in this study, childhood trauma was assessed in patients with OCD or TTM, and healthy controls.

METHODS

Seventy-four patients with obsessive-compulsive disorder, 36 with trichotillomania and 31 healthy controls, took part in the study. Inclusion into this project was subject to informed written consent from all partici-

MRC Unit on Stress and Anxiety Disorders, University of Stellenbosch, Tygerberg, Cape Town, South Africa

Contract grant sponsor: Medical Research Council (MRC) of South Africa.

*Correspondence to: Christine Lochner, MRC Unit on Stress and Anxiety Disorders, University of Stellenbosch, PO Box 19063, Tygerberg 7505, Cape Town, South Africa.

E-mail: cl2@gerga.sun.ac.za

Received 3 June 2001; Revised 27 December 2001; Accepted 25 September 2001

Published online in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/da.10028

pants. As TTM patients are predominantly female [Christenson et al., 1991], the study was restricted to women, their ages ranging between 12 and 71. Patients were recruited from an Anxiety Disorders Clinic, while normal controls were recruited from a college class.

All patients met the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) criteria [American Psychiatric Association, 1994] for OCD on the Structured Clinical Interview of Axis I Disorders (SCID-I) [First et al., 1998], or TTM on the Structured Clinical Interview for the Diagnosis of OCD Spectrum Disorders (SCID-OCSD) [du Toit et al., 2001], as assessed by an experienced clinician. Controls were not screened and were included irrespective of possible diagnostic status/family history of psychiatric illness.

The Childhood Trauma Questionnaire (CTQ), a self-report instrument developed to provide brief, reliable, and valid assessment of a broad range of traumatic experiences in childhood [Bernstein et al., 1994], was completed by all participants. This questionnaire is comprised of 28 items that fall into five categories/subscales, namely: emotional abuse, physical abuse, sexual abuse, emotional neglect, and physical neglect. A sixth category is included as a validity scale (see Table 1). Reliability and validity of the CTQ has been well researched and the scale appears to be a useful measure of childhood trauma [Bernstein et al., 1994].

One-way analysis of variance (ANOVA) was used to compare total score on the CTQ as well as scores on the five subscales across the three groups.

RESULTS

Data were tabulated (Table 1). Mean age differed significantly between the groups ($F = 2.0$; $P = 0.003$); OCD and TTM patients were both significantly older than controls ($P < 0.001$).

ANOVA demonstrated that the total score on the CTQ differed significantly across the three groups ($F = 4.3$, $P = 0.015$). Both OCD ($t = -3.5$; $P = 0.001$) and

TTM ($t = -2.9$; $P = 0.005$) patients reported significantly more childhood trauma than controls.

There were significant differences between groups on two subscales in particular: the emotional neglect ($F = 6.8$, $P = 0.002$) and the physical abuse subscales ($F = 3.2$, $P = 0.04$). Emotional neglect was significantly higher in both OCD ($t = -4.5$; $P < 0.001$) and TTM ($t = -3.9$; $P < 0.001$) than in controls, with no significant difference between OCD and TTM. The physical abuse score was significantly higher in TTM than controls ($t = -3.0$; $P = 0.005$).

DISCUSSION

Childhood trauma in general was significantly higher in patients with OCD and TTM than in healthy controls. Both OCD and TTM patients scored higher than controls on emotional neglect, indicating that some aspects of childhood trauma may be common to both disorders. These findings seem to be consistent with previous research studies suggesting an association between anxiety disorders (such as OCD) and childhood trauma.

However, the association between childhood trauma and later OCD/TTM does not necessarily imply a causal relationship. Moreover, the findings here may not be generalizable to males. Variables such as age, comorbidity, and socio-economic status may also influence the data. Other limitations should also be considered: the information regarding childhood trauma was retrospective and self-reported, and could thus be subject to distortion.

Despite these limitations, we feel that the data presented here are useful insofar as they suggest that childhood trauma may influence the development of later anxiety disorders. Further research to explore this relationship, and to assess its specificity across the different anxiety disorders, is warranted.

REFERENCES

- American Psychiatric Association. 1994. Diagnostic and Statistical Manual of Mental Disorders. Washington D.C.: American Psychiatric Press.
- Baxter LR Jr. 1994. Positron emission tomography studies of cerebral glucose metabolism in obsessive compulsive disorder. *J Clin Psychiatry* 55: Suppl, 54-59.
- Bernstein DP, Fink L, Handelsman L, Foote J, Lovejoy M, Wenzel K, Sapareto E, Ruggiero J. 1994. Initial reliability and validity of a new retrospective measure of child abuse and neglect. *Am J Psychiatry* 151:1132-1136.
- Bryer JB, Nelson BA, Miller JB, Krol PA. 1987. Childhood sexual and physical abuse as factors in adult psychiatric illness. *Am J Psychiatry* 144:1426-1430.
- Burnam MA, Stein JA, Golding JM, Siegel JM, Sorenson SB, Forsythe AB, Telles CA. 1988. Sexual assault and mental disorders in a community population. *J Consult Clin Psychol* 56:843-850.
- Christenson GA, Mackenzie TB, Mitchell JE. 1991. Characteristics of 60 adult chronic hair pullers. *Am J Psychiatry* 148:365-370.
- David D, Giron A, Mellman TA. 1995. Panic-phobic patients and developmental trauma. *J Clin Psychiatry* 56:113-117.

TABLE 1. Childhood interpersonal trauma in OCD, TTM, and controls

	OCD ^a n = 74	TTM n = 36	Normal controls n = 31	P
Age ± SD	36.1 ± 16.3	31.8 ± 12.3	21.5 ± 1.0	.003
Age range	12-71 yrs	13-71 yrs	20-24 yrs	
Emotional abuse	10.3 ± 5.5	9.9 ± 5.6	7.8 ± 3.6	NS
Physical abuse	7.2 ± 3.8	7.9 ± 4.2	5.8 ± 1.1	.04
Sexual abuse	7.3 ± 4.5	7.1 ± 5.3	6.4 ± 3.3	NS
Emotional neglect	10.1 ± 5.1	10.3 ± 4.8	6.7 ± 2.5	.002
Physical neglect	5.9 ± 1.7	6.5 ± 2.2	5.6 ± 1.2	NS
Validity scale	0.8 ± 1.1	0.8 ± 1.1	1.0 ± 1.1	NS
CTQ total	40.8 ± 15.6	41.8 ± 16.8	32.4 ± 8.7	.015

^aOCD, obsessive-compulsive disorder; TTM, trichotillomania; CTQ, Childhood Trauma Questionnaire; NS, not significant.

- du Toit PL, van Kradenburg J, Niehaus D, Stein DJ. 2001. Comparison of obsessive-compulsive disorder patients with and without comorbid putative obsessive-compulsive spectrum disorders using a structured clinical interview. *Compr Psychiatry* 42:291–300.
- First MB, Spitzer RL, Gibbon M, Williams JBW. 1998. Structured clinical interview for DSM-IV Axis I disorders—Patient edition (SCID-I/P, Version 2.0, 8/98 revision). New York: New York State Psychiatric Institute, Biometrics Research Department.
- Freud S. 1909. Three case histories. New York: Macmillan.
- Hofmann SG, Levitt JT, Hoffman EC, Greene K, Litz BT, Barlow DH. 2001. Potentially traumatizing events in panic disorder and other anxiety disorders. *Depress Anxiety* 13:101–102.
- Keaney JC, Farley M. 1996. Dissociation in an outpatient sample of women reporting childhood sexual abuse. *Psychol Rep* 78: 59–65.
- Kessler RC, Davis CG, Kendler KS. 1997. Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. *Psychol Med* 27:1101–1119.
- Leonard HL, Swedo SE. 2001. Paediatric autoimmune neuropsychiatric disorders associated with streptococcal infection (PANDAS). *Int J Neuropsychopharmacol* 4:191–198.
- McElroy SL, Phillips KA, Keck PE Jr. 1994. Obsessive compulsive spectrum disorder. *J Clin Psychiatry* 55: Suppl, 33–51.
- Micallef J, Blin O. 2001. Neurobiology and clinical pharmacology of obsessive-compulsive disorder. *Clin Neuropharmacol* 24: 191–207.
- Mulder RT, Beautrais AL, Joyce PR, Fergusson DM. 1998. Relationship between dissociation, childhood sexual abuse, childhood physical abuse, and mental illness in a general population sample. *Am J Psychiatry* 155:806–811.
- Murrey GJ, Bolen J, Miller N, Siemsted K, Robbins M, Truskowski F. 1993. History of childhood sexual abuse in women with depressive and anxiety disorders: a comparative study. *J Sex Ed Therapy* 19:13–19.
- Pato MT, Schindler KM, Pato CN. 2001. The genetics of obsessive-compulsive disorder. *Curr Psychiatry Rep* 3:163–168.
- Pauls DL, Alsobrook JP. 1999. The inheritance of obsessive-compulsive disorder. *Child Adolesc Psychiatr Clin N Am* 8:481–496, viii.
- Saunders BE, Villopontaux LA, Lipovsky JA, Kilpatrick DG, Veronen LJ. 1992. Child sexual assault as a risk factor for mental disorders among women: a community survey. *J Interpers Violence* 7:189–204.
- Stein DJ. 2000. Neurobiology of the obsessive-compulsive spectrum disorders. *Biol Psychiatry* 47:296–304.
- Stein DJ, Hollander E. 1993. The spectrum of obsessive-compulsive related disorders. In: Hollander E, editor. *Obsessive-compulsive related disorders*. Washington D.C.: American Psychiatric Press.
- Stein DJ, Stone MH. 1997. Essential papers on obsessive-compulsive disorder. New York: New York University Press. p 4.
- Stein MB, Walker JR, Anderson G, Hazen AL, Ross CA, Eldridge G, Forde DR. 1996. Childhood physical and sexual abuse in patients with anxiety disorders and in a community sample. *Am J Psychiatry* 153:275–277.
- Walker EA, Katon WJ, Hansom J, Harrop-Griffiths J, Holm L, Jones ML, Hickok L, Jemelka RP. 1992. Medical and psychiatric symptoms in women with childhood sexual abuse. *Psychosom Med* 54:658–664.