

Throughout this paper the word "algebra" signifies an associative algebra over a fixed  $F$  of characteristic zero.

A variety of algebras  $\mathcal{M}$  is called a nonmatrix variety if the algebra of second-order matrices  $M_2$  does not belong to  $\mathcal{M}$ . The main example of nonmatrix varieties are the following:

$\mathcal{N}_\kappa$ , the variety of all nilpotent algebras of index  $\leq \kappa$ ;

$\mathcal{A}_0$ , the variety of all commutative algebras;

$\mathcal{A}_1 = \text{Var}(G)$ , where  $G$  is a Grassmann algebra of countable rank;

$\mathcal{A}_2 = \text{Var}(G \otimes G)$ .

If  $\mathcal{A}, \mathcal{B}, \mathcal{M}$  are varieties and  $\mathcal{A}, \mathcal{B} \subseteq \mathcal{M}$ , then  $\mathcal{A} \circ_{\mathcal{M}} \mathcal{B}$  denotes the  $\mathcal{M}$ -product of varieties [1], i.e., the intersection of the usual product  $\mathcal{A} \circ \mathcal{B}$  with  $\mathcal{M}$ .

In the present paper we prove that any nonmatrix variety  $\mathcal{M}$  can be "assembled" from the above-mentioned varieties by means of the operation  $\circ_{\mathcal{M}}$ ; more precisely, we prove the following results:

THEOREM 1. Suppose  $\mathcal{M}$  is a nonmatrix variety,  $\mathcal{M} \neq G \otimes G$ , and  $\mathcal{A}$  is the largest variety in  $\{\mathcal{O}, \mathcal{A}_0, \mathcal{A}_1\}$  ( $\mathcal{O}$  is the zero variety) lying in  $\mathcal{M}$ . Then for some  $\kappa$  we have

$$\mathcal{M} = \mathcal{M}_\kappa \circ_{\mathcal{M}} \mathcal{A},$$

where  $\mathcal{M}_\kappa = \mathcal{M} \cap \mathcal{N}_\kappa$ .

THEOREM 2. Suppose  $\mathcal{M}$  is a nonmatrix variety and  $\mathcal{M} \ni G \otimes G$ . Then for certain  $\kappa, \ell$  we have

$$\mathcal{M} = \mathcal{M}_\kappa \circ_{\mathcal{M}} (\mathcal{A}_2 \circ_{\mathcal{M}} (\mathcal{M}_\ell \circ_{\mathcal{M}} \mathcal{A}_0)).$$

THEOREM 3. Suppose  $\mathcal{M}$  is a nonmatrix variety,  $\mathcal{A}$  is a finitely based variety, and  $\mathcal{A} \supseteq \mathcal{A}_2$ . Then for some  $\kappa$  we have

$$\mathcal{M} = \mathcal{M}_\kappa \circ_{\mathcal{M}} (\mathcal{A} \cap \mathcal{M}).$$

It is not known whether the variety  $\mathcal{A}_2$  is finitely based, but if it is, then, by Theorem 3, the conclusion of Theorem 2 can be rewritten in the form  $\mathcal{M} = \mathcal{M}_\kappa \circ_{\mathcal{M}} \mathcal{A}_2$ . Note also that the varieties  $\mathcal{N}_2, \mathcal{A}_0, \mathcal{A}_1, \mathcal{A}_2$  are indecomposable; hence Theorems 1 and 2 give a decomposition of an arbitrary nonmatrix variety  $\mathcal{M}$  into an  $\mathcal{M}$ -product of indecomposable varieties.

Translated from Algebra i Logika, Vol. 19, No. 3, pp. 255-283, May-June, 1980. Original article submitted July 9, 1979.

From Theorems 1, 2, and 3 we obtain:

COROLLARY 1. The algebras of an arbitrary nonmatrix variety satisfy the identity

$$[[x_1, y_1], [x_1, t_1], u_1] \cdot \dots \cdot [[x_n, y_n], [x_n, t_n], u_n] = 0$$

for some  $n$ .

COROLLARY 2. If  $\mathcal{M}$  is a nonmatrix variety and  $\mathcal{M} \not\subseteq G \otimes G$ , then  $\mathcal{M}$  satisfies the identity

$$[x_1, y_1, z_1] \cdot \dots \cdot [x_n, y_n, z_n] = 0 \quad (*)$$

for some  $n$ .

Latyshev [2] studied the varieties of algebras satisfying an identity of the form

$$[x_1^{(n)}, \dots, x_m^{(n)}, [y^{(n)}, z^{(n)}]] \cdot \dots \cdot [x_1^{(k)}, \dots, x_m^{(k)}, [y^{(k)}, z^{(k)}]] = 0. \quad (**)$$

It is easy to show that an identity of the form (\*\*) implies an identity of the form (\*); hence, in view of Corollary 2, V. N. Latyshev's class of varieties coincides with the class of nonmatrix varieties that do not contain  $G \otimes G$ . Therefore in view of the main result of [2], we obtain

COROLLARY 3. If  $\mathcal{M}$  is a nonmatrix variety and  $\mathcal{M} \not\subseteq G \otimes G$ , then  $\mathcal{M}$  is Spechtian.

From Corollary 3 we obtain

COROLLARY 4. The variety of algebras satisfying a nontrivial identity of the form

$$\sum_{i=1}^n \alpha_i x^i y x^{n-i} = 0,$$

where  $\alpha_i \in F$ , is Spechtian.

It follows easily from Corollary 1 that any nonmatrix variety satisfies a Lie solvability identity. From this fact and a theorem of Higgins [3] we obtain

COROLLARY 5. If an associative algebra satisfies the identity  $[x, y, \dots, y]_k = 0$  for some  $k$ , then it satisfies the identity  $[x, x_2, \dots, x_m]$  for some  $m$ .

The proof of Corollary 5, obtained independently of [3], provides an affirmative answer to a question of Latyshev [4].

From Corollary 3 we also obtain

COROLLARY 6. If  $\mathcal{M}$  is locally weakly Noetherian, then  $\mathcal{M}$  is Spechtian.

From Corollary 5 we obtain for locally weakly Noetherian varieties the following

COROLLARY 7. If  $\mathcal{M}$  is locally weakly Noetherian, then  $\mathcal{M}$  satisfies an identity of the form

$$[x_1, \dots, x_m] \cdot y_1 \cdot \dots \cdot y_n [x_1, \dots, x_m] = 0.$$

# 1. PRELIMINARIES

Suppose  $X$  is a fixed countable linearly ordered set and  $F\langle X \rangle$  is the free associative algebra (without unity) generated by  $X$ . Represent  $X$  in the form

$$X = Y \cup \left( \bigcup_{i=1}^{\infty} T_i \right),$$

where  $T_i, Y$  are countable, pairwise disjoint subsets. Fix the sets  $T_i, Y$ . Let

$$T = \bigcup_{i=1}^{\infty} T_i.$$

Consider the ideal  $\mathcal{Y}$  of  $F\langle X \rangle$  generated by all elements of the form

$$\alpha_1 u \alpha_2 + \alpha_2 u \alpha_1,$$

where  $\alpha_1, \alpha_2 \in T_i$ ,  $i=1, 2, \dots$ ;  $u \in F\langle X \rangle \cup \{1\}$ . The images of the sets  $T_i, T$  under the natural homomorphism  $F\langle X \rangle \rightarrow F\langle X \rangle / \mathcal{Y}$  will be denoted by  $E_i, E$ , respectively. Obviously,

$$E = \bigcup_{i=1}^{\infty} E_i.$$

The image of the set  $Y$  will again be denoted by  $Y$ . We denote the quotient algebra  $F\langle X \rangle / \mathcal{Y}$  by  $F_{E,Y}$ . Thus, the algebra  $F_{E,Y}$  has the following defining relations:

$$\alpha_1 u \alpha_2 = -\alpha_2 u \alpha_1, \quad (1)$$

where  $\alpha_1, \alpha_2 \in E_i$ ,  $i=1, 2, \dots$ ;  $u \in F_{E,Y} \cup \{1\}$ .

Suppose  $A \subseteq X$ . Consider the linear operator  $S_A; F\langle X \rangle \rightarrow F\langle X \rangle$  defined as follows:

1. If  $f$  is a monomial and  $\deg_a f \geq 2$  for some  $a \in A$  ( $\deg_a f$  is the degree of the polynomial  $f$  with respect to the variable  $a$ ), then  $S_A(f) = 0$ .
2. If  $f = f(a_1, \dots, a_n, x_1, \dots, x_m)$  is a monomial,  $\alpha_i \in A, x_i \notin A$ ;  $\alpha_1 < \alpha_2 < \dots < \alpha_n$ ;  $\deg_{\alpha_i} f = 1$ ;  $n \geq 0$ , then

$$S_A(f) = \sum_{\sigma \in S(n)} (-1)^{\sigma} f(a_{\sigma(n)}, \dots, a_{\sigma(1)}, x_1, \dots, x_m),$$

where  $S(n)$  is the symmetric group of degree  $n$ . The operator  $S_A$  will be called the symmetrizer with respect to the set  $A$ .

It is easy to verify the following properties of symmetrizers:

If  $A \cap B = \emptyset$ , then

$$S_A(S_B(f)) = S_B(S_A(f)). \quad (2)$$

If  $f$  is homogeneous and linear with respect to the variables in  $A$ , then

$$S_A(S_A(f)) = n! S_A(f), \quad (3)$$

where  $n$  is the number of variables in  $A$  on which  $f$  depends.

The operator  ${}_T S: F\langle X \rangle \rightarrow F\langle X \rangle$  is defined as follows: if the polynomial  $f$  does not depend on the variables in  $T_i$  for  $i > \kappa$ , then  ${}_T S(f) = S_T \dots S_{T_\kappa}(f)$ .

Put  ${}_T \mathcal{S}(F\langle X \rangle) = M_{T,Y}$ . The natural homomorphism  $F\langle X \rangle \rightarrow F_{E,Y}$  induces a homomorphism of spaces  $\varphi: M_{T,Y} \rightarrow F_{E,Y}$ .

LEMMA 1.  $\varphi$  is an isomorphism.

Proof. Suppose  $f \in \text{Ker } \varphi = M_{T,Y} \cap \mathcal{I}$ . Since the spaces  $M_{T,Y}, \mathcal{I}$  are homogeneous and  $f \in M_{T,Y}$ , we may assume that  $f = {}_T \mathcal{S}(g)$ , where  $g$  is a polynomial that is multilinear with respect to the variables in  $T$ . Then in view of (2) and (3) we have

$${}_T \mathcal{S}(f) = {}_T \mathcal{S}({}_T \mathcal{S}(g)) = \mathcal{S}_{T_1} \mathcal{S}_{T_2} \dots \mathcal{S}_{T_k} \mathcal{S}_{T_k}(g) = n_1! \dots n_k! f, \quad (4)$$

where  $n_i$  is the number of variables in  $T_i$  on which  $g$  depends ( $n_i = 0$  for  $i > k$ ). On the other hand, it is easy to see that  ${}_T \mathcal{S}(\mathcal{I}) = 0$ . Therefore,  $f \in \mathcal{I}$  implies  ${}_T \mathcal{S}(f) = 0$ . It follows from this and (4) that  $f = 0$ .

It remains to show that  $M_{T,Y} + \mathcal{I} = F\langle X \rangle$ . Suppose  $f$  is an arbitrary monomial. If  $\deg_t f \geq 2$  for some  $t \in T$ , then, by definition of the ideal  $\mathcal{I}$ , we have  $f \in \mathcal{I}$ . In the opposite case, it is easy to see, using (1), that  ${}_T \mathcal{S}(f) = n_1! \dots n_k! f \pmod{\mathcal{I}}$ , where  $n_i$  is the number of variables in  $T_i$  occurring in  $f$ .

The lemma is proved.

If  $T$  is an arbitrary  $T$ -ideal of the algebra  $F\langle X \rangle$  and  $A$  is an arbitrary algebra, we denote by  $\Gamma(A)$  the ideal of values of the polynomials of  $T$  on  $A$ , i.e., the ideal generated by the elements  $f(a_1, \dots, a_n)$ , where  $f \in T, a_i \in A$ .

LEMMA 2. Suppose  $T$  is an arbitrary  $T$ -ideal of  $F\langle X \rangle$ . Then

$$\varphi(T \cap M_{T,Y}) = \Gamma(F_{E,Y}).$$

Proof. Suppose  $f = f(x_1, \dots, x_n) \in T \cap M_{T,Y}$ . Then  $\varphi(f) = f(\varphi(x_1), \dots, \varphi(x_n)) \in \Gamma(F_{E,Y})$ . Conversely, suppose  $g \in \Gamma(F_{E,Y})$ . We may assume that  $g$  is homogeneous; but then  $T$  contains a homogeneous polynomial  $f = f(t_1^{(n)}, t_2^{(n)}, \dots, t_k^{(n)}, y_1, \dots, y_s), t_i^{(n)} \in T_i, y_i \in Y$ , such that  $g = f(e_1^{(n)}, \dots, e_n^{(n)}, \dots, e_1^{(k)}, \dots, e_{n_k}^{(k)}, y_1, \dots, y_s), e_i^{(n)} \in E_j, y_i \in Y$ . Put

$$h = \frac{1}{n_1! \dots n_k!} {}_T \mathcal{S}(f).$$

Obviously,  $h \in T \cap M_{T,Y}$ . Using (1), it is easy to show that  $\varphi(h) = g$ .

The lemma is proved.

Definition 1. Let  $Z$  be the subspace of  $F\langle X \rangle$  generated by the set  $X$ . An ideal  $I$  of the algebra  $F\langle X \rangle$  will be called an  $\mathcal{S}$ -ideal if for any  $f(x_1, \dots, x_n) \in I$  and any  $z_1, \dots, z_n \in Z$  we have  $f(z_1, \dots, z_n) \in I$ , i.e., if  $I$  is stable under linear substitutions.

It is clear that any  $\mathcal{S}$ -ideal is stable under linearizations, hence is generated and defined by a set of multilinear polynomials lying in  $I$ .

Definition 2. Suppose  $A$  is an algebra and  $Z$  is a subspace of  $A$  generating  $A$  as a ring. An ideal of  $F\langle X \rangle$  of the form  $T[A, Z] = \{f(x_1, \dots, x_n) \in F\langle X \rangle \mid \forall z_1, \dots, z_n \in Z f(z_1, \dots, z_n) = 0\}$  will be called the ideal of identities of the pair  $(A, Z)$ .

It is clear that  $\mathcal{T}[A, Z]$  is an S-ideal and contains the ideal of identities of the algebra  $A$ .

The subalgebra of  $F_{E, Y}$  generated by the set

$$\bigcup_{i=1}^t E_i \cup \{y_1, \dots, y_r\}$$

will be denoted by  $F_{t, r}$ .

Proposition 1. For any T-ideal  $\Gamma$  there exist natural numbers  $t, r$  such that  $\Gamma = \mathcal{T}[A, Z]$ , where  $A = F_{t, r} / \Gamma(F_{t, r})$  and  $Z$  is the space generated by the classes of elements

$$\bigcup_{i=1}^t E_i \cup \{y_1, \dots, y_r\}.$$

Proof. We will make use of the representation theory of the symmetric group  $S(n)$ . For the details of this theory see [5-7]. The facts we need are all given in [8].

Let  $P_n$  be the space of multilinear polynomials of degree  $n$  in the variables  $x_1, \dots, x_n \in X$ . Put  $\Gamma_n = \Gamma \cap P_n$ ,  $d_n = \dim_F P_n / \Gamma_n$ . It is known [9] that for some natural number  $\alpha > 1$  we have

$$d_n \leq \alpha^n \quad (5)$$

for any  $n$ .

The dimension over  $F$  of a minimal left  $FS(n)$ -submodule of the left  $FS(n)$ -module  $P_n$  corresponding to the Young table  $\mathcal{D}$  will be denoted by  $b_{\mathcal{D}}$ . A formula for  $b_{\mathcal{D}}$  can be found in [5, 7, 8].

If  $p, q$  are natural numbers, we denote by  $\mathcal{D}_{p, q}$  the Young table of degree  $pq$  consisting of  $p$  equal columns.

Put  $t = \alpha^2 + 1$  and choose  $r$  so that

$$b_{\mathcal{D}_{t, r+1}} > \alpha^{2t(r+1)}. \quad (6)$$

Such a choice is possible, since  $\lim_{q \rightarrow \infty} \frac{b_{\mathcal{D}_{p, q}}}{(p-1)^{pq}} = \infty$  (this can be proved by using the formula for  $b_{\mathcal{D}}$  and Stirling's formula).

We will prove that  $t, r$  are the desired natural numbers.

Let  $B_n = \mathcal{T}[A, Z] \cap P_n$ . Since  $\mathcal{T}[A, Z], \Gamma$  are S-ideals, it follows that  $B_n, \Gamma_n$  are left  $FS(n)$ -submodules of the module  $P_n$  and the S-ideals  $\mathcal{T}[A, Z], \Gamma$  are defined by these submodules. Therefore, it suffices to prove that  $B_n = \Gamma_n$  for any  $n$ . But since  $P_n$  is a completely reducible  $FS(n)$ -module, it suffices to show that for any  $n$  and any table  $\mathcal{D}$  of degree  $n$  we have

$$\Gamma_n \cap \mathcal{U}_{\mathcal{D}} = B_n \cap \mathcal{U}_{\mathcal{D}}, \quad (7)$$

where  $\mathcal{U}_{\mathcal{D}}$  is the homogeneous component of  $P_n$  corresponding to the table  $\mathcal{D}$ .

We begin the proof of (7). Suppose  $\mathcal{D}$  is an arbitrary table. There are only two possible cases.

Case 1. The table  $\mathcal{D}$  contains the table  $\mathcal{D}_{t, t+1}$ . We will show that  $\Gamma_n \cap \mathcal{U}_{\mathcal{D}} = \mathcal{U}_{\mathcal{D}}$ . Since  $\Gamma \subseteq T[A, Z]$ , equality (7) will follow.

Suppose first that  $n \leq 2t(t+1)$ . The formula (see [7])  $b_{\mathcal{D}} = \sum_{\mathcal{D}'} b_{\mathcal{D}'}$ , where the sum extends over all tables  $\mathcal{D}'$  of degree  $n-1$  that are contained in  $\mathcal{D}$ , implies  $b_{\mathcal{D}} \geq b_{\mathcal{D}_{t, t+1}}$ . Therefore, if  $\mathcal{U}_{\mathcal{D}} \not\subseteq \Gamma_n$ , it follows from (6) that

$$d_n = \dim_F P_n / \Gamma_n \geq \dim_F \mathcal{U}_{\mathcal{D}} / \mathcal{U}_{\mathcal{D}} \cap \Gamma_n \geq b_{\mathcal{D}} \geq b_{\mathcal{D}_{t, t+1}} > a^{2t(t+1)} \geq a^n,$$

which contradicts (5). Therefore  $\mathcal{U}_{\mathcal{D}} \subseteq \Gamma_n$ .

Suppose  $f(x_1, \dots, x_{t(t+1)}) = \sum_{\sigma \in S(t(t+1))} \alpha_{\sigma} x_{\sigma(1)} \dots x_{\sigma(t(t+1))}$  is an arbitrary polynomial in  $\mathcal{U}_{\mathcal{D}_{t, t+1}}$ . We will show that any polynomial of the form

$$g(x_1, \dots, x_n) = \sum_{\sigma \in S(t(t+1))} \alpha_{\sigma} x_{\sigma(1)} y_1 x_{\sigma(2)} y_2 \dots x_{\sigma(t(t+1))} y_{t(t+1)}, \quad (8)$$

where  $y_i \in \{x_{t(t+1)+1}, \dots, x_n\}$ ,  $y_i \neq y_j$ , belongs to  $\Gamma$ . Indeed, for some permutation  $\tau \in S(n)$  we have  $g(x_1, \dots, x_n) = (f(x_1, \dots, x_{t(t+1)}) x_{t(t+1)+1} \dots x_n) \tau$  (the action of  $\tau \in S$  on  $P_n$  from the right is defined by  $(x_{j_1} x_{j_2} \dots x_{j_n}) \tau = x_{j_{\tau(1)}} \dots x_{j_{\tau(n)}}$ ). It follows from Proposition 1 of [8] that

$$g \in \bigoplus_{\mathcal{D}' \supset \mathcal{D}_{t, t+1}} \mathcal{U}_{\mathcal{D}'} \tau = \bigoplus_{\mathcal{D}' \supset \mathcal{D}_{t, t+1}} \mathcal{U}_{\mathcal{D}'} \subseteq \Gamma.$$

If we make the substitution  $x_i = u_i$  in (8), where  $i > t(t+1)$ ,  $u_i \in F \langle X \rangle \cup \{1\}$ , we obtain

$$\sum_{\sigma \in S(t(t+1))} \alpha_{\sigma} x_{\sigma(1)} u_1 x_{\sigma(2)} u_2 \dots x_{\sigma(t(t+1))} u_{t(t+1)} \in \Gamma. \quad (9)$$

Now suppose that  $n$  is an arbitrary number and  $h$  is any polynomial in  $\mathcal{U}_{\mathcal{D}}$  generating a minimal submodule. It follows from Remarks 1 and 2 of [8] that  $h$  can be represented as a linear combination of polynomials of the form (8), where  $y_i \in F \langle X \rangle \cup \{1\}$ ,

$$\sum_{\sigma \in S(t(t+1))} \alpha_{\sigma} x_{\sigma(1)} \dots x_{\sigma(t(t+1))} \in \mathcal{U}_{\mathcal{D}_{t, t+1}},$$

which, in view of (9), lie in  $\Gamma$ . Therefore,  $h$ , hence also  $\mathcal{U}_{\mathcal{D}}$ , is contained in  $\Gamma$ . Equality (7) is proved.

Case 2. The table  $\mathcal{D}$  does not contain the table  $\mathcal{D}_{t, t+1}$ . Since  $\Gamma \subseteq T[A, Z]$ , it suffices to prove that

$$\mathcal{U}_{\mathcal{D}} \cap B_n \subseteq \mathcal{U}_{\mathcal{D}} \cap \Gamma_n. \quad (10)$$

Suppose  $\mathcal{D}^*$  is an arbitrary diagram of the table  $\mathcal{D}$  (i.e., some filling in of the cells of  $\mathcal{D}$  by the integers from 1 to  $n$ ), and  $P_{\mathcal{D}^*}$  (respectively,  $Q_{\mathcal{D}^*}$ ) is the subgroup of  $S(n)$  consisting of all permutations fixing all rows (respectively, columns) of the diagram  $\mathcal{D}^*$ . Put

$$e_{\mathcal{D}^*} = \sum_{\rho \in P_{\mathcal{D}^*}} \rho \cdot \sum_{q \in Q_{\mathcal{D}^*}} (-1)^q q.$$

It follows from the representation theory of the group  $S(n)$  that the minimal ideal  $V_{\mathcal{D}}$  of  $FS(n)$ , corresponding to the table  $\mathcal{D}$ , is generated as left ideal by the elements  $e_{\mathcal{D}^*}$ . Therefore, the module  $\mathcal{U}_{\mathcal{D}} \cap \mathcal{B}_n$  is generated by the elements  $e_{\mathcal{D}^*} f$ , where  $f \in \mathcal{U}_{\mathcal{D}} \cap \mathcal{B}_n$ .

Let  $\mathcal{L}_i$  (respectively,  $N_i$ ) be the set of numbers appearing in the  $i$ -th column (respectively, row) of the diagram  $\mathcal{D}^*$ . Since  $\mathcal{D}$  does not contain the table  $\mathcal{D}_{t, z+1}$ , we can choose integers  $\kappa, \ell$  such that

$$0 \leq \kappa \leq t, \quad 0 \leq \ell \leq z, \quad (11)$$

$$\bigcup_{i=1}^{\kappa} \mathcal{L}_i \cup \bigcup_{i=1}^{\ell} N_i = \{1, \dots, n\}. \quad (12)$$

Consider the subgroups  $\bar{P}_{\mathcal{D}^*}^{\circ} = \{\rho \in P_{\mathcal{D}^*} \mid \rho(j) = j \quad \forall j \in \bigcup_{i=1}^{\kappa} \mathcal{L}_i\}$ ,  $Q_{\mathcal{D}^*}^{\circ} = \{q \in Q_{\mathcal{D}^*} \mid q(j) = j \quad \forall j \notin \bigcup_{i=1}^{\kappa} \mathcal{L}_i\}$ .

We have

$$e_{\mathcal{D}^*} = \sum_{\rho \in P'} \rho' \cdot \sum_{\rho \in P_{\mathcal{D}^*}^{\circ}} \rho \cdot \sum_{q \in Q_{\mathcal{D}^*}^{\circ}} (-1)^q q \cdot \sum_{q' \in Q'} \alpha_{q, q'},$$

where  $P'$  (respectively,  $Q'$ ) is a set of representatives of the left (respectively, right) cosets of  $P_{\mathcal{D}^*}^{\circ}$  ( $Q_{\mathcal{D}^*}^{\circ}$ ) modulo the subgroup  $P_{\mathcal{D}^*}^{\circ}$  ( $Q_{\mathcal{D}^*}^{\circ}$ );  $\alpha_{q, q'} = \pm 1$ . Therefore, the module  $\mathcal{U}_{\mathcal{D}} \cap \mathcal{B}_n$  is generated by the elements

$$h_{\mathcal{D}^*, f} = \left( \sum_{\rho \in P_{\mathcal{D}^*}^{\circ}} \rho \cdot \sum_{q \in Q_{\mathcal{D}^*}^{\circ}} (-1)^q q \right) f,$$

where  $f \in \mathcal{U}_{\mathcal{D}} \cap \mathcal{B}_n$ .

Suppose  $f = f(x_{i_1, 1}, \dots, x_{i_{\kappa}, \kappa}, x_{j_1, 1}, \dots, x_{j_{\ell}, \ell})$ , where  $\{i_1, \dots, i_{\kappa}\} = \mathcal{L}_S$ ,  $\{j_1, \dots, j_{\ell}\} = N_S \setminus \bigcup_{i=1}^{\kappa} \mathcal{L}_i$ . In view of (12), such a representation is possible. It is easy to see that the polynomial  $h_{\mathcal{D}^*, f}$  is, to within sign and a renaming of the variables, a linearization of the polynomial

$$g = s_{\tau_1} s_{\tau_2} \dots s_{\tau_{\kappa}} (f(t_1^{(1)}, \dots, t_{\kappa_1}^{(1)}, \dots, t_1^{(\kappa)}, \dots, t_{\kappa_{\ell}}^{(\kappa)}, \underbrace{y_1, \dots, y_{\kappa_1}}_{m_1}, \dots, \underbrace{y_{\ell_1}, \dots, y_{\ell_{\ell}}}_{m_{\ell}})).$$

Since  $\mathcal{T}[A, Z]$  is an S-ideal, we have  $g \in \mathcal{T}[A, Z]$ ; but then, in view of the definition of  $\mathcal{T}[A, Z]$  and (11), we obtain

$$h = g|_{t_i^{(j)} = e_i^{(j)}, j=1, \dots, \kappa; i=1, \dots, n_j} \in \mathcal{T}(F_{t, z}),$$

where  $e_i^{(j)} = \varphi(t_i^{(j)})$  ( $\varphi$  is the canonical isomorphism  $M_{\tau, y} \rightarrow F_{E, y}$ ). Therefore, by Lemma 2,  $\varphi^{-1}(h) \in \mathcal{T} \cap M_{\tau, y} \subseteq \mathcal{T}$ . Since  $g \in M_{\tau, y}$ ,  $\varphi(g) = h$ , and  $\varphi$  is an isomorphism, it follows that  $g \in \mathcal{T} \cap M_{\tau, y} \subseteq \mathcal{T}$ . Therefore,  $h_{\mathcal{D}^*, f} \in \mathcal{T}_n$ . This proves (10), hence also (11).

The proposition is proved.

## 2. ALGEBRA $G_2$

Let  $I$  denote the  $S$ -ideal generated by the polynomials

$$[x_1, x_2, x_3], \quad (13)$$

$$\sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] [x_{\sigma(2)}, y_2] [x_{\sigma(3)}, y_3], \quad (14)$$

where  $x_i, y_j \in X$ . Put  $G_2 = F\langle X \rangle / I$ . In view of the definition, this algebra is defined by the relations

$$[x_1, x_2, x_3] = 0, \quad (15)$$

$$\sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] [x_{\sigma(2)}, y_2] [x_{\sigma(3)}, y_3] = 0, \quad (16)$$

where  $x_i, y_j$  are arbitrary generators.

In this section we will show that the algebra  $G_2$  is an analog in some sense of the algebra  $G \otimes G$ .

LEMMA 3. Suppose  $G^*$  is a Grassmann algebra with 1. Then the algebra  $G^* \otimes G^*$  is a homomorphic image of  $G_2$ .

Proof. Suppose  $e_i$  are generators of the algebra  $G^*$ . We will identify the element  $1 \otimes 1$  with 1, and  $e_i \otimes 1$  with  $e_i$ . We denote the element  $1 \otimes e_i$  by  $f_i$ . The algebra  $G^* \otimes G^*$  is generated by the countable set  $\{e_1, e_2, \dots\} \cup \{f_1, f_2, \dots\} \cup \{1\}$ , and we need only verify relations (15) and (16) for the generators; this verification is immediate.

The lemma is proved.

A polynomial  $f \in F\langle X \rangle$  will be called a commutator polynomial if  $f$  is homogeneous in all variables and can be represented as a linear combination of polynomials of the form  $[y_1, y_2] \cdots [y_{2k-1}, y_{2k}]$ , where  $y_i \in X$ .

LEMMA 4. Suppose  $A$  is an  $S$ -ideal and  $A \supseteq I$ . If  $A$  contains a commutator polynomial  $g$  that is not in  $I$ , then  $A$  contains a commutator polynomial  $f$  such that

1.  $f \notin I$ ;
2.  $f = \sum_{\sigma \in S(k)} (-1)^\sigma h(x_{\sigma(1)}, \dots, x_{\sigma(k)}, y_1, \dots, y_k)$ , where  $h$  is a commutator polynomial, and  $x_i, y_j$  are pairwise distinct elements of  $X$ ;
3.  $\deg_{x_i} f = 1$ ,  $\deg_{y_i} f = 2$ .

Proof. Let  $P_n$  be the left  $FS(n)$ -module of multilinear polynomials in the variables  $x_1, \dots, x_n$ . Since  $A, I$  are  $S$ -ideals, it follows that  $A_n = A \cap P_n$  and  $I_n = I \cap P_n$  are submodules. We may assume that  $g \in P_n$  and generates a minimal submodule  $K$ . Let  $u_g$  be the homogeneous component of  $P_n$  containing  $K$ . We will show that the table  $\mathcal{D}$  contains at most two columns.

Suppose this is not so. As we observed in the proof of Proposition 1,  $K$  is generated by the polynomials  $e_{\mathcal{D}^*} g$ , where  $\mathcal{D}^*$  is an arbitrary diagram of  $D$  and



$$e_{\mathcal{D}^*} = \sum_{\rho \in \mathcal{P}_{\mathcal{D}^*}} \rho \cdot \sum_{g \in Q_{\mathcal{D}^*}} (-1)^g g.$$

Let  $\mathcal{P}'$  be the subgroup of  $\mathcal{P}_{\mathcal{D}^*}$  consisting of the permutations fixing any number except for the first three (for definiteness we assume these three are 1, 2, 3) appearing in the first row. Then

$$e_{\mathcal{D}^*} = \sum_{\rho \in \mathcal{P}'} \rho \cdot \sum_{\sigma \in \mathcal{S}(n)} \alpha_{\sigma} \sigma$$

for certain  $\alpha_{\sigma} \in F$ . Put

$$g_1 = \left( \sum_{\sigma \in \mathcal{S}(n)} \alpha_{\sigma} \sigma \right) g.$$

It is easy to see that  $g_1$  is a commutator polynomial,  $g_1 \in K$ , and

$$e_{\mathcal{D}^*} g = \left( \sum_{\rho \in \mathcal{P}'} \rho \right) g_1(x_1, \dots, x_n) = \frac{|\mathcal{P}'|}{3} \sum_{\sigma \in \mathcal{S}(3)} g_1(x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}, x_4, \dots, x_n).$$

Also, using relations (16) and (15), it is easy to see that the right-hand side is equal to 0 in  $G_2$ , i.e.,  $e_{\mathcal{D}^*} g \in I$ . Therefore,  $K \subseteq I$ , which contradicts the choice of  $g$ .

It follows from what has been proved and from Remarks 1 and 2 of [8] that  $K$  is generated by a polynomial that is a linearization of one of the form

$$h = [\delta_{\kappa+\ell}(y_1, \dots, y_{\ell}, x_1, \dots, x_{\kappa}) \delta_{\ell}(y_1, \dots, y_{\ell})] \left( \sum_{\sigma \in \mathcal{S}(\kappa+2\ell)} \alpha_{\sigma} \sigma \right),$$

where  $\delta_{\kappa}(x_1, \dots, x_{\kappa})$  is a standard polynomial and  $\alpha_{\sigma} \in F$ . Putting

$$f = \sum_{\sigma \in \mathcal{S}(K)} (-1)^{\sigma} h(x_{\sigma(1)}, \dots, x_{\sigma(K)}, y_1, \dots, y_{\ell}),$$

we have  $f = \kappa! h$ . Therefore, a linearization of  $f$  generates  $K$ , hence  $f \notin I$ . Since  $K$  is generated by a commutator polynomial, it follows that all polynomials in  $K$ , and also  $h$  and  $f$ , are commutator polynomials.

The lemma is proved.

**LEMMA 5.** Suppose  $A$  is an  $\mathcal{S}$ -ideal and  $A \supseteq I$ . If  $A$  contains a commutator polynomial  $f$  that is not in  $I$ , then  $A$  contains a polynomial of the form  $[x_1, t_1]^2 \dots [x_{\rho}, t_{\rho}]^2$ , where  $x_i, t_i$  are pairwise distinct variables in  $X$ .

Proof. In view of Lemma 4, we may assume that  $f$  possesses properties 1-3 of Lemma 4.

The proof is by induction on the degree of  $f$ . The basis of the induction,  $\deg f = 2$ , is obvious.

Suppose

$$\deg f > 2, \quad f = \sum_{\sigma \in \mathcal{S}(K)} (-1)^{\sigma} h(x_{\sigma(1)}, \dots, x_{\sigma(K)}, y_1, \dots, y_{\ell}).$$

There are only three possibilities.

Case 1.  $\ell \geq 2$ . We will show that

$$f \equiv [y_{\ell-1}, y_\ell]^2 f_1(x_1, \dots, x_k, y_1, \dots, y_{\ell-2}) \pmod{I}, \quad (17)$$

where  $f_1$  is a commutator polynomial. Indeed, since  $f$  is a commutator polynomial, it follows from (15) that  $f$  can be represented as a linear combination of polynomials of the form

$$[x_1, x_2][x_3, x_4][x_5, x_6][x_7, x_8] g_x, \quad (18)$$

where  $x_i \in \{x_1, \dots, x_k, y_1, \dots, y_\ell\}$ , and  $g_x$  is a commutator polynomial not depending on  $y_{\ell-1}, y_\ell$ . We may assume that  $x_1 = x_3 = y_{\ell-1}$  (otherwise we use (15)). We may also assume that  $x_2 = y_\ell$ . For if  $x_2 \neq y_\ell$ , but  $x_4 = y_\ell$ , then we can use (15). If  $x_2 \neq y_\ell$  and  $x_4 \neq y_\ell$ , then we can assume  $x_6 = x_8 = y_\ell$ . Using (15) and (16), we obtain

$$\begin{aligned} [y_{\ell-1}, x_2][y_{\ell-1}, x_4][x_5, y_\ell][x_7, y_\ell] &\stackrel{(16)}{\equiv} -[y_{\ell-1}, x_2][y_{\ell-1}, y_\ell][x_5, x_4][x_7, y_\ell] - \\ &- [y_{\ell-1}, x_2][y_{\ell-1}, y_\ell][x_5, y_\ell][x_7, x_4] \stackrel{(15)}{\equiv} -[y_{\ell-1}, y_\ell][y_{\ell-1}, x_2][x_5, x_4][x_7, y_\ell] - \\ &- [y_{\ell-1}, y_\ell][y_{\ell-1}, x_2][x_5, y_\ell][x_7, x_4] \pmod{I}, \end{aligned}$$

from which it follows that the polynomial (18) can be represented modulo  $I$  as a linear combination of polynomials of the form (18) in which  $x_1 = x_3 = y_{\ell-1}, x_2 = y_\ell$ . If  $x_4 \neq y_\ell$ , then, as above, we may assume  $x_6 = y_\ell$ . Modulo  $I$  we have the congruences

$$\begin{aligned} [y_{\ell-1}, y_\ell][y_{\ell-1}, x_4][x_5, y_\ell] &\stackrel{(16)}{\equiv} -[y_{\ell-1}, x_4][y_{\ell-1}, y_\ell][x_5, y_\ell] - [y_{\ell-1}, y_\ell]^2 [x_5, x_4] \stackrel{(15)}{\equiv} \\ &\stackrel{(15)}{\equiv} -[y_{\ell-1}, y_\ell][y_{\ell-1}, x_4][x_5, y_\ell] - [y_{\ell-1}, y_\ell]^2 [x_5, x_4] \pmod{I}. \end{aligned}$$

Therefore

$$[y_{\ell-1}, y_\ell][y_{\ell-1}, x_4][x_5, y_\ell] \equiv -\frac{1}{2} [y_{\ell-1}, y_\ell]^2 [x_5, x_4] \pmod{I}.$$

Congruence (17) is proved.

Let  $A_l$  be the  $\mathcal{S}$ -ideal generated by the set  $\{f_i\} \cup I$ . Using (15) and (17), it is easy to see that  $[y_{\ell-1}, y_\ell]^2 A_l \subseteq A$ . Since  $f_i \notin I$  and  $\deg f_i < \deg f$ , it follows from the induction assumption that the assertion of the lemma holds for  $A_l$ , hence also, in view of what was said above, for  $A$ .

Case 2.  $\ell = 1$ . We will show that this case is impossible. Indeed, for some  $\alpha \in F$  we have

$$\begin{aligned} f &\equiv \alpha \sum_{\sigma \in S(k)} (-1)^\sigma [x_{\sigma(1)}, y_1][x_{\sigma(2)}, y_1][x_{\sigma(3)}, x_{\sigma(4)}] \dots [x_{\sigma(k-1)}, x_{\sigma(k)}] = \\ &= \frac{\alpha}{2} \sum_{\sigma \in S(k)} (-1)^\sigma [[x_{\sigma(1)}, y_1], [x_{\sigma(2)}, y_1]][x_{\sigma(3)}, x_{\sigma(4)}] \dots [x_{\sigma(k-1)}, x_{\sigma(k)}] \equiv 0 \pmod{I}. \end{aligned}$$

Case 3.  $\ell = 0$ . Here we may assume that  $f = \mathcal{S}_{2g}(x_1, \dots, x_{2g})$ . Since  $A$  is an  $\mathcal{S}$ -ideal we have  $g = \mathcal{S}_{2g}(y_1, y_2, x_3, \dots, x_{2g})(y_1, y_2) \in A$ , where  $x_i, y_j \in X$ . If  $g \notin I$ , then  $g$  satisfies the

conditions of the first case and  $g = [y_1, y_2]^2 f_1(x_3, \dots, x_{2q}) \pmod{I}$ , where  $f_1 \notin I$ . If  $A_1$  is the  $\mathcal{S}$ -ideal generated by the set  $\{f_1\} \cup I$ , then, using (15), it is easy to show that  $[y_1, y_2]^2 A_1 \subseteq A$ . Since  $\deg f_1 < \deg f$ , the assertion of the lemma follows from the induction assumption.

Thus, it remains to prove that  $g \notin I$ . In view of Lemma 3, it suffices to show that the algebra  $G^* \otimes G^*$  satisfies the relation

$$\sum_{\sigma, \tau \in \mathcal{S}(2)} S_{2q}(y_{\sigma(1)}, x_{\sigma(1)}, x_3, \dots, x_{2q}) [y_{\tau(2)}, x_{\tau(2)}] \neq 0 \quad (19)$$

for certain  $y_i, z_j, x_s \in \{e_1, e_2, \dots\} \cup \{f_1, f_2, \dots\}$  (the left-hand side of (19) is a linearization of the polynomial  $g$ ). Put  $y_1 = e_1, y_2 = f_1, x_1 = e_2, x_2 = f_2, x_i = e_i$ . Then the left-hand side of (19) is equal to

$$\begin{aligned} & S_{2q}(e_1, \dots, e_{2q}) [f_1, f_2] + S_{2q}(f_1, f_2, e_3, \dots, e_{2q}) [e_1, e_2] = 2(2q)! e_1 \dots e_{2q} f_1 f_2 + \\ & + 4(2q-2)! \left( \sum_{j=0}^{2q-2} \sum_{i=0}^j (-1)^{i+j} e_1 \dots e_i f_1 e_{i+1} \dots e_j f_2 e_{j+1} \dots e_{2q-2} e_{2q-1} e_{2q} \right) = \\ & = [2(2q)! + 4q(2q-2)!] e_1 \dots e_{2q} f_1 f_2 \neq 0. \end{aligned}$$

The lemma is proved.

If  $A$  is an algebra, we denote by  $T[A]$  the ideal of identities of  $A$ . Let  $\Gamma_2 = T[G \otimes G]$ , where  $G$  is a Grassmann algebra of countable rank, and let  $\Gamma_1 = T[G]$ .

Proposition 2.  $T[G_2] = \Gamma_2$ .

Proof. It is well known that  $\Gamma_1 = \{[x, y, z]\}^T$ . Therefore, since  $G^*$  (a Grassmann algebra with 1) satisfies the identity  $[x, y, z] = 0$ , it follows that  $T[G^*] = \Gamma_1$ . It is now easy to see that  $T[G^* \otimes G^*] = \Gamma_2$ . Therefore, it suffices to prove that  $T[G^* \otimes G^*] = T[G_2]$ .

Let  $\mathcal{L}$  be the subspace of  $G^*$  generated by the elements  $1 \otimes 1, e_i = e_i \otimes 1, f_i = 1 \otimes e_i$ . Consider the  $\mathcal{S}$ -ideal of identities  $I_1 = T[G^* \otimes G^*, \mathcal{L}]$  of the pair  $(G^* \otimes G^*, \mathcal{L})$  (see Definition 2). Obviously,  $T[G^* \otimes G^*]$  is the largest  $T$ -ideal contained in  $I_1$  and  $T[G_2]$  is the largest  $T$ -ideal contained in  $I$ . Therefore, it suffices to prove that  $I_1 = I$ .

It follows from Lemma 3 that  $I_1 \supseteq I$ . Suppose  $f \in I_1, f \notin I$ . Since  $I_1, I$  are  $\mathcal{S}$ -ideals, we may assume that  $f$  is a multilinear polynomial in the variables  $x_1, \dots, x_n$ .

If  $A = \{x_{i_1}, \dots, x_{i_k}\} \subseteq \{x_1, \dots, x_n\}$ , where  $i_1 < i_2 < \dots < i_k$ , then we denote by  $x_A$  the monomial  $x_{i_1} \dots x_{i_k}$ .

Modulo  $I$  the polynomial  $f$  can be represented in the form

$$f = \sum_A x_A f_A,$$

where  $f_A$  is a commutator polynomial in the variables in  $\{x_1, \dots, x_n\} \setminus A$ . Let  $A$  be a set of maximal cardinality such that  $f_A \notin I$ . We may assume without loss of generality that  $A = \{x_1, \dots, x_k\}$ . Consider the polynomial  $g(x_{k+1}, \dots, x_n) = f(x_1, \dots, x_n)|_{x_1=x_2=\dots=x_k=1}$ . Since  $f \in I_1$ , we have  $g \in I_1$ . Obviously,  $g = f_A \pmod{I}$ . Therefore,  $f_A \in I$ . Then, by Lemma 5,  $I_1$  con-

tains the polynomial  $h = [z_1, t_1]^2 \dots [z_p, t_p]^2$  for some  $p$ , where  $z_i, y_i$  are pairwise distinct variables in  $X$ . It remains to show that  $h \notin I_1$ . Put  $z_i = e_{2i-1} + f_{2i-1}$ ,  $t_i = e_{2i} + f_{2i}$ . Then we easily see that  $[z_1, t_1]^2 \dots [z_p, t_p]^2 = 4e_1^p \dots e_{2p}^p f_1 \dots f_{2p}$ , i.e.,  $h \notin I_1$ .

The proposition is proved.

Proposition 3. Suppose  $\Gamma$  is a  $T$ -ideal such that  $\Gamma \supseteq \Gamma_2$ ,  $\Gamma \neq \Gamma_2$ . Then for some natural number  $q$  we have  $\Gamma \supseteq \Gamma_1^q$ .

Proof. Consider the  $\mathcal{S}$ -ideal  $\mathcal{S} = \Gamma + I$ . Since  $\Gamma_2$  is the largest  $T$ -ideal contained in  $I$  and since  $\Gamma \neq \Gamma_2$ , it follows that  $\mathcal{S} \neq I$ . Suppose  $f \in \mathcal{S}$ ,  $f \notin I$ . Since  $\mathcal{S}, I$  are  $\mathcal{S}$ -ideals, we may assume that  $f$  is a multilinear polynomial in the variables  $x_1, \dots, x_n$ . As in Proposition 2, we represent  $f$  in the form

$$f = \sum_A x_A f_A,$$

where  $f_A$  is a commutator polynomial in the variables in  $\{x_1, \dots, x_n\} \setminus A$ . Let  $A$  be a set of maximal cardinality such that  $f_A \notin I$ . We may assume without loss of generality that  $A = \{x_1, \dots, x_k\}$ . Put

$$g(y_1, \dots, y_k, z_1, \dots, z_k, x_{k+1}, \dots, x_n) = f(x_1, \dots, x_n) \Big|_{\substack{x_i = [y_i, z_i] \\ x_k = [y_k, z_k]}}, \quad (20)$$

where  $y_i, z_j, x_s$  are pairwise distinct variables in  $X$ . Obviously,

$$g \equiv [y_1, z_1] \dots [y_k, z_k] f_A(x_{k+1}, \dots, x_n) \pmod{I}. \quad (21)$$

We will show that  $g \in \mathcal{S}$ . Since  $f \in \mathcal{S}$ , we have  $f = f_1 + f_2$ , where  $f_1 \in \Gamma$ ,  $f_2 \in I$ . We define polynomials  $g_1, g_2$  by means of (20) (instead of  $f$  use  $f_1$  and  $f_2$ ). Then  $g = g_1 + g_2$ . Since  $\Gamma$  is  $T$ -ideal  $g_1 \in \Gamma$ . The polynomial  $f_2$  can be represented in the form  $f_2 = \sum_i u_i h_i v_i$ , where  $u_i, v_i \in F\langle X \rangle \cup \{1\}$ , and the  $h_i$  are polynomials of the form (13) or (14). If  $x_1, \dots, x_k$  do not occur in  $h_i$ , then

$$u_i h_i v_i \Big|_{\substack{x_i = [y_i, z_i] \\ x_k = [y_k, z_k]}} = \sum u' h_i v'. \quad (22)$$

In the opposite case, the left-hand side of (22) lies in the  $\mathcal{S}$ -ideal generated by the polynomials of the form (13). Therefore,  $g_2 \in I$ . Then it follows from (21) that  $[y_1, z_1] \dots [y_k, z_k] f_A \in I$ , where  $f_A$  is a commutator polynomial,  $f_A \notin I$ . Using (15), it is easy to see that

$$[y_1, z_1] \dots [y_k, z_k] E \subseteq \mathcal{S}, \quad (23)$$

where  $E$  is the  $\mathcal{S}$ -ideal generated by  $\{f_A\} \cup I$ . By Lemma 5,  $E \ni [y_{k+1}, z_{k+1}]^2 \dots [y_p, z_p]^2$  for some  $p$ . Therefore, in view of (23) and (15), we obtain  $[y_1, z_1]^2 \dots [y_p, z_p]^2 \in \mathcal{S}$ .

Put

$$h(x_1, x_2, t_1, t_2) = \sum_{\sigma, \tau \in \mathcal{S}(2)} [x_{\sigma(1)}, t_{\sigma(1)}] \cdot [x_{\sigma(2)}, t_{\sigma(2)}],$$

where  $x_i, t_j \in X$  ( $h$  is a linearization of the polynomial  $[y, x]^2$ ). Since  $\mathcal{S}$  is an  $\mathcal{S}$ -ideal, it follows from what was said above that

$$u(x_1, \dots, x_{2p}, t_1, \dots, t_{2p}) = h(x_1, x_2, t_1, t_2) \dots h(x_{2p-1}, x_{2p}, t_{2p-1}, t_{2p}) \in \mathcal{S},$$

where  $x_i, t_j$  are pairwise distinct variables in  $X$ .

Put  $\sigma(x_1, \dots, x_{4p}, y_1, \dots, y_{4p}) = u(c_1, \dots, c_{4p})$ , where  $c_i = [x_i, y_i]$  and  $\sigma$  is multilinear. We will show that  $\sigma \in \mathcal{I}$ . Indeed, since  $u \in \mathcal{S} = \mathcal{I} + \mathcal{I}$ , we have  $u = u_1 + u_2$ , where  $u_1 \in \mathcal{I}$  and  $u_2$  lies in the  $\mathcal{S}$ -ideal generated by the polynomials  $[c_i, c_j, c_k]$ ,

$$\sum_{\sigma \in \mathcal{S}(3)} [c_{\sigma(1)}, c_4] [c_{\sigma(2)}, c_5] [c_{\sigma(3)}, c_6],$$

where  $c_i = [x_i, y_i]$ . Therefore, it suffices to show that these polynomials lie in  $\mathcal{I}_2$ , or, equivalently, that the algebra  $G \otimes G$  (or  $G^* \otimes G^*$ ) satisfies the identities

$$[c_1, c_2, c_3] = 0, \quad (24)$$

$$\sum_{\sigma \in \mathcal{S}(3)} [c_{\sigma(1)}, c_4] [c_{\sigma(2)}, c_5] [c_{\sigma(3)}, c_6] = 0, \quad (25)$$

where  $c_i = [x_i, y_i]$ .

The algebra  $G^* \otimes G^*$  can be represented in the form

$$G^* \otimes G^* = Z + \sum_i e_i Z + \sum_i f_i Z + \sum_{i,j} e_i f_j Z,$$

where  $Z$  is the center of  $G^* \otimes G^*$ . Using these decompositions, we see easily that for any  $x_i, y_i \in G^* \otimes G^*$  we have

$$c_i = [x_i, y_i] \in Z + \sum_i e_i Z + \sum_i f_i Z.$$

After this observation it is trivial to verify (24) and (25).

It is well known and easy to prove that if  $g = f_1 \dots f_n \in \mathcal{I}$ , where  $\mathcal{I}$  is a  $\mathcal{T}$ -ideal and the polynomials  $g, f_1, \dots, f_n$  are multilinear, then  $\mathcal{I}_1 \dots \mathcal{I}_n \subseteq \mathcal{I}$ , where  $\mathcal{I}_i = \{f_i\}^{\mathcal{I}}$ . In view of this observation and what was proved above, it follows that  $\mathcal{I}^p \subseteq \mathcal{I}$ , where  $\mathcal{I}$  is the  $\mathcal{T}$ -ideal generated by the set  $\{h(c_1, c_2, c_3, c_4)\} \cup \mathcal{I}_2$ , where  $c_i = [x_i, y_i]$ . It remains to prove that  $\mathcal{I} \supseteq \mathcal{I}_1^z$  for some  $z$ . We isolate this fact in a separate lemma.

**LEMMA 6.**  $\mathcal{I} \supseteq \mathcal{I}_1^z$ .

**Proof.** Since identity (24) holds modulo  $\mathcal{I}_2$  and since  $\mathcal{I} \supseteq \mathcal{I}_2$ , we have modulo  $\mathcal{I}$  the identities

$$\frac{1}{2} h(c_1, c_2, c_3, c_4) \stackrel{(24)}{=} [c_1, c_2] [c_3, c_4] + [c_1, c_4] [c_3, c_2] \equiv 0, \quad (26)$$

$$[c_1, c_2] [c_3, c_4] = -[c_1, c_4] [c_3, c_2]. \quad (27)$$

It is easy to see that in the algebra  $G \otimes G$ , hence also modulo  $\mathcal{I}$ , we have the identity

$$[[x_1, x_2], [x_3, x_4], x_5] = 0. \quad (28)$$

In (26) we make the replacement  $x_7 = x_6 c_5$ , where  $c_5 = [x_5, y_5]$ . Using (27) and (28), we obtain the following congruence modulo  $\tau$ :

$$0 \equiv [x_6, y_7] ([c_5, c_2] [c_3, c_4] + [c_5, c_4] [c_3, c_2]) + c_5 ([x_6, y_7, c_2] [c_3, c_4] + [x_6, y_7, c_4] [c_3, c_2]) + x_6 ([c_5, y_7, c_2] [c_3, c_4] + [c_5, y_7, c_4] [c_3, c_2]) + ([x_6, c_2] [c_3, c_4] + [x_6, c_4] [c_3, c_2]) [c_5, y_7].$$

Then

$$([x_6, c_2] [c_3, c_4] + [x_6, c_4] [c_3, c_2]) [c_5, y_7] \equiv 0 \pmod{\tau}. \quad (29)$$

Making the substitution  $x_6 = x_6 c_7$ , where  $c_7 = [x_7, y_7]$ , we obtain the congruence

$$([x_6, c_2] c_7 [c_3, c_4] + [x_6, c_4] c_7 [c_3, c_2]) [c_5, y_7] \equiv 0 \pmod{\tau},$$

from which it follows that

$$([x_6, c_2, c_7] [c_3, c_4] + [x_6, c_4, c_7] [c_3, c_2]) [c_5, y_7] \equiv 0 \pmod{\tau}. \quad (30)$$

Using (27) and (28), we obtain

$$[x_6, c_2, c_7] [c_3, c_4] \stackrel{(27)}{=} -[x_6, c_2, c_4] [c_3, c_7] \stackrel{(28)}{=} -[x_6, c_4, c_2] [c_3, c_7] \stackrel{(27)}{=} [x_6, c_4, c_7] [c_3, c_2] \pmod{\tau}.$$

In view of (30),

$$[x_6, c_4, c_7] [c_3, c_2] [c_5, y_7] \equiv 0 \pmod{\tau}. \quad (31)$$

Let  $\mathcal{B}$  be the  $\tau$ -ideal generated by the polynomial  $[x_1, x_2, x_3, [x_4, x_5]]$ , where  $x_i \in X$ . It follows from (31) that  $\tau \supseteq \mathcal{B}^2 \cdot \tau$ . It remains to show that  $\mathcal{B} \supseteq \tau$ .

In the identity (modulo  $\mathcal{B}$ )  $[x_1, x_2, x_3, [x_4, x_5]] = 0$  we make the substitution  $x_4 = x_4 \cdot [x_6, x_7]$  and obtain the congruence

$$0 \equiv [x_1, x_2, x_3, [x_4 \cdot [x_6, x_7], x_5]] \equiv [x_1, x_2, x_3, x_4 \cdot [x_6, x_7, x_5]] \equiv [x_1, x_2, x_3, x_4] [x_6, x_7, x_5] \equiv 0 \pmod{\mathcal{B}}. \quad (32)$$

Then making the substitution  $x_3 = x_3 \cdot [x_8, x_9]$  in (32) we obtain

$$0 \equiv [x_1, x_2, x_3 \cdot [x_8, x_9], x_4] [x_6, x_7, x_5] \equiv [x_1, x_2, x_3] \cdot [x_8, x_9, x_4] \times \\ \times [x_6, x_7, x_5] \equiv [x_1, x_2, x_3] [x_8, x_9, x_4] [x_6, x_7, x_5] \pmod{\mathcal{B}}.$$

The lemma and the proposition are proved.

### 3. NONMATRIX VARIETIES

In this section,  $\mathcal{M}$  is a fixed nonmatrix variety and  $\Gamma$  is the ideal of identities of  $\mathcal{M}$ . Suppose  $p, q$  are arbitrary nonnegative integers. Put  $\bar{F}_{p,q} = F_{p,q} / \Gamma(F_{p,q})$ . The images of the sets  $E_i$ ,  $Y_q = \{y_1, \dots, y_q\}$  under the natural homomorphism  $F_{p,q} \rightarrow \bar{F}_{p,q}$  will be denoted by  $\bar{E}_i$ ,  $\bar{Y}_q = \{z_1, \dots, z_q\}$  respectively.

Let  $\mathcal{U}_{p,q}$  be the ideal of  $\bar{F}_{p,q}$  generated by all elements of the form  $[z_i, z_j], [e, f, z_i], [ef, gh], [e, f, g]$ , where

$$z_i, z_j \in \bar{Y}_q, e, f, g, h \in \bigcup_{i=1}^p \bar{E}_i.$$

LEMMA 7. If  $p, q$  are arbitrary, then  $\mathcal{U}_{p,q}$  is a nilpotent ideal of the algebra  $\bar{F}_{p,q}$ .

Proof. Let  $V_{i,j}^{(1)}$ , where  $1 \leq i, j \leq q$ , be the ideal generated by the element  $[z_i, z_j]$ ,  $z_i, z_j \in \bar{Y}_q$ ;  $V_{\kappa, \ell, i}^{(2)}$ , where  $1 \leq \kappa, \ell \leq p, 1 \leq i \leq q$ , the ideal generated by all elements  $[e, f, z_i]$ , where  $e \in \bar{E}_\kappa, f \in \bar{E}_\ell, z_i \in \bar{Y}_q$ ;  $V_{\kappa, \ell, \mu, \nu}^{(3)}$  is the ideal generated by the elements  $[ef, gh]$ , where  $e \in \bar{E}_\kappa, f \in \bar{E}_\ell, g \in \bar{E}_\mu, h \in \bar{E}_\nu, 1 \leq \kappa, \ell, \mu, \nu \leq p$ ;  $V_{\kappa, \ell, \mu}^{(4)}$ , where  $1 \leq \kappa, \ell, \mu \leq p$ , the ideal generated by all elements  $[e, f, g]$ , where  $e \in \bar{E}_\kappa, f \in \bar{E}_\ell, g \in \bar{E}_\mu$ . Since  $\mathcal{U}_{p,q}$  is the sum of the ideals  $V_{i,j}^{(1)}, V_{\kappa, \ell, i}^{(2)}, V_{\kappa, \ell, \mu, \nu}^{(3)}, V_{\kappa, \ell, \mu}^{(4)}$ , the number of which is finite, it suffices to prove the nilpotency of which of these four ideals for fixed  $i, j, \kappa, \ell, \mu, \nu$ .

Since  $\mathcal{M}$  is a nonmatrix variety, the commutator ideal of the algebra  $A = F\langle X \rangle / \Gamma$  is a nil ideal. Therefore,  $A$  satisfies the identity  $([x, t]y)^m = 0$  for some  $m$ . This means that for any  $x, t$  the right ideal  $E = [x, t]A$  satisfies the identity  $x^m = 0$ , where  $m = m(\Gamma)$ . By the Nagata-Higman theorem [10],  $E$  is nilpotent. But then the ideal generated by  $[x, t]$  is nilpotent. Therefore, the algebra  $A$ , hence any algebra in  $\mathcal{M}$ , satisfies the identity

$$[x, t]y, [x, t]y_2 \dots y_n [x, t] = 0 \quad (33)$$

for an arbitrary  $n = n(m)$ , where some of the variables  $y_i$  can be absent. It follows at once from (33) that  $V_{i,j}^{(1)}$  is nilpotent.

Linearizing (33), we obtain an equivalent identity

$$\sum_{\sigma \in S(n+1)} [x_{\sigma(1)}, t_{\sigma(1)}] y_1 \dots y_n [x_{\sigma(n+1)}, t_{\sigma(n+1)}] = 0. \quad (34)$$

We make the substitutions  $x_s = [e_s, f_s]$  in (34), where  $e_s \in \bar{E}_\kappa, f_s \in \bar{E}_\ell; t_s = z_i$ , where  $z_i \in \bar{Y}_q$ ;  $y_s = a_s$ , where  $a_s \in \bar{F}_{p,q} \cup \{1\}$ . In the algebra  $\bar{F}_{p,q}$  we obtain

$$\begin{aligned} 0 &= (n+1)! \sum_{\sigma \in S(n+1)} [e_{\sigma(1)}, f_{\sigma(1)}, z_i] a_1 \dots a_n [e_{\sigma(n+1)}, f_{\sigma(n+1)}, z_i] \stackrel{(1)}{=} \\ &= [(n+1)!]^2 \sum_{\sigma \in S(n+1)} (-1)^\sigma [e_1, f_{\sigma(1)}, z_i] a_1 \dots a_n [e_{n+1}, f_{\sigma(n+1)}, z_i] \stackrel{(1)}{=} \\ &= [(n+1)!]^3 [e_1, f_1, z_i] a_1 \dots a_n [e_{n+1}, f_{n+1}, z_i], \end{aligned}$$

from which it follows at once that  $V_{\kappa, \ell, i}^{(2)}$  is nilpotent.

In exactly the same way we can prove that  $V_{\kappa, \ell, \mu, \nu}^{(3)}$  is nilpotent. To do this we make the replacement  $x_s = e_s f_s, t_s = g_s h_s, y_s = a_s$  in (34), where

$$e_s \in \bar{E}_\kappa, f_s \in \bar{E}_\ell, g_s \in \bar{E}_\mu, h_s \in \bar{E}_\nu, a_s \in \bar{F}_{p,q} \cup \{1\}.$$

It remains to show that the ideal  $V_{\kappa, \ell, \mu}^{(4)}$  is nilpotent. Put

$$V^{(3)} = \sum_{1 \leq \kappa, \ell, \mu, \nu \leq p} V_{\kappa, \ell, \mu, \nu}^{(3)}.$$

Since the ideals  $V_{\kappa, \ell, \mu, \nu}^{(3)}$  are nilpotent, so is  $V^{(3)}$ .

Suppose  $\ell_s \in \bar{E}_\kappa, g_s \in \bar{E}_\mu, f_s \in \bar{E}_\nu$ . Modulo  $V^{(3)}$  we have the congruences

$$\begin{aligned} & [\ell_1 \ell_2, g_1] f [\ell_3 \ell_4, g_2] = \ell_1 \ell_2 g_1 f [\ell_3 \ell_4, g_2] - g_1 \ell_1 \ell_2 f [\ell_3 \ell_4, g_2] = \\ & = g_1 f \ell_1 \ell_2 [\ell_3 \ell_4, g_2] - \ell_2 f g_1 \ell_1 [\ell_3 \ell_4, g_2] = g_1 f \ell_1 [\ell_3 \ell_4, \ell_2 g_2] - \ell_2 f g_1 [\ell_3 \ell_4, \ell_1 g_2] \equiv 0, \\ \text{i.e., } & [\ell_1 \ell_2, g_1] f [\ell_3 \ell_4, g_2] \equiv 0 \pmod{V^{(3)}}. \end{aligned}$$

Therefore, since  $V^{(3)}$  is nilpotent,  $\Gamma(F_{p,q})$  (hence also  $\Gamma(F_{E,Y})$ ) contains, for some  $m$ , all elements of the form

$$[\ell_1 \ell_2, g_1] f_1 [\ell_3 \ell_4, g_2] f_2 \dots f_{m-1} [\ell_{2m-1} \ell_{2m}, g_m] f_m,$$

where  $\ell_s \in E_\kappa, g_s \in E_\mu, f_s \in E_\nu$ . It follows from this and Lemmas 1 and 2 that the algebra  $A = F\langle X \rangle / \Gamma$  (and also any algebra in  $\mathcal{M}$ ) satisfies the identity

$$\sum_{\substack{\sigma \in S(2m) \\ \tau, \pi \in S(m)}} (-1)^\sigma (-1)^\tau (-1)^\pi [x_{\sigma(1)} x_{\sigma(2)}, y_{\tau(1)}] t_{\pi(1)} \dots [x_{\sigma(2m-1)} x_{\sigma(2m)}, y_{\tau(m)}] t_{\pi(m)} = 0. \quad (35)$$

If in (35) we put  $x_\tau = \ell_\tau, y_\tau = g_\tau, t_\tau = a_\tau$ , where  $\ell_\tau \in \bar{E}_\kappa, g_\tau \in \bar{E}_\mu, a_\tau \in \bar{F}_{E,Y}$  ( $\bar{F}_{E,Y} = F_{E,Y} / \Gamma(F_{E,Y})$ ) we obtain in the algebra  $\bar{F}_{E,Y}$  the equality

$$0 = \sum_{\sigma \in S(m)} (-1)^\sigma [\ell_1 \ell_2, g_1] a_{\sigma(1)} \dots [\ell_{2m-1} \ell_{2m}, g_m] a_{\sigma(m)} \stackrel{(1)}{=} \sum_{\sigma \in S(m)} (-1)^\sigma h_\sigma(\ell_1, \dots, \ell_m) a_{\sigma(1)} \dots a_{\sigma(m)},$$

$$\stackrel{(1)}{=} h_m(\ell_1, \dots, \ell_m) = 0, \text{ where } h_m(x_1, \dots, x_m) = \sum_{\sigma \in S(m)} x_{\sigma(1)} \dots x_{\sigma(m)}; \ell_s = [\ell_{2s-1} \ell_{2s}, g_s] a_s.$$

Therefore, the right ideal  $\mathcal{L}_{\kappa, \mu}$  of  $\bar{F}_{E,Y}$  generated by all elements  $[\ell_1 \ell_2, g] a$ , where  $\ell_s \in \bar{E}_\kappa, g \in \bar{E}_\mu, a \in \bar{F}_{E,Y}$ , satisfies the identity  $h_m(x_1, \dots, x_m) = 0$ . By the Nagata-Higman theorem [10], we see that  $\mathcal{L}_{\kappa, \mu}$ , hence also the ideal  $W_{\kappa, \mu}$  generated by the elements  $[\ell_1 \ell_2, g]$ , is nilpotent, i.e., for some  $\nu = \nu(m)$  the ideal  $\Gamma(F_{E,Y})$  contains all elements of the form

$$[\ell_1 \ell_2, g_1] a_1 \dots a_{\nu-1} [\ell_{2\nu-1} \ell_{2\nu}, g_\nu],$$

where  $\ell_s \in E_\kappa, g_s \in E_\mu, a_s \in F_{E,Y} \cup \{1\}$ . Again applying Lemmas 1 and 2, we see that any algebra in  $\mathcal{M}$  satisfies the identity

$$\sum_{\substack{\sigma \in S(2\nu) \\ \tau \in S(\nu)}} (-1)^\sigma (-1)^\tau [x_{\sigma(1)} x_{\sigma(2)}, y_{\tau(1)}] t_1 \dots t_{\nu-1} [x_{\sigma(2\nu-1)} x_{\sigma(2\nu)}, y_{\tau(\nu)}] = 0, \quad (36)$$

where some of the variables  $t_s$  can be absent.

Put  $W'_{\kappa, \mu} = W_{\kappa, \mu} \cap F_{p,q}$ ,  $W = \sum_{\kappa, \mu \leq p} W'_{\kappa, \mu}$ . In view of what was said above,  $W$  is nilpotent.

If in (36) we make the substitution  $x_{2s-1} = \ell_s, x_{2s} = f_s, y_s = g_s, t_s = a_s$ , where  $\ell_s \in \bar{E}_\kappa, f_s \in \bar{E}_\ell, g_s \in \bar{E}_\mu, a_s \in \bar{F}_{p,q} \cup \{1\}$  we obtain modulo  $W$  a congruence in the algebra  $\bar{F}_{p,q}$ :



$$\begin{aligned}
0 &= \sum_{\substack{\sigma \in S(2t) \\ \tau \in S(t)}} (-1)^\sigma (-1)^\tau [x_{\sigma(1)} x_{\sigma(2)} y_{\tau(1)}] t_1 \dots t_{t-1} [x_{\sigma(2t-1)} x_{\sigma(2t)} y_{\tau(t)}] \begin{matrix} x_{2s-1} = e_s \\ x_{2s} = f_s \\ y_s = g_s \\ t_s = a_s \end{matrix} = \\
&= \sum_{\sigma, \tau, \pi \in S(t)} (-1)^\sigma (-1)^\tau (-1)^\pi [e_{\sigma(1)} f_{\tau(1)} g_{\pi(1)}] a_1 \dots a_{t-1} [e_{\sigma(t)} f_{\tau(t)} g_{\pi(t)}],
\end{aligned}$$

hence, in view of (1),

$$[e_1, f_1, g_1] a_1 [e_2, f_2, g_2] a_2 \dots a_{t-1} [e_t, f_t, g_t] \equiv 0 \pmod{W}.$$

This implies that the ideal  $V_{\kappa, \ell, \mu}^{(4)}$  is nilpotent.

The lemma is proved.

Proposition 4. For any  $\rho$  there exist  $\alpha = \alpha(\rho)$  such that  $T[\bar{F}_{\rho,0}] \supseteq \Gamma_2^\alpha$  ( $T[\bar{F}_{\rho,0}]$  is the ideal of identities of the algebra  $\bar{F}_{\rho,0}$ ).

Proof. In view of Lemma 7, it suffices to show that  $T[\bar{F}_{\rho,0}/\mathcal{U}_{\rho,0}] \supseteq \Gamma_2$ . In view of Proposition 2, it is enough to prove that in the algebra  $\bar{F}_{\rho,0}/\mathcal{U}_{\rho,0}$  the generators in  $\bigcup_{i=1}^P E_i$  satisfy (15) and (16).

In this case the ideal  $\mathcal{U}_{\rho,0}$  is generated by the elements  $[e, f, g]$ ,  $[e f, g h]$ , where  $e, f, g, h \in \bigcup_{i=1}^P E_i$ . Relation (15) follows trivially from the definition of  $\mathcal{U}_{\rho,0}$ . We will show that in the algebra  $\bar{F}_{\rho,0}$  the generators in  $\bigcup_{i=1}^P E_i$  satisfy modulo  $\mathcal{U}_{\rho,0}$  the relation

$$\sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] x_{\sigma(2)} x_{\sigma(3)} = 0. \quad (37)$$

Indeed,

$$\begin{aligned}
\sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] x_{\sigma(2)} x_{\sigma(3)} &\equiv \sum_{\sigma \in S(3)} x_{\sigma(1)} [x_{\sigma(2)}, y_1] x_{\sigma(3)} = \sum_{\sigma \in S(3)} x_{\sigma(1)} x_{\sigma(2)} y_1 x_{\sigma(3)} - \\
&- \sum_{\sigma \in S(3)} x_{\sigma(1)} y_1 x_{\sigma(2)} x_{\sigma(3)} \equiv \sum_{\sigma \in S(3)} y_1 x_{\sigma(1)} x_{\sigma(2)} x_{\sigma(3)} - \sum_{\sigma \in S(3)} x_{\sigma(1)} y_1 x_{\sigma(2)} x_{\sigma(3)} = \sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] x_{\sigma(2)} x_{\sigma(3)},
\end{aligned}$$

which implies (37). From (37) we obtain

$$0 \equiv \sum_{\sigma \in S(3)} [[x_{\sigma(1)}, y_1] x_{\sigma(2)} x_{\sigma(3)}] y_2 y_3 = 2 \sum_{\sigma \in S(3)} [[x_{\sigma(1)}, y_1] [x_{\sigma(2)}, y_2] x_{\sigma(3)}] y_3 \equiv 2 \sum_{\sigma \in S(3)} [x_{\sigma(1)}, y_1] [x_{\sigma(2)}, y_2] [x_{\sigma(3)}, y_3].$$

Relation (16) is proved.

The proposition is proved.

From now on,  $t, z$  are fixed natural numbers such that the  $T$ -ideal  $\Gamma$  satisfies the conclusion of Proposition 1, and  $\rho = t + tz$ .

LEMMA 8. Suppose  $f = f(x_1, \dots, x_n)$  is a multilinear polynomial and  $f \in T[\bar{F}_{\rho,0}]$ . Let  $T_f$  denote the  $T$ -ideal generated by the multilinear polynomial

$$g = f(x_1, \dots, x_n) s_{2(z+n)}(y_1, \dots, y_{2(z+n)}),$$

where  $x_i, y_j \in X$ , and  $S_K$  is a standard polynomial, and let  $T_2$  denote the  $T$ -ideal generated by all multilinear polynomials

$$h_W = f(w_1 S_{2(r+1)}(y_1^{(1)}, \dots, y_{2(r+1)}^{(1)}), \dots, w_n S_{2(r+1)}(y_1^{(n)}, \dots, y_{2(r+1)}^{(n)})),$$

where  $w_i, y_j^{(i)} \in X$ , some of the variables  $w_i$  can be absent, and  $W$  is the set of variables  $w_i$  occurring in  $h_W$ . Then  $T_1^\rho, T_2^\rho \subseteq T$  for some natural number  $\rho$ .

Proof. By Proposition 1,  $T = T[A, Z]$ , where  $A = \bar{F}_{t,r}$ , and  $Z$  is the subspace of  $A$  generated by the set  $\bigcup_{i=1}^t \bar{E}_i \cup \bar{Y}_r$ . By Lemma 7, the ideal  $\mathcal{U}_{t,r}$  is nilpotent. Let  $\rho$  be the index of nilpotency of  $\mathcal{U}_{t,r}$ . In view of what was said above, it suffices to show that  $g, h_W \in \mathcal{U}_{t,r}$  for any  $x_i, y_i, y_j^{(i)}$ ,  $w_i \in \bigcup_{i=1}^t \bar{E}_i \cup \bar{Y}_r$ .

Consider the subalgebra  $\mathcal{D}$  of  $\bar{F}_{t,r}$  generated by the set  $\bigcup_{i=1}^t \bar{E}_i \cup (\bigcup_{j=1}^r \bigcup_{i=1}^t x_j \bar{E}_i)$ , where  $x_j \in \bar{Y}_r$ . Since the elements of  $x_j \bar{E}_i$  satisfy (1), the algebra  $\mathcal{D}$  is a homomorphic image of  $\bar{F}_{p,0}$ , where  $p = t + tr$ .

If more than  $r$  elements among  $y_1, \dots, y_{2(r+k)}$  belong to  $\bar{Y}_r$ , then at least two of these elements are equal to and we have  $g = 0$ . Similarly, if more than  $r$  elements among  $y_1^{(i)}, \dots, y_{2(r+1)}^{(i)}$  belong to  $\bar{Y}_r$  for some  $i$ , then  $h_W = 0$ . In the opposite case, since

$$S_{2K}(x_1, \dots, x_{2K}) = \frac{1}{2^K} \sum_{\sigma \in S(2K)} (-1)^\sigma [x_{\sigma(1)}, x_{\sigma(2)}] \dots [x_{\sigma(2K-1)}, x_{\sigma(2K)}],$$

it follows that  $S_{2(r+n)}(y_1, \dots, y_{2(r+n)})$  can be represented modulo  $\mathcal{U}_{t,r}$  as a linear combination of elements of the form  $c_1 \dots c_n d$ , where  $c_i = [a_i, b_i]$ ,

$$a_i, b_i \in \bigcup_{i=1}^t \bar{E}_i, d \in \mathcal{D},$$

and  $S_{2(r+1)}(y_1^{(i)}, \dots, y_{2(r+1)}^{(i)})$  can be represented as a linear combination of elements of the form  $cd$ , where  $c = [a, b]$ ,  $a, b \in \bigcup_{i=1}^t \bar{E}_i$ ,  $d \in \mathcal{D}$ . Therefore, since modulo  $\mathcal{U}_{t,r}$  the elements  $c_i, c$  lie in the center, it follows that  $g$  and  $h_W$  can be represented modulo  $\mathcal{U}_{t,r}$  as a linear combination of elements of the form  $f(d_1, \dots, d_n) d_{n+1}$ , where  $d_i \in \mathcal{D}$ . Since  $f \in T[\bar{F}_{p,0}] \subseteq T[\mathcal{D}]$ , we have  $f(d_1, \dots, d_n) = 0$ . This means that  $g, h_W \in \mathcal{U}_{t,r}$ .

The lemma is proved.

To prove the main results we will need

LEMMA 9. If a nonmatrix variety  $\mathcal{L}$  does not contain the Grassmann algebra  $G$ , then  $\mathcal{L}$  satisfies the identity  $[x_1, x_2] \dots [x_{2n-1}, x_{2n}] = 0$  for some  $n$ .

Proof. Since  $G \notin \mathcal{L}$ , it follows that  $\mathcal{L}$  satisfies the Capelli identities of some order  $K$ , as shown in [11]. Then, as is well known [13],  $\mathcal{L}$  is generated by a  $(k-1)$ -generator algebra. Since  $\mathcal{L}$  is a nonmatrix variety, the commutator ideal of any finitely generated algebra in  $\mathcal{L}$  is nilpotent.

The lemma is proved.

We can now turn to the proof of the main results.

THEOREM 1. Suppose  $\mathcal{M}$  is a nonmatrix variety  $\mathcal{M} \not\supset G \otimes G$ , and  $\mathcal{A}$  is the largest variety in  $\{\mathcal{O}, \mathcal{A}_0, \mathcal{A}_1\}$  contained in  $\mathcal{M}$ . Then for some  $\kappa$  we have  $\mathcal{M} = \mathcal{M}_\kappa \circ_m \mathcal{A}$ , where  $\mathcal{M}_\kappa = \mathcal{M} \cap \mathcal{N}_\kappa$ .

Proof. If  $\mathcal{A} = \mathcal{O}$ , the conclusion of the theorem follows from the Nagata-Higman theorem, and if  $\mathcal{A} = \mathcal{A}_0$ , from Lemma 9. Suppose  $\mathcal{A} = \mathcal{A}_1$ ,  $t, \nu$  are natural numbers such that the ideal of identities  $\Gamma$  of the variety  $\mathcal{M}$  satisfies the conclusion of Proposition 1, and  $\rho = t + t\nu$ .

By Proposition 4, the algebra  $\bar{F}_{\rho,0}$  contains a nilpotent ideal  $\mathcal{U}$  such that  $T[\bar{F}_{\rho,0}/\mathcal{U}] \supseteq \Gamma_2 + \Gamma$ . Since  $\mathcal{M} \not\supset G \otimes G$ , we have  $\Gamma + \Gamma_2 \neq \Gamma_2$ . Applying Proposition 3 to the  $\Gamma$ -ideal  $T[\bar{F}_{\rho,0}/\mathcal{U}]$ , we obtain  $T[\bar{F}_{\rho,0}] \supseteq \Gamma_1^\delta$  for some  $\delta$ . Since  $\Gamma_1 = \{[x, y, z]\}^\Gamma$ , we obtain, applying Lemma 8 to the polynomial  $f = [x, y, z] \cdot \dots \cdot [x, y, z]$ , that  $(\Gamma_1^\delta \Gamma)^\beta \subseteq \Gamma$  for some  $\beta$ , where  $\Gamma$  is the  $\Gamma$ -ideal generated by  $\{S_{2(\nu+3y)}(y, \dots, y_{2(\nu+3y)})\} \cup \Gamma$ . Since the identity  $S_t(x, \dots, x_t) = 0$  is not satisfied on  $G$ , it follows from Lemma 9 that  $T \supseteq \Gamma_0^n$  for some  $n$ , where  $\Gamma_0 = \{[x, y]\}^\Gamma$ . In view of what was said above,  $\Gamma \supseteq (\Gamma_1^\delta \Gamma)^\beta \supseteq (\Gamma_1^\delta \Gamma_0^n)^\beta \supseteq \Gamma_1^{\beta(\delta+n)}$ .

The theorem is proved.

THEOREM 2. Suppose  $\mathcal{M}$  is a nonmatrix variety and  $\mathcal{M} \ni G \otimes G$ . Then  $\mathcal{M} = \mathcal{M}_\kappa \circ_m (\mathcal{A}_2 \circ_m (\mathcal{M}_\ell \circ_m \mathcal{A}_0))$  for certain  $\kappa, \ell$ .

Proof. Suppose  $t, \nu$  are natural numbers such that the ideal of identities  $\Gamma$  of the variety  $\mathcal{M}$  satisfies the conclusion of Proposition 1, and let  $\rho = t + t\nu$ . Put  $M = F\langle X \rangle / \Gamma$ . By Lemma 8, the algebra  $M$  contains ideals  $\mathcal{I}_0, \mathcal{I}_2, \mathcal{I}_0 \subseteq \mathcal{I}_2$ , such that  $\mathcal{I}_0$  is nilpotent,  $T[\mathcal{I}_2/\mathcal{I}_0] \supseteq T[\bar{F}_{\rho,0}]$ , and algebra  $M/\mathcal{I}_2$  satisfies the standard identity of degree  $\nu+1$ . By Proposition 4,  $M$  contains an ideal  $\mathcal{J}_1, \mathcal{I}_0 \subseteq \mathcal{J}_1 \subseteq \mathcal{I}_2$ , such that  $\mathcal{J}_1$  is nilpotent and  $T[\mathcal{I}_2/\mathcal{J}_1] \supseteq \Gamma_2$ . By Lemma 8,  $M$  contains an ideal  $\mathcal{J}_3, \mathcal{I}_2 \subseteq \mathcal{J}_3$ , such that the algebra  $\mathcal{J}_3/\mathcal{I}_2$  is nilpotent and  $M/\mathcal{J}_3$  is commutative.

The theorem is proved.

THEOREM 3. Suppose  $\mathcal{M}$  is a nonmatrix variety,  $\mathcal{A}$  is a finitely based variety, and  $\mathcal{A} \ni \mathcal{A}_2$ . Then  $\mathcal{M} = \mathcal{M}_\kappa \circ_m (\mathcal{A} \cap \mathcal{M})$  for some  $\kappa$ .

Proof. This theorem is equivalent to the following assertion: If  $V$  is a finitely based  $\Gamma$ -ideal and  $V \subseteq \Gamma_2$ , then  $V$  is nilpotent modulo  $\Gamma = T[\mathcal{M}]$ . We may assume that  $V$  is generated by a single multilinear polynomial  $f_0(x_1, \dots, x_m)$ .

Suppose  $t, \nu$  are natural numbers such that the  $\Gamma$ -ideal  $\Gamma$  satisfies the conclusion of Proposition 1, and let  $\rho = t + t\nu$ . By Proposition 4,  $T[\bar{F}_{\rho,0}] \supseteq V^\alpha$  for some  $\alpha$ . Applying Lemma 8 to the polynomial  $f = f_0(x_1^{(\alpha)}, \dots, x_m^{(\alpha)}) \cdot \dots \cdot f_0(x_1^{(\alpha)}, \dots, x_m^{(\alpha)})$ , we obtain  $(V^\alpha \Gamma)^\beta \subseteq \Gamma$  for some  $\beta$ , where  $\Gamma = \{S_{2(\nu+\alpha m)}(x_1, \dots, x_{2(\nu+\alpha m)})\}^\Gamma + \Gamma$ . By Lemma 9,  $T \supseteq \Gamma_0^n$  for some  $n$ , where  $\Gamma_0 = \{[x, y]\}^\Gamma$ . In view of what was said above,  $\Gamma \supseteq (V^\alpha \Gamma)^\beta \supseteq (V^\alpha \Gamma_0^n)^\beta \supseteq V^{\beta(\alpha+n)}$ .

The theorem is proved.

We turn to the corollaries.

Corollary 1 follows from Theorem 3, since the variety of algebras satisfying the identity  $[[x, y], [x, t], h] = 0$  contains the variety  $\mathcal{A}_2$ .

Corollary 2 follows from Theorem 1, since  $T[\mathcal{A}_1] = \{[x, y, z]\}^T$ .

Corollary 3 follows from Corollary 2 and a theorem of Latyshev [2].

Corollary 4 follows from Corollary 3, since an identity of the form

$$\sum_i \alpha_i x^i y x^{n-i} = 0$$

is not satisfied by the algebras  $M_2$  and  $G \otimes G$ .

Indeed, since  $T[G \otimes G] = T[G^* \otimes G^*]$ , where  $G^*$  is a Grassmann algebra with unity, it suffices to show that this identity is not satisfied by the algebras  $M_2$  and  $G^* \otimes G^*$ . Suppose the algebra  $M_2$  (or  $G^* \otimes G^*$ ) satisfies the identity  $\sum_i \alpha_i x^i y x^{n-i} = 0$ . Since our algebra has a unity, this identity implies the identity  $[y, x, \dots, x] = 0$  (we need only make the substitution  $x \rightarrow x+1$  and take the homogeneous component of smallest degree). Now to see that the identity  $[y, x, \dots, x] = 0$  is not satisfied by  $M_2$  and  $G^* \otimes G^*$ , it suffices to make the following substitutions: in the first case,  $x = e_{11}$ ,  $y = e_{12} + e_{21}$  and in the second  $x = e_1 \otimes 1$ ,  $y = \sum_{i=2}^n e_i \otimes f_i$ .

Proof of Corollary 5. Put

$$W_m = \{[x_1, \dots, x_m]\}^T, \quad \mathcal{U}_{q,n} = \{[x_1, \dots, x_q, y, \dots, y]\}_n^T.$$

It suffices to show that for any  $q, n \geq 1$  there exists  $\rho = \rho(q, n)$  such that  $\mathcal{U}_{q,n} \supseteq W_{\rho(q,n)}$ . The proof is by induction on  $n$ . For  $n=1$  the assertion is trivial. Fix  $n$ . Suppose for any  $q$  there exists  $\rho(q, n)$  such that  $\mathcal{U}_{q,n} \supseteq W_{\rho(q,n)}$  (induction assumption). We fix  $q$  and will prove that there exists  $s = \rho(q, n+1)$  such that

$$\mathcal{U}_{q,n+1} \supseteq W_s.$$

It is easy to see that the identity  $[x_1, \dots, x_q, y, \dots, y] = 0$  is not satisfied by  $G \otimes G$ . Therefore, it follows from Theorem 1 that  $\mathcal{U}_{q,n+1} \supseteq \Gamma_1^{\alpha}$ . Thus, it suffices to show that for any  $\beta \geq 1$  there exists  $t = t(\beta)$  such that

$$\mathcal{U}_{q,n+1} + \Gamma_1^{\beta} \supseteq W_{t(\beta)}. \quad (38)$$

The proof is by induction on  $\beta$ . For  $\beta=1$  the assertion is trivial. Fix  $\beta$ . Suppose (38) holds (induction assumption). We will show that there exists  $\tau = t(\beta+1)$  such that

$$\mathcal{U}_{q,n+1} + \Gamma_1^{\beta+1} \supseteq W_{\tau}.$$

It follows from (38) that

$$\mathcal{U}_{q,n+1} + \Gamma_1^{\beta+1} \supseteq (\mathcal{U}_{q,n+1} + \Gamma_1^{\beta}) \Gamma_1 \supseteq W_{t(\beta)} \Gamma_1.$$

Analogously,

$$\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1} \supseteq \Gamma_1 W_{t(\beta)}.$$

Therefore,

$$\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1} \supseteq W_{t(\beta)} \Gamma_1 + \Gamma_1 W_{t(\beta)} + \mathcal{U}_{g,n+1}. \quad (39)$$

Put  $j = \max(g, t(\beta))$ . Then, modulo  $\mathcal{U}_{g,n+1}$  and  $\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$ , we have the identity

$$[x_1, \dots, x_j, \underbrace{y, \dots, y}_{n+1}] = 0. \quad (40)$$

Making a partial linearization in (40), we obtain

$$0 = \sum_{i=1}^{n+1} [x_1, \dots, x_j, \underbrace{y, \dots, y}_{i-1}, z, \underbrace{y, \dots, y}_{n-i+1}] = n [x_1, \dots, x_j, \underbrace{y, \dots, y}_n, z] + \sum_{i=1}^n \alpha_i [x_1, \dots, x_j, \underbrace{z, y, \dots, y}_i, \underbrace{y, \dots, y}_{n-i}] \quad (41)$$

for certain  $\alpha_i \in F$ .

Making the substitution  $z = [u, v]$  in (41), by virtue of (39) we obtain modulo  $\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$  the identity

$$[x_1, \dots, x_j, \underbrace{y, \dots, y}_n, [u, v]] = 0. \quad (42)$$

From (42) modulo  $\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$  we obtain the identity

$$[x_1, \dots, x_j, \underbrace{y, \dots, y}_n, t, [u, v]] = 0. \quad (43)$$

It follows from (42) and (43) that  $[\mathcal{U}_{g,n}, [u, v]] \subseteq \mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$ . By the induction assumption,  $\mathcal{U}_{g,n} \supseteq W_{p(g,n)}$ . Therefore, modulo  $\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$  we have the identity

$$[x_1, \dots, x_\mu, [u, v]] = 0, \quad \mu = p(g, n), \quad \mu \geq j. \quad (44)$$

From (40) modulo  $\mathcal{U}_{g,n+1} + \Gamma_1^{\beta+1}$  we obtain the identity

$$\sum_{\sigma \in S(n+1)} [x_1, \dots, x_\mu, y_{\sigma(1)}, \dots, y_{\sigma(n+1)}] = 0. \quad (45)$$

It follows from (44) and (45) that

$$0 = \sum_{\sigma \in S(n+1)} [x_1, \dots, x_\mu, y_{\sigma(1)}, \dots, y_{\sigma(n+1)}] = (n+1)! [x_1, \dots, x_\mu, y_1, \dots, y_{n+1}]$$

as required.

Corollary 6 follows from Corollary 3. Actually, it is known [12] that since  $\mathcal{M}$  is locally weakly Noetherian, the algebras in  $\mathcal{M}$  satisfy an identity of the form

$$[x, y, \dots, y] x^k [u, t, \dots, t] = 0, \quad (46)$$

but, as is easily seen, (46) is not satisfied by the algebras  $M_2$  and  $G \otimes G$ .

Corollary 7 follows immediately from (46) and Corollary 5.

The author would like to thank A. I. Shirshov and L. A. Bokut' for their interest and I. V. L'vov for some useful discussions.

#### LITERATURE CITED

1. A. I. Mal'tsev, Algebraic Systems [in Russian], Nauka, Moscow (1970).
2. V. N. Latyshev, "The finite basis property for the identities of certain rings," Usp. Mat. Nauk, 32, No. 4, 259-266 (1977).
3. P. J. Higgins, "Lie rings satisfying the Engel condition," Proc. Cambridge Phil. Soc., 50, No. 1, 8-15 (1954).
4. V. N. Latyshev, "Nonmatrix varieties of associative algebras," Doctoral Dissertation, Moscow State Univ., Moscow (1978).
5. G. de B. Robinson, Representation Theory of the Symmetric Group, Univ. of Toronto Press (1961).
6. C. Curtis and I. Reiner, Representation Theory of Finite Groups and Associative Algebras, Interscience, New York (1962).
7. H. Boerner, Representations of Groups, North-Holland, Amsterdam (1963).
8. A. R. Kemer, "The Specht property of T-ideals with power growth of codimensions," Sib. Mat. Zh., 19, No. 1, 54-69 (1978).
9. A. Regev, "Existence of identities in  $A \otimes B$ ," Israel J. Math., 11, No. 2, 131-152 (1972).
10. G. Higman, "On a conjecture of Nagata," Proc. Cambridge Phil. Soc., 52, No. 1, 1-4 (1956).
11. A. R. Kemer, "Remark on the standard identity," Mat. Zametki, 23, No. 5, 753-757 (1978).
12. A. Z. Anan'in, "Locally finitely approximable and locally representable varieties of algebras," Algebra Logika, 16, No. 1, 3-23 (1977).
13. Yu. P. Razmyslov, "The Jacobson radical in PI-algebras," Algebra Logika, 13, No. 3, 337-360 (1974).