

## LETTER TO THE EDITOR

### Serum MicroRNA Screening for *DICER1*-Associated Pleuropulmonary Blastoma

To the Editor: We read with interest the article ‘Judicious *DICER1* Testing and Surveillance Imaging Facilitates Early Diagnosis and Cure of Pleuropulmonary Blastoma’ [1], as it raised pertinent issues for the management of families known to carry *DICER1* mutations. The authors suggest that to detect early-stage (i.e., Type I) pleuropulmonary blastoma (PPB), for which survival rates are > 90% [2], children known to harbor a germline *DICER1* mutation should receive CT chest scan at 3 months of age, and again at 1–2 years if the first scan is negative [1]. Although the majority of patients with PPB are found to have germline *DICER1* mutations, penetrance is low. The majority of mutation carriers are unaffected [3], with only 10–20% estimated to develop PPB. Consequently, any screening programmes for PPB in patients with germline *DICER1* mutations needs to be as non-invasive as possible, minimizing exposure to ionizing radiation. Serum microRNA profiling may be an important addition to any programme of radiological surveillance. Serum microRNAs show considerable promise as cancer biomarkers [4], particularly as they are highly stable and resistant to degradation [5]. We recently identified a panel of microRNAs that were more abundant in the serum of a 2-year-old female at the time-of-diagnosis of an advanced (Type III) PPB, compared with patients with other solid tumors of childhood and a non-malignant control group [6]. The patient carried a germline *DICER1* mutation, and the PPB cells showed a further somatic ‘hotspot’ mutation in the *DICER1* RNaseIIIb domain, consistent with other reports [7]. Amongst the over-expressed serum microRNAs, there was significant over-representation of -3p strands, in keeping with the observation that *DICER1* RNaseIIIb hotspot mutations result in a -3p strand bias in affected tissues [8]. Two specific microRNAs from this panel (miR-125a-3p/miR-125b-2-3p), had highly elevated serum levels at PPB diagnosis and demonstrated early treatment-related reductions [6]. Importantly, in healthy family members with germline *DICER1* mutations, serum levels of these two microRNAs were similar to the control group, suggesting that the changes in the patient were directly attributable to release of microRNAs from the PPB tumor cells into the bloodstream and not from the germline *DICER1* mutation per se. Comprehensive evaluation of the clinical utility of serum microRNAs is now warranted in two patient groups. First, as a longitudinal screening-tool in patients with germline *DICER1* mutations, initially in parallel with judicious radiological imaging, to identify whether levels of PPB-specific serum microRNAs [6] are elevated in early-stage disease, where outcomes are more favorable [2]. Second, in patients presenting de novo with a lung lesion, in order to resolve diagnostic dilemmas, for example, distinguishing PPB from developmental anomalies such as congenital cystic adenomatous malformation (CCAM) [9]. As CCAMs are not associated with germline and somatic *DICER1* mutations, we hypothesize that the serum profiles obtained would not show the PPB-associated -3p strand bias. In summary, if the utility of longitudinal serum microRNA monitoring is confirmed in a larger cohort of patients with

germline *DICER1* mutations, the resultant decrease in CT scans will reduce the associated radiation-risk to babies and very young children [10].

Shivani Bailey, MD  
Department of Paediatric  
Haematology and Oncology  
Addenbrooke’s Hospital  
Hills Road, Cambridge, UK

Department of Pathology  
University of Cambridge  
Tennis Court Road  
Cambridge, UK

Katie L. Raby, MSc  
Department of Pathology  
University of Cambridge  
Tennis Court Road  
Cambridge, UK

Nicholas Coleman, MD, PhD\*  
Department of Pathology  
University of Cambridge  
Tennis Court Road  
Cambridge, UK

Department of Histopathology  
Addenbrooke’s Hospital  
Hills Road, Cambridge, UK

Matthew J. Murray, MD, PhD\*  
Department of Paediatric  
Haematology and Oncology  
Addenbrooke’s Hospital  
Hills Road, Cambridge, UK

Department of Pathology  
University of Cambridge  
Tennis Court Road  
Cambridge, UK

Conflict of interest: None

\*Correspondence to: Dr. Matthew J. Murray and Professor Nicholas Coleman, Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge CB2 1QP, UK. Email: mjm16@cam.ac.uk (Murray), nc109@cam.ac.uk (Coleman)

Received 16 June 2014; Accepted 9 July 2014

## REFERENCES

- Schultz KA, Harris A, Williams GM, et al. Judicious DICER1 testing and surveillance imaging facilitates early diagnosis and cure of pleuropulmonary blastoma. *Pediatr Blood Cancer* 2014; In Press.
- Bisogno G, Brennan B, Orbach D, et al. Treatment and prognostic factors in pleuropulmonary blastoma: An EXPeRT report. *Eur J Cancer* 2014;50:178–184.
- Slade I, Bacchelli C, Davies H, et al. DICER1 syndrome: Clarifying the diagnosis, clinical features and management implications of a pleiotropic tumour predisposition syndrome. *J Med Genet* 2011;48:273–278.
- Murray MJ, Coleman N. Testicular cancer: A new generation of biomarkers for malignant germ cell tumours. *Nat Rev Urol* 2012;9:298–300.
- Mitchell PS, Parkin RK, Kroh EM, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A* 2008;105:10513–10518.
- Murray MJ, Bailey S, Raby KL, et al. Serum levels of mature microRNAs in DICER1-mutated pleuropulmonary blastoma. *Oncogenesis* 2014;3:e87.
- Heravi-Moussavi A, Anglesio MS, Cheng SW, et al. Recurrent somatic *DICER1* mutations in nonepithelial ovarian cancers. *N Engl J Med* 2012;366:234–242.
- Anglesio MS, Wang Y, Yang W, et al. Cancer-associated somatic DICER1 hotspot mutations cause defective miRNA processing and reverse-strand expression bias to predominantly mature 3p strands through loss of 5p strand cleavage. *J Pathol* 2013;229:400–409.
- Hill DA, Dehner LP. A cautionary note about congenital cystic adenomatoid malformation (CCAM) type 4. *Am J Surg Pathol* 2004;28:554–555 author reply 555.
- Ahmed BA, Connolly BL, Shroff P, et al. Cumulative effective doses from radiologic procedures for pediatric oncology patients. *Pediatrics* 2010;126:e851–858.