

11. Bossuyt PM, Reitsma JB, Bruns DE et al: The STARD statement for reporting studies of diagnostic accuracy: explanation and elaboration. *Ann Intern Med* 2003; **138**: W1.
12. Scales CD Jr, Dahm P, Sultan S et al: How to use an article about a diagnostic test. *J Urol* 2008; **180**: 469.
13. Coplen DE, Austin PF, Yan Y et al: Correlation of prenatal and postnatal ultrasound findings with the incidence of vesicoureteral reflux in children with fetal renal pelvic dilatation. *J Urol* 2008; **180**: 1631.
14. Weinstein S, Obuchowski NA and Lieber ML: Clinical evaluation of diagnostic tests. *AJR Am J Roentgenol* 2005; **184**: 14.
15. Woodward M and Frank D: Postnatal management of antenatal hydronephrosis. *BJU Int* 2002; **89**: 149.
16. Quirino IG, Diniz JS, Bouzada MC et al: Clinical course of 822 children with prenatally detected nephrouropathies. *Clin J Am Soc Nephrol* 2012; **7**: 444.
17. Ismaili K, Avni FE, Piepsz A et al: Current management of infants with fetal renal pelvis dilatation: a survey by French-speaking pediatric nephrologists and urologists. *Pediatr Nephrol* 2004; **19**: 966.
18. Peters CA: Congenital obstructive nephropathy: is the fog lifting? *Kidney Int* 2005; **67**: 371.
19. Grignon A, Filion R, Filiatrault D et al: Urinary tract dilatation in utero: classification and clinical applications. *Radiology* 1986; **160**: 645.
20. Coplen DE, Austin PF, Yan Y et al: The magnitude of fetal renal pelvic dilatation can identify obstructive postnatal hydronephrosis, and direct postnatal evaluation and management. *J Urol* 2006; **176**: 724.
21. de Kort EH, Bambang Oetomo S and Zegers SH: The long-term outcome of antenatal hydronephrosis up to 15 millimetres justifies a noninvasive postnatal follow-up. *Acta Paediatr* 2008; **97**: 708.
22. Chipde SS, Lal H, Gambhir S et al: Factors predicting improvement of renal function after pyeloplasty in pediatric patients: a prospective study. *J Urol* 2012; **188**: 262.
23. Romao RL, Farhat WA, Pippi Salle JL et al: Early postoperative ultrasound after open pyeloplasty in children with prenatal hydronephrosis helps identify low risk of recurrent obstruction. *J Urol* 2012; **188**: 2347.
24. Fischer JE, Bachmann LM and Jaeschke R: A readers' guide to the interpretation of diagnostic test properties: clinical example of sepsis. *Intensive Care Med* 2003; **29**: 1043.
25. Mesrobian HG: Editorial comment. *J Urol* 2006; **176**: 727.
26. Peters CA: Editorial: The long-term followup of prenatally detected severe bilateral newborn hydronephrosis initially managed nonoperatively. *J Urol* 2002; **168**: 1121.
27. Dhillon HK: Prenatally diagnosed hydronephrosis: the Great Ormond Street experience. *Br J Urol* 1998; **81**: 39.
28. Hashim H and Woodhouse CJ: Ureteropelvic junction obstruction. *Eur Urol Suppl* 2012; **11**: 8.
29. Ransley PG, Dhillon HK, Gordon I et al: The postnatal management of hydronephrosis diagnosed by prenatal ultrasound. *J Urol* 1990; **144**: 584.
30. Bandin F, Siwy J, Breuil B et al: Urinary proteome analysis at 5-year followup of patients with non-operated ureteropelvic junction obstruction suggests ongoing kidney remodeling. *J Urol* 2012; **187**: 1006.

EDITORIAL COMMENT

The authors have demonstrated a positive correlation between prenatal and postnatal renal pelvic dilatation, Society for Fetal Urology grading and the need for surgical intervention for UPJO. Noteworthy is the diligence required to successfully identify prenatally and evaluate postnatally 312 of 371 patients with hydronephrosis. Paramount to the success of such a program is the multidisciplinary approach which transcends departmental and at some

institutions physical boundaries that separate adult and pediatric health care centers. This approach should serve as a model for other programs pursuing research endeavors in the field of fetal medicine.

C. D. Anthony Herndon

*Department of Urology
University of Virginia School of Medicine
Charlottesville, Virginia*