

The coenzyme Q system in the classification of the  
ascosporogenous yeast genus *Dekkera* and the asporogenous  
yeast genus *Brettanomyces*

Y. YAMADA, HISANO TAKINAMI-NAKAMURA, Y. TAHARA  
AND MAUDY TH. SMITH<sup>1</sup>

*Laboratory of Applied Microbiology, Department of Agricultural Chemistry,  
Shizuoka University, Shizuoka 422, Japan*

<sup>1</sup> *Yeast Division, Centraalbureau voor Schimmelcultures, Laboratory of  
Microbiology, Delft University of Technology, Delft, The Netherlands*

YAMADA, Y., TAKINAMI-NAKAMURA, H., TAHARA, Y. and SMITH, M. TH. 1980.  
The coenzyme Q system in the classification of the ascosporogenous yeast genus  
*Dekkera* and the asporogenous yeast genus *Brettanomyces*. *Antonie van  
Leeuwenhoek* 46: 595-599.

Fourteen strains of the genera *Dekkera* and *Brettanomyces* were examined for  
the coenzyme Q system. Without exception they contained the Q-9 system. The  
results are discussed from the taxonomic point of view.

#### INTRODUCTION

With respect to their physiology, representatives of the ascosporogenous yeast  
genus *Dekkera* and of its asporogenous counterpart *Brettanomyces*, are charac-  
terized by producing large amounts of acetic acid through an incomplete oxy-  
biontic dissimilation of carbohydrate. Furthermore, both genera show a so-  
called "negative Pasteur effect" ("Custers effect"), i.e. a stimulation of fermenta-  
tion by molecular oxygen (Wikén et al., 1961; Scheffers and Wikén, 1969).

These properties have been used as taxonomic criteria (van der Walt,  
1970*a, b*). In previous papers we stated that the coenzyme Q system is very useful  
for classifying yeasts and yeast-like organisms (Yamada and Kondo, 1972*a, b, c*,  
1973; Yamada et al., 1973*a, b, c*, 1976*a, b*, 1977).

This paper deals with the significance of the coenzyme Q system<sup>a</sup> in the  
taxonomy of organisms which belong to the genera *Dekkera* and *Brettanomyces*.

<sup>a</sup>The abbreviations used for coenzyme Q or ubiquinone in this paper are: Co-Q, coenzyme Q; Q-*n*,  
coenzyme Q with *n* isoprene units in the side chain, e.g. Q-9.

## MATERIALS AND METHODS

*Microorganisms and cultivation*

The yeast strains used in this study are listed in Table 1. The cultures were cultivated as described previously (Yamada and Kondo, 1973). However, the psychrophilic strains of *Brettanomyces nanus* nom. nud. were cultured at room temperature.

*Extraction and purification of coenzyme Q*

The Co-Q was extracted from intact cells and partially purified by preparative thin layer chromatography (Yamada and Kondo, 1973).

*Determination of the coenzyme Q system*

Reversed phase paper chromatography was used for the preparation of Co-Q. Mass spectrometry was applied for the determination of the Co-Q system (Yamada and Kondo, 1973).

*Reagents and chemicals*

Authentic preparations of the Co-Q series (Q-6 to Q-10) were obtained as described in our preceding papers (Yamada and Kondo, 1973; Yamada et al., 1973a,b,c, 1976a,b). Other reagents and chemicals were commercial preparations (Nakarai).

Table 1. The coenzyme Q system in the genera *Dekkera* and *Brettanomyces*.

Species and strain	Other remarks	Co-Q system
<i>D. bruxellensis</i> CBS 74	ATCC 36234, IFO 1590, type	Q-9
<i>D. intermedia</i> CBS 4914	ATCC 36235, type	Q-9
<i>B. abstinens</i> CBS 6055	ATCC 22341, IFO 1589, type	Q-9
<i>B. anomalus</i> CBS 77	ATCC 10559, NRRL Y-1415, IFO 0796, type	Q-9
<i>B. bruxellensis</i> CBS 72	ATCC 10560, NRRL Y-1411, type	Q-9
<i>B. clausenii</i> CBS 76	ATCC 10562, NRRL Y-1414, IFO 0627, type	Q-9
<i>B. custersianus</i> CBS 4805	IFO 1585, type	Q-9
<i>B. custersii</i> CBS 5512	IFO 1586, neotype	Q-9
<i>B. intermedius</i> CBS 73	IFO 1587, type	Q-9
<i>B. lambicus</i> CBS 75	ATCC 10563, NRRL Y-1413, IFO 0797, type	Q-9
<i>B. naardenensis</i> CBS 6042	ATCC 22075, IFO 1588, type	Q-9
<i>B. nanus</i> nom. nud. CBS 1945	Scheffers (1966)	Q-9
CBS 1955		Q-9
CBS 1956		Q-9

CBS, Centraalbureau voor Schimmelcultures, Delft, The Netherlands; IFO, Institute for Fermentation, Osaka, Japan; ATCC, American Type Culture Collection, Rockville, Maryland, U.S.A.; NRRL, ARS Culture Collection, Northern Regional Research Center, U.S. Department of Agriculture, Peoria, Illinois, U.S.A.

## RESULTS AND DISCUSSION

The strains examined in the genera *Dekkera* and *Brettanomyces* were found to have Q-9 without exception (Table 1). This indicates that these two genera are, in this respect, taxonomically homogeneous. They have the Q-9 system in common with representatives of the genera *Debaryomyces*, *Schwanniomyces*, *Lodderomyces*, *Lipomyces*, *Wickerhamia*, *Endomyces* and *Metschnikowia*, and with some strains in the genera *Pichia* and *Schizosaccharomyces* (Yamada et al., 1973*b,c*, 1976*a,b*, 1977). However, the genera *Dekkera* and *Brettanomyces* differ in the Co-Q system from representatives of the genera *Hansenula* (Q-7 and Q-8), *Saccharomyces* (Q-6), *Kluyveromyces* (Q-6), *Nadsonia* (Q-6), *Saccharomycodes* (Q-6), *Hanseniaspora* (Q-6), *Kloeckera* (Q-6), *Nematospora* (Q-6) and from the majority of the strains in the genera *Pichia* (Q-7 and Q-8; some strains Q-9) and *Schizosaccharomyces* (Q-10; some strains Q-9).

Van der Walt (1963, 1970*a*) suggested that the genera *Dekkera* and *Brettanomyces* and the genera *Hanseniaspora* and *Kloeckera* would be derived from a common progenitor, based on the facts that all four genera utilize nitrite slowly in low concentration, that some species of the ascosporeogenous genera *Dekkera* and *Hanseniaspora* produce hat-shaped, easily liberated ascospores, and that *Hanseniaspora valbyensis* is able to form notable amounts of acetic acid. Fiol and Billon-Grand (1978) proposed several phylogenetic lines among the species of the genera *Dekkera*, *Brettanomyces*, *Hanseniaspora* and *Kloeckera* on the basis of activities of intracellular glycosidases and nitrate and nitrite reductases.

Spencer and Gorin (1969) determined the proton magnetic resonance spectra of mannans of several *Brettanomyces* species. They accommodated the *Brettanomyces* species in subgroup b of group X or the *Saccharomyces microellipsoides* group. These *Brettanomyces* species have the spectra with signals at  $\tau$  4.30 and  $\tau$  4.40, which are common to *Kluyveromyces lodderi*, *Hanseniaspora osmophila* and *Saccharomyces carlsbergensis*. These three species have coenzyme Q-6 (Yamada et al., 1976*a,b*; unpublished data).

Having tested strains of all the known species of the genera *Dekkera* and *Brettanomyces*, we have found that their Co-Q system differs considerably from that of the genera *Hanseniaspora* and *Kloeckera*. Despite the similarity of the production of hat-shaped ascospores which are easily liberated from asci, the difference from Q-9 to Q-6 tends to exclude the possibility that these genera are closely related (Fig. 1).

Received 20 June 1980




Fig. 1. Schematic representation of the coenzyme Q system in the genera *Dekkera*, *Brettanomyces*, *Hanseniaspora* and *Kloeckera*. These genera are divided into two groups, namely, the Q-9 and the Q-6 possessing organisms.

#### REFERENCES

- FIOL, J. B. and BILLON-GRAND, G. 1978. Etude de quelques enzymes intra-cellulaires dans les genres *Dekkera* et *Brettanomyces*; conséquences systématiques. — *Mycopathologia* **64**: 183–186.
- SCHEFFERS, W. A. 1966. Stimulation of fermentation in yeasts by acetoin and oxygen. — *Nature* **210**: 533–534.
- SCHEFFERS, W. A. and WIKÉN, T. O. 1969. The Custers effect (negative Pasteur effect) as a diagnostic criterion for the genus *Brettanomyces*. — *Antonie van Leeuwenhoek* **35**, Supplement Yeast Symposium: A 31–A 32.
- SPENCER, J. F. T. and GORIN, P. A. J. 1969. Systematics of the genera *Saccharomyces*, *Schizosaccharomyces*, *Endomycopsis*, *Kluyveromyces*, *Schwanniomyces* and *Brettanomyces*: proton magnetic resonance spectra of the mannans and mannose-containing polysaccharides as an aid in classification. — *Antonie van Leeuwenhoek* **35**: 361–378.
- VAN DER WALT, J. P. 1963. Nitrite utilization by *Brettanomyces*. — *Antonie van Leeuwenhoek* **29**: 52–56.
- VAN DER WALT, J. P. 1970a. Genus 4. *Dekkera* van der Walt, p. 157–165. In J. Lodder (ed.), *The yeasts. A taxonomic study*. — North-Holland Publishing Co., Amsterdam.
- VAN DER WALT, J. P. 1970b. Genus 1. *Brettanomyces* Kufferath et van Laer, p. 863–892. In J. Lodder (ed.), *The yeasts. A taxonomic study*. — North-Holland Publishing Co., Amsterdam.
- WIKÉN, T., SCHEFFERS, W. A. and VERHAAR, A. J. M. 1961. On the existence of a negative Pasteur effect in yeasts classified in the genus *Brettanomyces* Kufferath et van Laer. — *Antonie van Leeuwenhoek* **27**: 401–433.
- YAMADA, Y. and KONDO, K. 1972a. Taxonomic significance of coenzyme Q system in yeasts and yeast-like fungi (1). — *Proc. 1st Special. Int. Symp. Yeasts*, Smolenice: 363–373.
- YAMADA, Y. and KONDO, K. 1972b. Taxonomic significance of coenzyme Q system in yeasts and yeast-like fungi (2). — *Proc. 4th Int. Ferment. Symp.*, Kyoto: 781–784.
- YAMADA, Y. and KONDO, K. 1972c. Taxonomic significance of the coenzyme Q system in yeasts and yeast-like fungi (3). — *Proc. 2nd Special. Int. Symp. Yeasts*, Tokyo: 63–69.
- YAMADA, Y. and KONDO, K. 1973. Coenzyme Q system in the classification of the yeast genera *Rhodotorula* and *Cryptococcus*, and the yeast-like genera *Sporobolomyces* and *Rhodospiridium*. — *J. Gen. Appl. Microbiol.* **19**: 59–77.
- YAMADA, Y., KANEMATSU, Y., OHASHI, M. and KONDO, K. 1973a. On the partly reduced coenzyme Q isolated from “*Rhodotorula lactosa*” IFO 1058 and its relation to the taxonomic position. — *Agr. Biol. Chem.* **37**: 621–628.

- YAMADA, Y., OKADA, T., UESHIMA, O. and KONDO, K. 1973*b*. Coenzyme Q system in the classification of the ascosporeogenous yeast genera *Hansenula* and *Pichia*. — J. Gen. Appl. Microbiol. **19**: 189–208.
- YAMADA, Y., ARIMOTO, M. and KONDO, K. 1973*c*. Coenzyme Q system in the classification of the ascosporeogenous yeast genus *Schizosaccharomyces* and yeast-like genus *Endomyces*. — J. Gen. Appl. Microbiol. **19**: 353–358.
- YAMADA, Y., ARIMOTO, M. and KONDO, K. 1976*a*. Coenzyme Q system in the classification of apiculate yeasts in the genera *Nadsonia*, *Saccharomycodes*, *Hanseniaspora*, *Kloeckera*, and *Wickerhamia*. — J. Gen. Appl. Microbiol. **22**: 293–299.
- YAMADA, Y., NOJIRI, M., MATSUYAMA, M. and KONDO, K. 1976*b*. Coenzyme Q system in the classification of the ascosporeogenous yeast genera *Debaryomyces*, *Saccharomyces*, *Kluyveromyces*, and *Endomycopsis*. — J. Gen. Appl. Microbiol. **22**: 325–337.
- YAMADA, Y., ARIMOTO, M. and KONDO, K. 1977. Coenzyme Q system in the classification of some ascosporeogenous yeast genera in the families Saccharomycetaceae and Sporophthoraceae. — Antonie van Leeuwenhoek **43**: 65–71.