

Reduced IGF-1 Signaling Delays Age-Associated Proteotoxicity in Mice

Ehud Cohen, Johan F. Paulsson, Pablo Blinder, Tal Burstyn-Cohen, Deguo Du, Gabriela Estepa, Anthony Adame, Hang M. Pham, Martin Holzenberger, Jeffery W. Kelly, Eliezer Masliah, and Andrew Dillin*

*Correspondence: dillin@salk.edu

DOI 10.1016/j.cell.2010.02.031

(Cell 139, 1157–1169; December 11, 2009)

During the revision of our manuscript, two research groups independently reported that reduced IGF signaling is protective toward Alzheimer's disease-related pathology. We would like to direct readers to the following references:

Freude, S., Hettich, M.M., Schumann, C., Stöhr, O., Koch, L., Köhler, C., Udelhoven, M., Leeser, U., Müller, M., Kubota, N., et al. (2009). Neuronal IGF-1 resistance reduces Abeta accumulation and protects against premature death in a model of Alzheimer's disease. *FASEB J.* 23, 3315–3324.

Killick, R., Scales, G., Leroy, K., Causevic, M., Hooper, C., Irvine, E.E., Choudhury, A.I., Drinkwater, L., Kerr, F., Al-Qassab, H., et al. (2009). Deletion of *Irs2* reduces amyloid deposition and rescues behavioural deficits in APP transgenic mice. *Biochem. Biophys. Res. Commun.* 386, 257–262.