


Current awareness on yeast

(5 weeks journals - Search completed at 13th. Aug. 2008)

Current Awareness – Use the links online

View this bibliography online at www.interscience.wiley.com

The articles use  the innovative multi-publisher reference linking. This enables readers to move seamlessly from the reference to the cited publication, typically located on a different server and published by a different publisher. Links to MEDLINE abstracts are also available for key journals in the biosciences.

7487

I Reviews

- Dickson RC. (2008). Univ Kentucky, Dept Mol & Cellular Biochem, Lexington, Ky 40536, USA. New insights into sphingolipid metabolism and function in budding yeast. *J Lipid Res* **49**, (5) 909
- Fonseca GG, Heinze E, Wittmann C, Gombert AK*. (2008). *Delft Univ Technol, Dept Biotechnol, Julianalaan 67, NL-2628 BC Delft, The Netherlands. The yeast *Kluyveromyces marxianus* and its biotechnological potential. *Appl Microbiol Biotechnol* **79**, (3) 339
- Hedbacker K, Carlson M*. (2008). *Columbia Univ, Dept Genet & Dev, 701 West 168th St, New York, NY 10032, USA. SNF1/AMPK pathways in yeast. *Front Biosci* **13**, 2408
- Kutty SN, Philip R*. (2008). *Cochin Univ Sci & Technol, Sch Ocean Sci & Technol, Dept Marine Biol Microbiol & Biochem, Fine Arts Ave, IN-682016 Kochi, India. Marine yeasts - A review. *Yeast* **25**, (7) 465
- Matmati N, Hannun YA*. (2008). *Med Univ Sth Carolina, Dept Biochem & Mol Biol, Charleston, SC 29425, USA. Thematic review series: Sphingolipids. *ISC1* (inositol phosphosphingolipid-phospholipase C), the yeast homologue of neutral sphingomyelinases. *J Lipid Res* **49**, (5) 922
- Wang YC. (2008). Florida State Univ, Dept Biomed Sci, 1115 West College St, Tallahassee, FL 32306, USA. Chromosome instability in yeast and its implications to the study of human cancer. *Front Biosci* **13**, 2091
- Wedlich-Soldner R, Li R. (2008). Max Planck Inst Biochem, Klopferspitz 18, DE-82152 Martinsried, Germany. Yeast and fungal morphogenesis from an evolutionary perspective. *Semin Cell Dev Biol* **19**, (3) 224
- Zhang YQ, Rao R*. (2008). *Johns Hopkins Univ, Dept Physiol, 725 Nth Wolfe St, Baltimore, Md 21205, USA. A spoke in the wheel - Calcium spikes disrupt yeast cell cycle. *Cell Cycle* **7**, (7) 870

2 General

- Jacobsen MD, Davidson AD, Li SY, Shaw DJ, Gow NAR, Odds FC*. (2008). *Institute Med Sci, Aberdeen Fungal Grp, Aberdeen AB25 2ZD, Scotland. Molecular phylogenetic analysis of *Candida tropicalis* isolates by multi-locus sequence typing. *Fungal Genet Biol* **45**, (6) 1040

Nagahama T, Abdel-Wahab MA, Nogi Y, Miyazaki M, Uematsu K, Hamamoto M, Horikoshi K. (2008). Japan Agcy Marine Earth Sci & Technol, Extremobiosphere Res Ctr, 2-15 Natsushimacho, Yokosuka, Kanagawa 237 006, Japan. *Dipodascus tetrasporeus* sp nov., an ascosporogenous yeast isolated from deep-sea sediments in the Japan Trench. *Int J Syst Evol Microbiol* **58**, (4) 1040

Tsai IJ, Bensasson D, Burt A, Koufopanou V*. (2008). *Imperial Coll London, Div Biol, Silwood Pk, Ascot SL5 7PY, England. Population genomics of the wild yeast *Saccharomyces paradoxus*: Quantifying the life cycle. *Proc Natl Acad Sci U S A* **105**, (12) 4957

3 Biochemistry

- Bayliak M, Gospodaryov D, Semchyshyn H, Lushchak V*. (2008). *Vassyl Stefanyk Precarpathian Natl Univ, Dept Biochem, 57 Shevchenko Str, UA-76025 Ivano Frankivsk, Ukraine. Inhibition of catalase by aminotriazole *in vivo* results in reduction of glucose-6-phosphate dehydrogenase activity in *Saccharomyces cerevisiae* cells. *Biochemistry (Mosc)* **73**, (4) 420
- Dian EA, Papatheodorou P, Emmrich K, Randel O, Geissler A, Kolling R, Rassow J*, Motz C. (2008). *Ruhr Univ Bochum, Inst Physiol Chem, Abt Zellbiochem, DE-44780 Bochum, Germany. Role of γ -subunit N- and C-termini in assembly of the mitochondrial ATP synthase in yeast. *J Mol Biol* **377**, (5) 1314
- Grintsevich EE, Benchaar SA, Warshaviak D, Boonthueung P, Halgand F, Whitelegge JP, Faull KF, Loo RRO, Sept D, Loo JA, Reisler E*. (2008). *UCLA, Dept Chem & Biochem, 405 Hilgard Ave, Los Angeles, Ca 90024, USA. Mapping the cofilin binding site on yeast G-actin by chemical cross-linking. *J Mol Biol* **377**, (2) 395
- Huang LY, Umanah G, Hauser M, Son C, Arshava B, Naider F, Becker JM*. (2008). *Univ Tennessee, Dept Microbiol, Knoxville, Tn 37996, USA. Unnatural amino acid replacement in a yeast G protein-coupled receptor in its native environment. *Biochemistry* **47**, (20) 5638
- Naidoo N, Harrop SJ, Sobti M, Haynes PA, Szymczynska BR, Williamson JR, Curmi PMG, Mabbutt BC*. (2008). *MacQuarie Univ, Dept Chem & Biomol Sci, Nth Ryde, NSW 2109, Australia. Crystal structure of Lsm3 octamer from *Saccharomyces cerevisiae*: Implications for Lsm ring organisation and recruitment. *J Mol Biol* **377**, (5) 1357
- Perrot M, Massoni A, Boucherie H*. (2008). *IBGC, UMR CNRS 5095, 1 rue Camille Saint-Saens, FR-33077 Bordeaux, France. Sequence requirements for N^o-terminal acetylation of yeast proteins by NatA. *Yeast* **25**, (7) 513

In order to keep subscribers up-to-date with the latest developments in their field, this current awareness service is provided by John Wiley & Sons and contains newly-published material on yeasts. Each bibliography is divided into 10 sections. 1 Reviews; 2 General; 3 Biochemistry; 4 Biotechnology; 5 Cell Biology; 6 Gene Expression; 7 Genetics & Genomics; 8 Physiology; 9 Medical Mycology; 10 Recombinant DNA Technology. Within each section, articles are listed in alphabetical order with respect to author. If, in the preceding period, no publications are located relevant to any one of these headings, that section will be omitted.

- Poveda A, Sendra R*. (2008). *Univ Valencia, Dept Bioquim & Biol Mol, C/ Dr Moliner 50, Valencia, Spain. Site specificity of yeast histone acetyltransferase B complex *in vivo*. *FEBS J* **275**, (9) 2122
- Ruckenstuhl C, Poschenel A, Possert R, Baral PK, Gruber K, Turnowsky F*. (2008). *Karl Franzens Univ, Inst Mol Biosci, Humboldtstr 50, AT-8010 Graz, Austria. Structure-function correlations of two highly conserved motifs in *Saccharomyces cerevisiae* squalene epoxidase. *Antimicrob Agents Chemother* **52**, (4) 1496
- Vashishtha AK, West AH, Cook PF*. (2008). *Univ Oklahoma, Dept Chem & Biochem, Parrington Oval, Norman, Ok 73019, USA. Overall kinetic mechanism of saccharopine dehydrogenase (L-glutamate forming) from *Saccharomyces cerevisiae*. *Biochemistry* **47**, (19) 5417
- Yokoyama K, Nakagawa M, Satoh M, Saitoh S, Dohmae N, Harada A, Satoh N, Karasawa K, Takio K, Yanagida M, Inoue K. (2008). Teikyo Univ, Fac Pharmaceut Sci, Sagami-hara, Kanagawa 229 019, Japan. Expression of a novel 90-kDa protein, Lsd90, involved in the metabolism of very long-chain fatty acid-containing phospholipids in a mitosis-defective fission yeast mutant. *J Biochem* **143**, (3) 369
- Yu ZH, Gonciarz MD, Sundquist WI, Hill CP, Jensen GJ*. (2008). *Calif Inst Technol, Div Biol, 1200 East California Blvd, Pasadena, Ca 91125, USA. Cryo-EM structure of dodecameric Vps4p and its 2:1 complex with Vta1p. *J Mol Biol* **377**, (2) 364
- Zupancic ML, Frieman M, Smith D, Alvarez RA, Cummings RD, Cormack BP*. (2008). *Johns Hopkins Univ, Dept Mol Biol & Genet, Baltimore, Md 21205, USA. Glycan microarray analysis of *Candida glabrata* adhesin ligand specificity. *Mol Microbiol* **68**, (3) 547
- ## 4 Biotechnology
- Alloue WAM, Destain J, El Medjoub T, Ghalfi H, Kabran P, Thonart P. (2008). Fac Univ Sci Agron Gembloux, Unite Bioind, CWBI, Passage Deportes 2, BE-5030 Gembloux, Belgium. Comparison of *Yarrowia lipolytica* lipase immobilization yield of entrapment, adsorption, and covalent bond techniques. *Appl Biochem Biotechnol* **150**, (1) 51
- Antosova M, Illeova V, Vandakova M, Druzkovska A, Polakovic M*. (2008). *Slovak Tech Univ Bratislava, Inst Chem & Environm Engn, SK-81237 Bratislava, Slovakia. Chromatographic separation and kinetic properties of fructosyltransferase from *Aureobasidium pullulans*. *J Biotechnol* **135**, (1) 58
- Aragon CC, Ferreira-Dias S, Gattas EAD, Peres MDS*. (2008). *UNESP, Dept Food & Nutr, Rod Araraquara-Jau km 1, BR-14801-902 Sao Paulo, Brazil. Characterization of glycerol kinase from baker's yeast: Response surface modeling of the enzymatic reaction. *J Mol Catal B Enzym* **52-53**, 113
- Arroyo-Lopez FN, Bautista-Gallego J, Duran-Quintana MC, Garrido-Fernandez A. (2008). Inst Grasa - CSIC, Dept Biotechnol Alimentos, Avda Padre Garcia Tejero 4, ES-41012 Seville, Spain. Modelling the inhibition of sorbic and benzoic acids on a native yeast cocktail from table olives. *Food Microbiol* **25**, (4) 566
- Baerends RJS, De Hulster E, Geertman JMA, Daran JM, Van Maris AJA, Veenhuis M, Van der Klei IJ, Pronk JT*. (2008). *Delft Univ Technol, Dept Biotechnol, Julianalaan 67, NL-2628 BC Delft, The Netherlands. Engineering and analysis of a *Saccharomyces cerevisiae* strain that uses formaldehyde as an auxiliary substrate. *Appl Environ Microbiol* **74**, (10) 3182
- Ben Akacha N, Zehlila A, Merji S, Jerbi T, Gargouri M. (2008). Natl Inst Appl Sci & Technol - INSAT, Biocatalysis & Bioprocess Unit, Enzyme Engn Grp, Tunis, Tunisia. Effect of gamma-ray on activity and stability of alcohol-dehydrogenase from *Saccharomyces cerevisiae*. *Biochem Eng J* **40**, (1) 184
- Bender JP, Mazutti MA, Di Luccio M, Treichel H*. (2008). *Univ Regional Integrada Aldo Uruguai & Missoes, Dept Engn & Alimentos, BR-99700-000 Erechim, Brazil. Extraction of inulinase obtained by solid state fermentation of sugarcane bagasse by *Kluyveromyces marxianus* NRRL Y-7571. *Appl Biochem Biotechnol* **149**, (3) 195
- Blanco CA, Rayo J, Giralda JM. (2008). Univ Valladolid, Dept Ingn Agrícola & Forestal, Area Tecnol Alimentos, ES-34004 Palencia, Spain. Improving industrial full-scale production of baker's yeast by optimizing aeration control. *J AOAC Int* **91**, (3) 607
- Blazejak S, Duszkievicz-Reinhard W, Gniewosz M*, Wiatrik P. (2008). *Agric Univ, Dept Biotechnol Microbiol & Food Evaluation, ul Nowoursynowska 159 C, PL-02776 Warsaw, Poland. Impact of magnesium and mannose in the cultivation media on the magnesium biosorption, the biomass yield and on the cell wall structure of *Candida utilis* yeast. *Eur Food Res Technol* **227**, (3) 695
- Bolumar T, Sanz Y, Aristoy MC, Toldra F*. (2008). *Inst Agroquim & Tecnol Alimentos - CSIC, POB 73, ES-46100 Valencia, Spain. Purification and characterisation of proteases A and D from *Debaryomyces hansenii*. *Int J Food Microbiol* **124**, (2) 135
- Camargo LA, Ribeiro MHL, Peres MDS, Gattas EAD*. (2008). *UNESP, Dept Food & Nutr, Rod Araraquara-Jau km 1, BR-14801-902 Sao Paulo, Brazil. Kinetic properties of glycerophosphate oxidase isolated from dry baker's yeast. *J Mol Catal B Enzym* **52-53**, 140
- Chen ZY, Wang ZY, He XP*, Guo X, Li WW, Zhang B. (2008). *CAS, Inst Microbiol, Lab Yeast Mol Genet & Breeding, Datun Rd, CN-100101 Beijing, Peoples Rep China. Uricase production by a recombinant *Hansenula polymorpha* strain harboring *Candida utilis* uricase gene. *Appl Microbiol Biotechnol* **79**, (4) 545
- Cus F, Raspor P. (2008). Agric Inst Slovenia, Cent Labs, Hacquetova 17, SI-1000 Ljubljana, Slovenia. The effect of pyrimethanil on the growth of wine yeasts. *Lett Appl Microbiol* **47**, (1) 54
- De Paula FC, Cazetta ML, Monti R, Contiero J*. (2008). *Sao Paulo State Univ, Biol Sci Inst, Dept Biochem & Microbiol, Av 24A 1515, BR-13506-900 Rio Claro, Brazil. Sucrose hydrolysis by gelatin-immobilized inulinase from *Kluyveromyces marxianus* var. *bulgaricus*. *Food Chem* **111**, (3) 691
- Elbandy M, Shinde PB, Dang HT, Hong J, Bae KS, Jung JH*. (2008). *Pusan Natl Univ, Coll Pharm, Pusan, South Korea. Furan metabolites from the sponge-derived yeast *Pichia membranifaciens*. *J Nat Prod* **71**, (5) 869
- Escalante-Minakata P, Blaschek HP, De la Rosa APB, Santos L, De Leon-Rodriguez A*. (2008). *Inst Potosino Invest Cient & Tecnol, Div Mol Biol, AP 3-74 Tangamanga, MX-78231 San Luis Potosi, Mexico. Identification of yeast and bacteria involved in the mezcalt fermentation of *Agave salmiana*. *Lett Appl Microbiol* **46**, (6) 626
- Garofalo C, Silvestri G, Aquilanti L, Clementi F. (2008). Polytech Univ Marche, Dept Food Sci, Via Brecce Bianche, IT-60131 Ancona, Italy. PCR-DGGE analysis of lactic acid bacteria and yeast dynamics during the production processes of three varieties of Panettone. *J Appl Microbiol* **105**, (1) 243
- Gibson BR, Prescott KA, Smart KA*. (2008). *Univ Nottingham, Div Food Sci, Sutton Bonington Campus, Loughborough LE12 5RD, England. Petite mutation in aged and oxidatively stressed ale and lager brewing yeast. *Lett Appl Microbiol* **46**, (6) 636
- Guimaraes PMR, Multanen JP, Domingues L, Teixeira JA, Londerborough J. (2008). Univ Minho, Inst Biotechnol & Bioengn, Ctr Biol Engn, Campus Gualtar, PT-4710-057 Braga, Portugal. Stimulation of zero-*trans* rates of lactose and maltose uptake into yeasts by preincubation with hexose to increase the adenylate energy charge. *Appl Environ Microbiol* **74**, (10) 3076
- Harms J, Wang XY, Kim T, Yang XM, Rathore AS*. (2008). *Amgen Inc, Process Dev, Thousand Oaks, Ca 91320, USA. Defining process design space for biotech products: Case study of *Pichia pastoris* fermentation. *Biotechnol Prog* **24**, (3) 655
- Hernalsteens S, Mauger F. (2008). Univ Estadual Campinas, Dept Food Engn, CP 6121, BR-13083-970 Campinas, SP, Brazil. Purification and characterisation of a fructosyltransferase from *Rhodotorula* sp. *Appl Microbiol Biotechnol* **79**, (4) 589
- Holmgren M, Sellstedt A*. (2008). *Umea Univ, Dept Plant Physiol, Plant Sci Ctr, SE-90187 Umea, Sweden. Identification of white-rot and soft-rot fungi increasing ethanol production from spent sulfite liquor in co-culture with *Saccharomyces cerevisiae*. *J Appl Microbiol* **105**, (1) 134
- Huang XJ, Yu AG, Xu ZK*. (2008). *Zhejiang Univ, Inst Polymer Sci, MoE Key Lab Macromol Synth & Functionalization, CN-310027 Hangzhou, Peoples Rep China. Covalent immobilization of lipase from *Candida rugosa* onto poly(acrylonitrile-co-2-hydroxyethyl methacrylate) electrospun fibrous membranes for potential bioreactor application. *Bioresour Technol* **99**, (13) 5459

- Jeyaram K, Singh WM, Capece A, Romano P. (2008). IBSD, Takyelpat Inst Area, Microbial Resources Div, IN-795001 Imphal, India. Molecular identification of yeast species associated with 'Hamei'—A traditional starter used for rice wine production in Manipur, India. *Int J Food Microbiol* **124**, (2) 115
- Kinay P, Yildiz M. (2008). Ege Univ, Dept Plant Protect, TR-35100 Izmir, Turkey. The shelf life and effectiveness of granular formulations of *Metschnikowia pulcherrima* and *Pichia guilliermondii* yeast isolates that control postharvest decay of citrus fruit. *Biol Control* **45**, (3) 433
- Leskinen P, Hilscherova K, Sidlova T, Kiviranta H, Pessala P, Salo S, Verta M, Virta M*. (2008). *Univ Helsinki, Dept Appl Chem & Microbiol, POB 56, FI-00014 Helsinki, Finland. Detecting AhR ligands in sediments using bioluminescent reporter yeast. *Biosens Bioelectron* **23**, (12) 1850
- Li GY, Jiang YR*, Huang KL, Ding P, Yao LL. (2008). *Cent Sth Univ Technol, Coll Chem & Chem Engr, CN-410083 Changsha, Hunan, Peoples Rep China. Kinetics of adsorption of *Saccharomyces cerevisiae* mandelated dehydrogenase on magnetic Fe₃O₄-chitosan nanoparticles. *Colloid Surface A* **320**, (1-3) 11
- Libkind D, Gadanho M, Van Broock M, Sampalo JP. (2008). Univ Nacl Comahue, CONICET, Lab Microbiol Aplicada & Biotecnol, CRUB, Quintral 1250, AR-8400 San Carlos de Bariloche, Rio Negro, Argentina. Studies on the heterogeneity of the carotenogenic yeast *Rhodotorula mucilaginosa* from Patagonia, Argentina. *J Basic Microbiol* **48**, (2) 93
- Marques S, Alves L, Roseiro JC, Girio FM*. (2008). *INETI, Dept Biotechnol, Estrada Paco do Lumiar 22, PT-1649-038 Lisbon, Portugal. Conversion of recycled paper sludge to ethanol by SHF and SSF using *Pichia stipitis*. *Biomass Bioeng* **32**, (5) 400
- Maskow T, Roellich A, Fetzler I, Ackermann JU, Harms H. (2008). UFZ, Dept Environm Microbiol, Helmholtz Ctr Environm Res, Permoserstr 15, DE-04318 Leipzig, Germany. On-line monitoring of lipid storage in yeasts using impedance spectroscopy. *J Biotechnol* **135**, (1) 64
- Matsushika A, Watanabe S, Kodaki T, Makino K, Sawayama S. (2008). AIST, BTRC, 2-2-2 Hirosuehiro, Hiroshima 737 0197, Japan. Bioethanol production from xylose by recombinant *Saccharomyces cerevisiae* expressing xylose reductase, NADP⁺-dependent xylitol dehydrogenase, and xylulokinase. *J Biosci Bioeng* **105**, (3) 296
- Mijatovic S, Gadda G*. (2008). *Georgia State Univ, Dept Chem, POB 4098, Atlanta, Ga 30302, USA. Oxidation of alkyl nitronates catalyzed by 2-nitropropane dioxygenase from *Hansenula mrakii*. *Arch Biochem Biophys* **473**, (1) 61
- Modig T, Almeida JRM, Gorwa-Grauslund MF, Liden G. (2008). Lund Univ, Dept Chem Engr, POB 124, SE-22100 Lund, Sweden. Variability of the response of *Saccharomyces cerevisiae* strains to lignocellulose hydrolysate. *Biotechnol Bioeng* **100**, (3) 423
- Musgrove MT, Jones DR, Hinton A, Ingram KD, Northcutt JK. (2008). NSDA/ARS, Egg Safety & Qual Res Unit, 950 College Stn Rd, Athens, Ga 30605, USA. Identification of yeasts isolated from commercial shell eggs stored at refrigerated temperatures. *J Food Prot* **71**, (6) 1258
- Ni XM, Chi ZM*, Liu ZQ, Yue LX. (2008). *Ocean Univ Qingdao, UNESCO, Chinese Ctr Marine Biotechnol, CN-266003 Qingdao, Peoples Rep China. Screening of protease-producing marine yeasts for production of the bioactive peptides. *Acta Oceanol Sin* **27**, (2) 116
- Nie Y, Xu Y*, Wang HY, Xu N, Xiao R, Sun ZH. (2008). *Sthn Yangtze Univ, MoE Key Lab Ind Biotechnol, 170 Huihe Rd, CN-214036 Wuxi, Peoples Rep China. Complementary selectivity to (S)-1-phenyl-1,2-ethanediol-forming *Candida parapsilosis* by expressing its carbonyl reductase in *Escherichia coli* for (R)-specific reduction of 2-hydroxyacetophenone. *Biocatal Biotransform* **26**, (3) 210
- Oba T, Suenaga H, Muta S, Tashiro K, Kuhara S. (2008). Fukuoka Ind Technol Ctr, Biotechnol & Food Res Inst, 1465-5 Aikawa, Fukuoka 839 0861, Japan. Properties of a sucrose-tolerant mutant of *Saccharomyces cerevisiae*. *World J Microbiol Biotechnol* **24**, (7) 1233
- Paez G, Jimenez E, Marmol Z, Ferrer J, Sulbaran B, Ojeda G, Araujo K*, Rincon M. (2008). *Univ Zulia, Lab Tecnol Alimentos & Fermentacion Ind, Apdo 526, VE-4011 Maracaibo, Estado Zulia, Venezuela. Amino acids profile of the single cell protein from *Kluyveromyces marxianus* var. *marxianus* (Spanish, English Abstract). *Interiencia* **33**, (4) 297
- Panda T, Gowrishankar BS. (2008). IIT, Biochem Engr Lab, IN-600036 Chennai, India. Clinical analysis of application of generalized distance function for optimization of important variables for esterase synthesis by *Saccharomyces cerevisiae*. *Bioresource Technol* **99**, (13) 5545
- Panesar PS. (2008). Sant Longowal Inst Engr & Technol, Dept Food Technol, IN-148106 Longowal, India. Application of response surface methodology for maximal lactose hydrolysis in whole milk using permeabilised yeast cells. *Acta Aliment* **37**, (2) 191
- Perles CE, Moran PJS, Volpe PLO. (2008). UNICAMP, Inst Quim, CP 6154, BR-13083-970 Campinas, SP, Brazil. Bioreduction of ethyl 3 oxybutyrate by *Saccharomyces cerevisiae*. A metabolic *in vivo* study. *J Mol Catal B Enzym* **52-53**, 82
- Pickering GJ, Nikfardjam MSP. (2008). Brock Univ, Dept Biol Sci, 500 Glenridge Ave, St Catharines, Ontario, Canada L2S 3A1. Influence of variety and commercial yeast preparation on red wine made from autochthonous Hungarian and Canadian grapes. Part II. Oral sensations and sensory : instrumental relationships. *Eur Food Res Technol* **227**, (3) 925
- Plessas S, Bosnea L, Psarianos C, Koutinas AA, Marchant R, Banat IM. (2008). Univ Patras, Dept Chem, Food Biotechnol Grp, GR-26500 Patras, Greece. Lactic acid production by mixed cultures of *Kluyveromyces marxianus*, *Lactobacillus delbrueckii* ssp *bulgaricus* and *Lactobacillus helveticus*. *Bioresource Technol* **99**, (13) 5951
- Popp A, Nguyen HTT, Boulahya K, Bideaux C, Alfenore S, Guillouet SE, Nevoigt E*. (2008). *KUL, Inst Bot & Microbiol, Kasteelpk Arenberg 31, Bus 2438, BE-3001 Heverlee, Belgium. Fermentative production of *l*-glycerol 3-phosphate utilizing a *Saccharomyces cerevisiae* strain with an engineered glycerol biosynthetic pathway. *Biotechnol Bioeng* **100**, (3) 497
- Raman N, Pothiraj C. (2008). VHNSN Coll, Dept Chem, IN-626001 Virudunagar, India. One step process for starch fermentation employing the starch digesting yeast *Saccharomyces diastaticus*. *J Indian Chem Soc* **85**, (4) 452
- Rao RS, Bhadra B, Shivaji S*. (2008). *Ctr Cellular & Mol Biol, Uppal Rd, IN-500007 Hyderabad, India. Isolation and characterization of ethanol-producing yeasts from fruits and tree barks. *Lett Appl Microbiol* **47**, (1) 19
- Romano A, Perello MC, De Revel G, Lonvaud-Funel A. (2008). Via 25 Aprile 3, IT-20036 Meda, Italy. Growth and volatile compound production by *Brettanomyces/Dekkera bruxellensis* in red wine. *J Appl Microbiol* **104**, (6) 1577
- Rustoy EM, Cerrutti P, Galvagno MA, Baldessari A*. (2008). *UBA, Fac Ciencias Exactas & Nat, Dept Quim Organ, AR-1428 Buenos Aires, DF, Argentina. An efficient biotransformation of dialkyl esters of 2-oxoglutaric acid by *Rhodotorula minuta* whole cells. *Biocatal Biotransform* **26**, (3) 204
- Rymowicz W, Rywinska A, Gladkowski W. (2008). Wroclaw Univ Environm & Life Sci, Dept Biotechnol & Food Microbiol, PL-50375 Wroclaw, Poland. Simultaneous production of citric acid and erythritol from crude glycerol by *Yarrowia lipolytica* Wratislavia K1. *Chem Pap* **62**, (3) 239
- Sangorin MP, Lopes CA, Jofre V, Querol A, Caballero AC*. (2008). *Univ Nacl Comahue, Dept Quim, Lab Microbiol & Biotechnol, AR-1400(8300) Buenos Aires, Neuquen, Argentina. Spoilage yeasts from Patagonian cellars: Characterization and potential biocontrol based on killer interactions. *World J Microbiol Biotechnol* **24**, (7) 945
- Saxon RE, Leisch H, Hudlicky T*. (2008). *Brock Univ, Dept Chem, 500 Glenridge Ave, St Catharines, Ontario, Canada L2S 3A1. Preliminary investigation of the yeast-mediated reduction of β -keto amides derived from cyclic amines as potential resolution methodology. *Tetrahedron Asymmetry* **19**, (6) 672
- Schafer B, Neffgen A, Kliner U. (2008). Rhein Westfal TH Aachen, Dept Biol Microbiol & Genet, 4 Worringer Weg, DE-52056 Aachen, Germany. A novel yeast-based tool to detect mutagenic and recombinogenic effects simultaneously. *Mutat Res* **652**, (1) 20
- Scharnawski M, Pongdontri P, Mora G, Hoppert M, Fulda M*. (2008). *Univ Gottingen, Dept Plant Biochem, Justus von Liebig Weg 11, DE-37077 Gottingen, Germany. Mutants of *Saccharomyces cerevisiae* deficient in acyl-CoA synthetases secrete fatty acids due to interrupted fatty acid recycling. *FEBS J* **275**, (11) 2765

- Slagowski NL, Kramer RW, Morrison MF, LaBaer J, Lesser CF*. (2008). *Harvard Med Sch, MGH, Div Infect Dis, Cambridge, Ma 02138, USA. A functional genomic yeast screen to identify pathogenic bacterial proteins. *PLoS Pathog* **4**, (1) 9
- Titu D, Chadha A*. (2008). *IIT, Dept Biotechnol, Bioorgan Chem Lab, IN-600036 Chennai, India. Preparation of optically pure alkyl 3-(hetero-2-yl)-3-hydroxypropionate by *Candida parapsilosis* ATCC 7330 mediated deracemisation. *J Mol Catal B Enzym* **52-53**, 168
- Tondee T, Sirianuntapiboon S*, Ohmomo S. (2008). *King Mongkut's Univ Technol Thonburi, Dept Environm Technol, TH-10140 Bangkok, Thailand. Decolorization of molasses wastewater by yeast strain, *Issatchenkia orientalis* No. SF9-246. *Bioresource Technol* **99**, (13) 5511
- Torrens J, Urpi P, Riu-Aurnatell M, Vichi S*, Lopez-Tamames E, Buxaderas S. (2008). *Univ Barcelona, Dept Nutr & Bromatol, XaRTA, Av Joan XXIII s/n, ES-08028 Barcelona, Spain. Different commercial yeast strains affecting the volatile and sensory profile of cava base wine. *Int J Food Microbiol* **124**, (1) 48
- Vermeulen A, Dang TDT, Geeraerd AH, Bernaerts K, Debevere J, Van Impe J, Devlieghere F*. (2008). *Univ Ghent, Dept Food Safety & Food Qual, Lab Food Microbiol & Food Preservation, BE-9000 Ghent, Belgium. Modelling the unexpected effect of acetic and lactic acid in combination with pH and a_w on the growth/no growth interface of *Zygosaccharomyces bailii*. *Int J Food Microbiol* **124**, (1) 79
- Wang L, Chi ZM*, Wang XH, Ju L, Chi Z, Guo N. (2008). *Ocean Univ China, UNESCO Chinese Ctr Marine Biotechnol, Yushan Rd 5, CN-266003 Qingdao, Peoples Rep China. Isolation and characterization of *Candida membranifaciens* subsp *flavinogenie* W14-3, a novel riboflavin-producing marine yeast. *Microbiol Res* **163**, (3) 255
- Watanabe K, Fujimoto J, Sasamoto M, Dugersuren J, Tumursuh T, Demberel S. (2008). Yakult Cent Inst, Microbiol Res, 1796 Yaho, Tokyo 186 8650, Japan. Diversity of lactic acid bacteria and yeasts in Airag and Tarag, traditional fermented milk products of Mongolia. *World J Microbiol Biotechnol* **24**, (8) 1313
- Wenger MD, DePhillips P, Braceivell DG. (2008). UCL, Dept Biochem Engr, Advanced Ctr Biochem Engr, Torrington Pl, London WC1E 7JE, England. A microscale yeast cell disruption technique for integrated process development strategies. *Biotechnol Prog* **24**, (3) 606
- Wentz AE, Shusta EV*. (2008). *Univ Wisconsin, Dept Chem & Biol Engr, Madison, WI 53706, USA. Enhanced secretion of heterologous proteins from yeast by overexpression of ribosomal subunit RPP0. *Biotechnol Prog* **24**, (3) 748
- Wu TK, Wen HY, Chang CH, Liu YT. (2008). Natl Chiao Tung Univ, Inst Mol Sci, Dept Biol Sci & Technol, Hsin-Chu 300, Taiwan. Protein plasticity: A single amino acid substitution in the *Saccharomyces cerevisiae* oxidosqualene-lanosterol cyclase generates protosta-13(17),24-dien-3 β -ol, a rearrangement product. *Org Lett* **10**, (12) 2529
- Xue FY, Miao JX, Zhang X, Luo H, Tan TW*. (2008). *Beijing Univ Chem Technol, Beijing Key Lab Bioprocess, CN-100029 Beijing, Peoples Rep China. Studies on lipid production by *Rhodotorula glutinis* fermentation using monosodium glutamate wastewater as culture medium. *Bioresource Technol* **99**, (13) 5923
- Yoshimoto M, Sato M, Yoshimoto N, Nakao K. (2008). Yamaguchi Univ, Dept Appl Mol Biosci, 2-16-1 Tokiwadai, Ube, Yamaguchi 755 86, Japan. Liposomal encapsulation of yeast alcohol dehydrogenase with cofactor for stabilization of the enzyme structure and activity. *Biotechnol Prog* **24**, (3) 576
- Zhao X, Kong XL, Hua YY, Feng B, Zhao ZB*. (2008). *CAS, Dalian Inst Chem Phys, Div Biotechnol, CN-116023 Dalian, Peoples Rep China. Medium optimization for lipid production through co-fermentation of glucose and xylose by the oleaginous yeast *Lipomyces starkeyi*. *Eur J Lipid Sci Technol* **110**, (5) 405
- Altman K, Frank M, Neumann D, Jakobs S, Westermann B*. (2008). *Univ Bayreuth, Inst Zellbiol, POB 101251, DE-95440 Bayreuth, Germany. The class V myosin motor protein, Myo2, plays a major role in mitochondrial motility in *Saccharomyces cerevisiae*. *J Cell Biol* **181**, (1) 119
- Aspuria PJ, Tamanoi F*. (2008). *UCLA, Dept Microbiol Immunol & Mol Genet, Jonsson Comprehensive Canc Ctr, Los Angeles, Ca 90095, USA. The Tsc/Rheb signaling pathway controls basic amino acid uptake via the Cat1 permease in fission yeast. *Mol Genet Genomics* **279**, (5) 441
- Bachelier-Bassi S, Gadal O, Bourouta G, Nehrbass U. (2008). Inst Pastuer, CNRS URA 2582, Unite Biol Cellulaire Noyau, 25 rue Dr Roux, FR-75724 Paris, France. Cell cycle-dependent kinetochore localization of condensin complex in *Saccharomyces cerevisiae*. *J Struct Biol* **162**, (2) 248
- Barba G, Soto T, Madrid M, Nunez A, Vicente J, Gacto M*, Cansado J. (2008). *Univ Murcia, Dept Genet & Microbiol, Campus Univ Espinardo, ES-30071 Murcia, Spain. Activation of the cell integrity pathway is channelled through diverse signalling elements in fission yeast. *Cell Signal* **20**, (4) 748
- Caudron F, Andrieux A, Job D, Boscheron C*. (2008). *INSERM U836, Grp Physiopathol Cytoskeleton, FR-38043 Grenoble, France. A new role for kinesin-directed transport of Bik1p (CLIP-170) in *Saccharomyces cerevisiae*. *J Cell Sci* **121**, (9) 1506
- Clemençon B, Rey M, Dianoux AC, Trezeguet R, Lauquin GJM, Brandolin G, Pelosi L*. (2008). *UMR 5092 CNRS-CEA-UJF, FR-38054 Grenoble 9, France. Structure-function relationships of the C-terminal end of the *Saccharomyces cerevisiae* ADP/ATP carrier isoform 2. *J Biol Chem* **283**, (17) 11218
- Clifford DM, Wolfe BA, Roberts-Galbraith RH, McDonald WH, Yates JR, Gould KL*. (2008). *Vanderbilt Univ Sch Med, Howard Hughes Med Inst, 221 Kirkland Hall, Nashville, TN 37232, USA. The Clp1/Cdc14 phosphatase contributes to the robustness of cytokinesis by association with anillin-related Mid1. *J Cell Biol* **181**, (1) 79
- Codlin S, Haines RL, Mole SE*. (2008). *UCL, MRC Mol Cell Biol Lab, Mortimer St, London WC1E 6BT, England. *btn1* affects endocytosis, polarization of sterol-rich membrane domains and polarized growth in *Schizosaccharomyces pombe*. *Traffic* **9**, (6) 936
- Dmitruk KV, Stnutok OV, Gonchar MV, Sibiryi AA*. (2008). *Natl Acad Sci Ukraine, Inst Cell Biol, ul Drahomanova 14-16, UA-79005 Lvov, Ukraine. Construction of flavocytochrome b_5 -overproducing strains of the thermotolerant methylotrophic yeast *Hansenula polymorpha* (*Pichia angusta*). *Microbiology (Mosc)* **77**, (2) 181
- Fox JE, Burow ME, McLachlan JA, Miller CA*. (2008). *Tilane Univ, Ctr Bioenvironm Res, New Orleans, La 70112, USA. Detecting ligands and dissecting nuclear receptor-signaling pathways using recombinant strains of the yeast *Saccharomyces cerevisiae*. *Nat Protoc* **3**, (4) 637
- Hartmann C, Chami M, Zachariae U, De Groot BL, Engel A*, Grutter MG. (2008). *Univ Basel, ME Inst Struct Biol, Biozentrum, CH-4056 Basel, Switzerland. Vacuolar protein sorting: Two different functional states of the AAA-ATPase Vps4p. *J Mol Biol* **377**, (2) 352
- Iwaki T, Iefuji H, Hiraga Y, Hosomi A, Morita T, Giga-Hama Y, Takegawa K*. (2008). *Kagawa Univ, Dept Life Sci, Miki cho, Kagawa 761 0795, Japan. Multiple functions of ergosterol in the fission yeast *Schizosaccharomyces pombe*. *Microbiology* **154**, (3) 830
- Johnson BS, McCaffery JM, Lindquist S*, Gitler AD. (2008). *Whitehead Inst Biomed Res, 9 Cambridge Ctr, Cambridge, Ma 02142, USA. A yeast TDP-43 proteinopathy model: Exploring the molecular determinants of TDR-43 aggregation and cellular toxicity. *Proc Natl Acad Sci U S A* **105**, (17) 6439
- Kim JM, Lee CD, Rath A, Davidson AR*. (2008). *Univ Toronto, Dept Mol Genet, 100 College St, Toronto, Ontario, Canada M5S 1A8. Recognition of non-canonical peptides by the yeast Fus1p SH3 domain: Elucidation of a common mechanism for diverse SH3 domain specificities. *J Mol Biol* **377**, (3) 889
- Kraft C, Deplazes A, Sohrmann M, Peter M. (2008). ETH Honggerberg, Inst Biochem, HPM, CH-8093 Zurich, Switzerland. Mature ribosomes are selectively degraded upon starvation by an autophagy pathway requiring the Ubp3p/Bre5p ubiquitin protease. *Nat Cell Biol* **10**, (5) 602

5 Cell biology

- Aguilar-Arnal L, Marsellach FX, Azorin F*. (2008). *Inst Mol Biol Barcelona - CSIC, Parc Cientif Barcelona, Josep Samitier 1-5,### The fission yeast homologue of CENP-B, Abp1, regulates directionality of mating-type switching. *EMBO J* **27**, (7) 1029

- Lange A, Mills RE, Devine SE, Corbett AH*. (2008). *Emory Univ, Dept Biochem, 1510 Clifton Rd NE, Atlanta, Ga 30322, USA. A PY-NLS nuclear targeting signal is required for nuclear localization and function of the *Saccharomyces cerevisiae* mRNA-binding protein Hrp1. *J Biol Chem* **283**, (19) 12926
- Li F, Palecek SP*. (2008). *Univ Wisconsin, Dept Chem & Biol Engr, Madison, WI 53706, USA. Distinct domains of the *Candida albicans* adhesin Eap1p mediate cell-cell and cell-substrate interactions. *Microbiology* **154**, (4) 1193
- Maslyk M, Kochanowicz E, Zielinski R, Kubinski K, Hellman U, Szyszka R*. (2008). *John Paul II Catholic Univ, Environm Protect Inst, Dept Mol Biol, Krasnicka Ave 102, PL-20718 Lublin, Poland. Yeast surviving factor Svf1 as a new interacting partner, regulator and *in vitro* substrate of protein kinase CK2. *Mol Cell Biochem* **312**, (1-2) 61
- Motley AM, Ward GP, Hettema EH*. (2008). *Univ Sheffield, Dept Mol Biol & Biotechnol, Sheffield S10 2TN, England. Dnm1p-dependent peroxisome fission requires Caf4p, Mdv1p and Fis1p. *J Cell Sci* **121**, (10) 1633
- Nagotu S, Saraya R, Otzen M, Veenhuis M, Van der Klei IJ*. (2008). *Univ Groningen, Mol Cell Biol Groningen Biomol Sci & Biotechnol Inst, POB 14, NL-9750 AA Haren, The Netherlands. Peroxisome proliferation in *Hansenula polymorpha* requires Dnm1p which mediates fission but not *de novo* formation. *Biochim Biophys Acta* **1783**, (5) 760
- Ohtaka A, Okuzaki D, Nojima H*. (2008). *Osaka Univ, Microbial Dis Res Inst, Dept Mol Genet, 3-1 Yamadaoka, Suita, Osaka 565 0871, Japan. Mug27 is a meiosis-specific protein kinase that functions in fission yeast meiosis II and sporulation. *J Cell Sci* **121**, (9) 1547
- Oyamada H, Ogawa Y, Shibata N, Okawa Y, Suzuki S, Kobayashi H*. (2008). *Nagasaki Int Univ, Dept Microbiol, 2825-7 Huis Ten Bosch, Nagasaki, Yamanashi 859 32, Japan. Structural analysis of cell wall mannann of *Candida sojae*, a new yeast species isolated from defatted soybean flakes. *Arch Microbiol* **189**, (5) 483
- Sari F, Heinrich M, Meyeri W, Braus GH, Irniger S*. (2008). *Univ Gottingen, Inst Microbiol & Genet, Griesbachstr 8, DE-37077 Gottingen, Germany. The C-terminal region of the meiosis-specific protein kinase Ime2 mediates protein instability and is required for normal spore formation in budding yeast. *J Mol Biol* **378**, (1) 31
- Shimon L, Hynes GM, McCormack EA, Willison KR*, Horovitz A. (2008). *Inst Canc Res, Chester Beatty Labs, CRUK Ctr Cell & Mol Biol, London SW3 6JB, England. ATP-induced allostery in the eukaryotic chaperonin CCT is abolished by the mutation G345D in CCT4 that renders yeast temperature-sensitive for growth. *J Mol Biol* **377**, (2) 469
- Steffen KK, MacKay VI, Kerr EO, Tsuchiya M, Hu D, Fox LA, Dang N, Johnston ED, Oakes JA, Tchao BN, Pak DN, Fields S, Kennedy BK*, Kaerberlein M. (2008). *Univ Washington, Dept Biochem, Seattle, WA 98195, USA. Yeast life span extension by depletion of 60S ribosomal subunits is mediated by Gcn4. *Cell* **133**, (2) 292
- Stuart D. (2008). Univ Alberta, Dept Biochem, 561 Med Sci Bldg, Edmonton, Alberta, Canada T6G 2H7. The meiotic differentiation program uncouples S-phase from cell size control in *Saccharomyces cerevisiae*. *Cell Cycle* **7**, (6) 777
- Tarasov K, Messier V, Landry CR, Radinovic S, Molina MMS, Shames I, Malitskaya Y, Vogel J, Bussey H, Michnick SW*. (2008). *Univ Montreal, Dept Biochim, Casier Postal 6128, Succursale Ctr-ville, Montreal, Quebec, Canada H3C 3J7. An *in vivo* map of the yeast protein interactome. *Science* **320**, (5882) 1465
- Toret CP, Lee L, Sekiya-Kawasaki M, Drubin DG*. (2008). *Univ Calif Berkeley, Dept Mol & Cell Biol, 16 Barker Hall, Berkeley, Ca 94720, USA. Multiple pathways regulate endocytic coat disassembly in *Saccharomyces cerevisiae* for optimal downstream trafficking. *Traffic* **9**, (5) 848
- Van Heusden GPH, Steensma HY. (2008). Leiden Univ, Inst Biol, Sect Mol & Dev Genet, NL-2333 AL Leiden, The Netherlands. The *Saccharomyces cerevisiae* Wss1 protein is only present in mother cells. *FEMS Microbiol Lett* **282**, (1) 100
- Wang T, Craig EA*. (2008). *Univ Wisconsin, Dept Biochem, 433 Babcock Dr, Madison, WI 53706, USA. Binding of yeast frataxin to the scaffold for Fe-S cluster biogenesis. *Isu. J Biol Chem* **283**, (18) 12674
- Wurtz C, Schliebs W, Erdmann R, Rottensteiner H*. (2008). *Ruhr Univ Bochum, Inst Physiol Chem, Abt Syst Biochem, DE-44780 Bochum, Germany. Dynamin-like protein-dependent formation of Woronin bodies in *Saccharomyces cerevisiae* upon heterologous expression of a single protein. *FEBS J* **275**, (11) 2932

6 Gene expression

- Abascal F, Carmona-Saez P, Carazo JM, Pascual-Montano A*. (2008). *Univ Complutense Madrid, Fac CC Fisicas, Computer Architecture Dept, ES-28040 Madrid, Spain. ChIPCodis: Mining complex regulatory systems in yeast by concurrent enrichment analysis of chip-on-chip data. *Bioinformatics* **24**, (9) 1208
- Brown KM, Landry CR, Hartl DL, Cavalieri D. (2008). Harvard Univ, Dept Organism & Evolutionary Biol, 16 Divinity Ave, Cambridge, Ma 02138, USA. Cascading transcriptional effects of a naturally occurring frameshift mutation in *Saccharomyces cerevisiae*. *Mol Ecol* **17**, (12) 2985
- Cholet O, Henaut A, Hebert A, Bonnarme P*. (2008). *AgroParisTech, INRA, UMR Genie & Microbiol Precedes Alimentaires 782, FR-78850 Thiverval Grignon, France. Transcriptional analysis of L-methionine catabolism in the cheese-ripening yeast *Yarrowia lipolytica* in relation to volatile sulfur compound biosynthesis. *Appl Environ Microbiol* **74**, (11) 3356
- Contreras-Levicoy J, Urbina F, Maldonado E*. (2008). *Univ Chile, Inst Ciencias Biomed, Programa Biol Celular & Mol, Casilla 70086, Santiago 7, Chile. *Schizosaccharomyces pombe* positive cofactor 4 stimulates basal transcription from TATA-containing and TATA-less promoters through Mediator and transcription factor IIA. *FEBS J* **275**, (11) 2873
- Decatur WA, Schnare MN. (2008). Univ Massachusetts, Dept Biochem & Mol Biol, 903 Lederle Grad Res Tower, Amherst, Ma 01003, USA. Different mechanisms for pseudouridine formation in yeast 5S and 5.8S rRNAs. *Mol Cell Biol* **28**, (10) 3089
- El Hage A, Koper M, Kufel J, Tollervey D*. (2008). *Univ Edinburgh, Wellcome Trust Ctr Cell Biol, Edinburgh EH9 3JR, Scotland. Efficient termination of transcription by RNA polymerase I requires the 5' exonuclease Rat1 in yeast. *Genes Dev* **22**, (8) 1069
- Gobeil LA, Plante P, Rohani M, Ouellette M, Provost P*. (2008). *Universite Laval, Fac Med, Ctr Rech CHUL-CHUQ, Ctr Rech Rhumatol & Immunol, 2705 Blvd Laurier, Quebec City, Quebec, Canada G1V 4G2. Involvement of Dcr1 in post-transcriptional regulation of gene expression in *Schizosaccharomyces pombe*. *Front Biosci* **13**, 2203
- Gradolatto A, Rogers RS, Lavender H, Taverna SD, Allis CD, Aitchison JD, Tackett AJ*. (2008). *Univ Arkansas Med Sci, Dept Biochem & Mol Biol, 4301 West Markham St, Little Rock, Ar 72205, USA. *Saccharomyces cerevisiae* Yta7 regulates histone gene expression. *Genetics* **179**, (1) 291
- Hoffmann B, Nickel J, Speer F, Schafer B*. (2008). *Rhein Westfal TH Aachen, Dept Biol Microbiol & Genet, 4 Worringer Weg, DE-52056 Aachen, Germany. The 3'-ends of mature transcripts are generated by a processosome complex in fission yeast mitochondria. *J Mol Biol* **377**, (4) 1024
- Jensen MM, Christensen MS, Bonven B*, Jensen TH. (2008). *Aarhus University, Department Molecular Biology, Centre MRNP Biogenesis & Metab, CF Mollers Alle, DK-8000 Aarhus C, Denmark. Requirements for chromatin reassembly during transcriptional downregulation of a heat shock gene in *Saccharomyces cerevisiae*. *FEBS J* **275**, (11) 2956
- Kawauchi J, Mischo H, Braglia P, Rondon A, Proudfoot NJ*. (2008). *Univ Oxford, Sir William Dunn Sch Pathol, Sth Parks Rd, Oxford OX1 3RE, England. Budding yeast RNA polymerases I and II employ parallel mechanisms of transcriptional termination. *Genes Dev* **22**, (8) 1082
- Kim KY, Truman AW, Levin DE*. (2008). *Johns Hopkins Univ, Dept Biochem & Mol Biol, Baltimore, Md 21205, USA. Yeast Mpk1 mitogen-activated protein kinase activates transcription through Swi4/Swi6 by a noncatalytic mechanism that requires upstream signal. *Mol Cell Biol* **28**, (8) 2579

- Linder T, Rasmussen NN, Samuelson CO, Chatzidakis E, Baraznenok V, Beve J, Henriksen P, Gustafsson CM, Holmberg S*. (2008). *Univ Copenhagen, Dept Mol Biol, Copenhagen Bioctr, Ole Maloes Vej 5, DK-2200 Copenhagen N, Denmark. Two conserved modules of *Schizosaccharomyces pombe* Mediator regulate distinct cellular pathways. *Nucleic Acids Res* **36**, (8) 2489
- Lotito L, Russo A, Chillemi G, Bueno S, Cavalieri D, Capranico G*. (2008). *Univ Bologna, Dept Biochem G Moruzzi, Via Imerio 48, IT-40126 Bologna, Italy. Global transcription regulation by DNA topoisomerase I in exponentially growing *Saccharomyces cerevisiae* cells: Activation of telomere-proximal genes by *TOP1* deletion. *J Mol Biol* **377**, (2) 311
- Manukyan A, Zhang J, Thippeswamy U, Yang JY, Zavala N, Mudannayake MP, Asmussen M, Schneider C, Schneider BL*. (2008). *Texas Tech Univ HSC, Dept Biochem & Cell Biol, Lubbock, Tx 79430, USA. Ccr4 alters cell size in yeast by modulating the timing of *CLN1* and *CLN2* expression. *Genetics* **179**, (1) 345
- Merz K, Hondele M, Goetze H, Gmelch K, Stoeckl U, Griesenbeck J*. (2008). *Univ Regensburg, Inst Biochem Genet & Mikrobiol, DE-93053 Regensburg, Germany. Actively transcribed rRNA genes in *S. cerevisiae* are organized in a specialized chromatin associated with the high-mobility group protein Hmo1 and are largely devoid of histone molecules. *Genes Dev* **22**, (9) 1190
- Mohammed S, Lorenzen K, Kerkhoven R, Van Breukelen B, Vannini A, Cramer P, Heck AJR*. (2008). *Univ Utrecht, Biomol Mass Spectrometry & Proteomics Grp, Bijvoet Ctr Biomol Res, Sorbonnelaan 16, NL-3584 CA Utrecht, The Netherlands. Multiplexed proteomics mapping of yeast RNA polymerase II and III allows near-complete sequence coverage and reveals several novel phosphorylation sites. *Anal Chem* **80**, (10) 3584
- Nagalakshmi U, Wang Z, Waern K, Shou C, Raha D, Gerstein M, Snyder M*. (2008). *Yale Univ, Dept Mol Cellular & Dev Biol, New Haven, Ct 06520, USA. The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science* **320**, (5881) 1344
- Nguyen VQ, Sil A*. (2008). *UCSF, Dept Microbiol & Immunol, San Francisco, Ca 94143, USA. Temperature-induced switch to the pathogenic yeast form of *Histoplasma capsulatum* requires Ryp1, a conserved transcriptional regulator. *Proc Natl Acad Sci U S A* **105**, (12) 4880
- Peng WM, Togawa C, Zhang KL, Kurdastani SK*. (2008). *UCLA, Dept Biol Chem, Los Angeles, Ca 90095, USA. Regulators of cellular levels of histone acetylation in *Saccharomyces cerevisiae*. *Genetics* **179**, (1) 277
- Ran F, Bali M, Michels CA*. (2008). *CUNY Queens Coll, Dept Biol, 65-30 Kissena Blvd, Flushing, NY 11367, USA. Hsp90/Hsp70 chaperone machine regulation of the *Saccharomyces MAL*-activator as determined *in vivo* using noninducible and constitutive mutant alleles. *Genetics* **179**, (1) 331
- Safi A, Wallace KA, Rusche LN*. (2008). *Duke Univ, Inst Genome Sci & Policy, Box 3382, Durham, NC 27710, USA. Evolution of new function through a single amino acid change in the yeast repressor Sum1p. *Mol Cell Biol* **28**, (8) 2567
- Sbia M, Parnell EJ, Yu Y, Olsen AE, Kretschmann KL, Voth WP, Stillman DJ*. (2008). *Univ Utah, Dept Pathol, 15 Nth Med Dr East, Salt Lake City, Ut 84112, USA. Regulation of the yeast Ace2 transcription factor during the cell cycle. *J Biol Chem* **283**, (17) 11135
- Sheth CC, Mogensen EG, Fu MS, Blomfield IC, Muhlschlegel FA*. (2008). *Univ Kent, Dept Biosci, Biomed Sci Grp, Canterbury CT2 7NJ, England. *Candida albicans* HSP12 is co-regulated by physiological CO₂ and pH. *Fungal Genet Biol* **45**, (7) 1075
- Su NY, Ouni I, Papagiannis CV, Kaiser P*. (2008). *240D Med Sci I, Irvine, Ca 92697, USA. A dominant suppressor mutation of the *met30* cell cycle defect suggests regulation of the *Saccharomyces cerevisiae* Met4-Cbf1 transcription complex by Met32. *J Biol Chem* **283**, (17) 11615
- Theodoro RC, De Moraes Gimenes Bosco S, Araujo JP, Candeias JMG, Da Graca Macoris SA, Trinca LA, Bagagli E*. (2008). *UNESP, IBB, Dept Microbiol & Immunol, BR-18618-000 Botucatu, SP, Brazil. Dimorphism, thermal tolerance, virulence and heat shock protein 70 transcription in different isolates of *Paracoccidioides brasiliensis*. *Mycopathologia* **165**, (6) 355
- Triana JL, Mendez Y, Navarro L, Reggio R, Ferreras AC, Edmundo C, Triana-Alonso FJ, Herrera F*. (2008). *Apartado Postal 2351, Maracay, Aragua, Venezuela. Prior interaction of ATP with yeast mRNAs enhances protein synthesis at the initiation step. *Biochim Biophys Acta* **1779**, (3) 175
- Viphakone N, Voisinnet-Hakil F, Minvielle-Sebastia L*. (2008). *Univ Victor Segalen Bordeaux 2, Inst Biochim & Genet Cellulaires, CNRS, Bordeaux, France. Molecular dissection of mRNA poly(A) tail length control in yeast. *Nucleic Acids Res* **36**, (7) 2418
- Zhu R, Janetzko F, Zhang Y, Van Duin ACT, Goddard WA, Salahub DR*. (2008). *Univ Calgary, Inst Biocomplex & Bioinformatic, 2500 University Dr, Calgary, Alberta, Canada T2N 1N4. Characterization of the active site of yeast RNA polymerase II by DFT and ReaxFF calculations. *Theor Chem Acc* **120**, (4-6) 479

7 Genetics & genomics

- Adachi Y, Kokubu A, Ebe M, Nagao K, Yanagida M*. (2008). *Kyoto Univ, Grad Sch Biostudies, Japan Sci & Technol Corp, CREST Res Program, Sakyo ku, Kyoto 606 8501, Japan. Cut1/separase-dependent roles of multiple phosphorylation of fission yeast cohesion subunit Rad21 in post-replicative damage repair and mitosis. *Cell Cycle* **7**, (6) 765
- Au WC, Crisp MJ, DeLuca SZ, Rando OJ, Basrai MA*. (2008). *NIH/NCI, Genet Branch Ctr Canc Res, 8901 Wisconsin Ave, NIMC Bldg 8, Rm 5101, Bethesda, Md 20889, USA. Altered dosage and mislocalization of histone H3 and Cse4p lead to chromosome loss in *Saccharomyces cerevisiae*. *Genetics* **179**, (1) 263
- Batova M, Borecka-Melkusova S, Simockova M, Dzugasova V, Goffa E, Subik J*. (2008). *Comenius Univ, Dept Microbiol & Virol, Mlynska dolina B2, SK-84215 Bratislava 4, Slovakia. Functional characterization of the *CgPGS1* gene reveals a link between mitochondrial phospholipid homeostasis and drug resistance in *Candida glabrata*. *Curr Genet* **53**, (5) 313
- Ben-Aroya S, Coombes C, Kwok T, O'Donnell KA, Boeke JD*, Hieter P. (2008). *Johns Hopkins Univ, Sch Med, High Throughput Biol Ctr, Baltimore, Md 21205, USA. Toward a comprehensive temperature-sensitive mutant repository of the essential genes of *Saccharomyces cerevisiae*. *Mol Cell* **30**, (2) 248
- Botet J, Rodriguez-Mateos M, Ballesta JPG, Revuelta JL, Remacha M*. (2008). *Univ Autonoma Madrid, Ctr Biol Mol, ES-28049 Madrid, Spain. A chemical genomic screen in *Saccharomyces cerevisiae* reveals a role for diphthamidation of translation elongation factor 2 in inhibition of protein synthesis by sordarin. *Antimicrob Agents Chemother* **52**, (5) 1623
- Cai J, Zhao RP, Jiang HF, Wang W*. (2008). *CAS, Kunming Inst Zool, State Key Lab Genet Resources & Evolution, CN-650223 Kunming, Yunnan, Peoples Rep China. *De novo* origination of a new protein-coding gene in *Saccharomyces cerevisiae*. *Genetics* **179**, (1) 487
- Chamilos G, Lewis RE, Lamaris GA, Albert ND, Kontoyiannis DP*. (2008). *UT MD Anderson Canc Ctr, Dept Infectious Dis Infect Control & Employee Hlth, Unit 402, Houston, Tx 77030, USA. Genomewide screening for genes associated with gliotoxin resistance and sensitivity in *Saccharomyces cerevisiae*. *Antimicrob Agents Chemother* **52**, (4) 1325
- Coic E, Feldman T, Landman AS, Haber JE*. (2008). *Brandeis Univ, Dept Biol, Rosenstiel Basic Med Sci Res Ctr, MS029, 415 South St, Waltham, Ma 02454, USA. Mechanisms of Rad52-independent spontaneous and UV-induced mitotic recombination in *Saccharomyces cerevisiae*. *Genetics* **179**, (1) 199
- Croy JE, Fast JL, Grimm NE, Wuttke DS*. (2008). *Univ Colorado, Dept Chem & Biochem, CB 215, Boulder, Co 80309, USA. Deciphering the mechanism of thermodynamic accommodation of telomeric oligonucleotide sequences by the *Schizosaccharomyces pombe* protection of telomeres 1 (Pot1pN) protein. *Biochemistry* **47**, (15) 4345
- Ehmsen KT, Heyer WD*. (2008). *Univ Calif Davis, Microbiol Sect, Davis, Ca 95616, USA. *Saccharomyces cerevisiae* Mus81-Mms4 is a catalytic DNA structure-selective endonuclease. *Nucleic Acids Res* **36**, (7) 2182

- Evans ML, Bostelman LJ, Albrecht AM, Keller AM, Strande NT, Thompson JS*. (2008). *Denison Univ, Dept Biol, Granville, Oh 43023, USA. UV sensitive mutations in histone H3 in *Saccharomyces cerevisiae* that alter specific K79 methylation states genetically act through distinct DNA repair pathways. *Curr Genet* **53**, (5) 259
- Gribun A, Cheung KLY, Huen J, Ortega J, Houry WA*. (2008). *Univ Toronto, Dept Biochem, Toronto, Ontario, Canada M5S 1A8. Yeast Rvb1 and Rvb2 are ATP-dependent DNA helicases that form a heterohexameric complex. *J Mol Biol* **376**, (5) 1320
- Hoffman PD, Curtis MJ, Iwai S, Hays JB*. (2008). *Oregon State Univ, Dept Environm & Mol Toxicol, Corvallis, Or 97331, USA. Biochemical evolution of DNA polymerase η : Properties of plant, human, and yeast proteins. *Biochemistry* **47**, (16) 4583
- Koivistoinen OM, Hilditch S, Voutilainen SP, Boer H, Penttilä M, Richard P*. (2008). *VTT Tech Res Ctr Finland, Tietotie 2, POB 1000, FI-02044 VTT-Espoo, Finland. Identification in the yeast *Pichia stipitis* of the first L-rhamnose-1-dehydrogenase gene. *FEBS J* **275**, (10) 2482
- Malay AD, Umehara T, Matsubara-Malay K, Padmanabhan B*, Yokoyama S. (2008). *Aptuit Laurus Private Ltd, ICICI Knowledge Park, Shameerpet, IN-500078 Hyderabad, India. Crystal structures of fission yeast histone chaperone Asf1 complexed with the Hip1 B-domain or the Cac2 C terminus. *J Biol Chem* **283**, (20) 14022
- Matsuzaki K, Shinohara A, Shinohara M*. (2008). *Osaka Univ, Inst Protein Res, Dept Integrated Protein Funct, 3-2 Yamadaoka, Suita, Osaka 565 0871, Japan. Forkhead-associated domain of yeast Xrs2, a homolog of human Nbs1, promotes nonhomologous end joining through interaction with a ligase IV partner protein, Lif1. *Genetics* **179**, (1) 213
- Miyoshi T, Kanoh J, Saito M, Ishikawa F*. (2008). *Kyoto Univ, Dept Gene Mechanisms, Yoshida Konoe cho, Sakyo ku, Kyoto 606 8501, Japan. Fission yeast Pot1-Tpp1 protects telomeres and regulates telomere length. *Science* **320**, (5881) 1341
- Rodriguez CE, Sobol Z, Schiestl RH*. (2008). *UCLA, Dept Pathol, Los Angeles, Ca 90024, USA. 9,10-Phenanthrenequinone induces DNA deletions and forward mutations via oxidative mechanisms in the yeast *Saccharomyces cerevisiae*. *Toxicol In Vitro* **22**, (2) 296
- Seong C, Sehorn MG, Plate I, Shi I, Song B, Chi P, Mortensen U, Sung P, Krejci L*. (2008). *Masaryk Univ, Natl Ctr Biomol Res, Kamenice 5/A4, CZ-62500 Brno, Czech Republic. Molecular anatomy of the recombination mediator function of *Saccharomyces cerevisiae* Rad52. *J Biol Chem* **283**, (18) 12166
- Solieri L, Antunez O, Perez-Ortin JE, Barrio E, Giudici P. (2008). Univ Modena & Reggio Emilia, Dept Agric & Food Sci, Via Amendola 2, Padiglione Besta, Reggio Emilia, Italy. Mitochondrial inheritance and fermentative : oxidative balance in hybrids between *Saccharomyces cerevisiae* and *Saccharomyces uvarum*. *Yeast* **25**, (7) 485
- Stein A, Aloy P*. (2008). *BSC/IRB, C/ Josep Samitier 1-5, Barcelona, Spain. A molecular interpretation of genetic interactions in yeast. *FEBS Lett* **582**, (8) 1245
- Vernon M, Lobachev K, Petes TD*. (2008). *Duke Univ Sch Med, Dept Mol Genet & Microbiol, Durham, NC 27710, USA. High rates of "unselected" aneuploidy and chromosome rearrangements in *tel1 mecl1* haploid yeast strains. *Genetics* **179**, (1) 237
- Wang Q, Zhang L, Lynn B, Rymond BC*. (2008). *Univ Kentucky, Dept Biol, Lexington, Ky 40506, USA. A BBP-Mud2p heterodimer mediates branchpoint recognition and influences splicing substrate abundance in budding yeast. *Nucleic Acids Res* **36**, (8) 2787
- Zhang M, Liang YP, Zhang XH, Xu Y, Dai HP*, Xiao W. (2008). *CAS, Inst Hydrol, 7 Donghu Rd Sth, CN-430072 Wuhan, Peoples Rep China. Deletion of yeast *CWP* genes enhances cell permeability to genotoxic agents. *Toxicol Sci* **103**, (1) 68
- Braconi D, Possenti S, Laschi M, Geminiani M, Lusini P, Bernardini G, Santucci A*. (2008). *Univ Siena, Dipt Biol Mol, Via Fiorentina 1, IT-53100 Siena, Italy. Oxidative damage mediated by herbicides on yeast cells. *J Agric Food Chem* **56**, (10) 3836
- Chen SL, He Q, Greenberg ML*. (2008). *Wayne State Univ, Dept Biol Sci, Detroit, Mi 48202, USA. Loss of tafazzin in yeast leads to increased oxidative stress during respiratory growth. *Mol Microbiol* **68**, (4) 1061
- Dani C, Bonatto D, Salvador M, Pereira MD, Henriques JAP, Eleutherio E*. (2008). *UFRJ, Univ Caxias do Sul, Inst Quim, Rio de Janeiro, Brazil. Antioxidant protection of resveratrol and catechin in *Saccharomyces cerevisiae*. *J Agric Food Chem* **56**, (11) 4268
- Ernst R, Kueppers P, Klein CM, Schwarzmüller T, Kuchler K, Schmitt L*. (2008). *Univ Dusseldorf, Inst Biochem, Universitätsstr 1, DE-40225 Dusseldorf, Germany. A mutation of the H-loop selectively affects rhodamine transport by the yeast multidrug ABC transporter Pdr5. *Proc Natl Acad Sci U S A* **105**, (13) 5069
- Folmer V, Pedroso N, Matias AC, Lopes SCDN, Antunes F, Cyrne L, Marinho HS*. (2008). *Univ Lisbon, Dept Quim & Bioquim, Ctr Quim & Bioquim, PT-1749-016 Lisbon, Portugal. H₂O₂ induces rapid biophysical and permeability changes in the plasma membrane of *Saccharomyces cerevisiae*. *Biochim Biophys Acta* **1778**, (4) 1141
- Golubev WI, Kulakovskaya TV, Shashkov AS, Kulakovskaya EV, Golubev NV. (2008). Russian Acad Sci, Skryabin Inst Biochem & Physiol Microorg, pr Nauki 5, RU-142290 Pushchino, Russia. Antifungal cellobiose lipid secreted by the epiphytic yeast *Pseudozyma graminicola*. *Microbiology (Mosc)* **77**, (2) 171
- Irazusta V, Moreno-Cermeno A, Cabisco E, Ros J*, Tamarit J. (2008). *Univ Lleida, Dept Ciencias Med Basicas, Grp Bioquim Estres Oxidatiu, ES-25008 Lleida, Spain. Major targets of iron-induced protein oxidative damage in frataxin-deficient yeasts are magnesium-binding proteins. *Free Radic Biol Med* **44**, (9) 1712
- Jadhav SR, Sui DX, Garavito RM, Worden RM*. (2008). *Michigan State Univ, Dept Chem Engn & Materials Sci, East Lansing, Mi 48824, USA. Fabrication of highly insulating tethered bilayer lipid membrane using yeast cell membrane fractions for measuring ion channel activity. *J Colloid Interface Sci* **322**, (2) 465
- Jennings ML, Cui J. (2008). Univ Arkansas Med Sci, Dept Physiol & Biophys, Little Rock, Ar 72205, USA. Chloride homeostasis in *Saccharomyces cerevisiae*: High affinity influx, V-ATPase-dependent sequestration, and identification of a candidate Cl-sensor. *J Gen Physiol* **131**, (4) 379
- Jung WH, Sham A, Lian TS, Singh A, Kosman DJ, Kronstad JW*. (2008). *UBC, Dept Microbiol & Immunol, Michael Smith Lab, Vancouver, BC, Canada V5Z 1M9. Iron source preference and regulation of iron uptake in *Cryptococcus neoformans*. *PLoS Pathog* **4**, (2) e45
- Lopez-Mirabal HR, Winther JR*, Kielland-Brandt MC. (2008). *Univ Copenhagen, Dept Biol, Universitetsparken 13, DK-2100 Copenhagen, Denmark. Oxidant resistance in a yeast mutant deficient in the Sit4 phosphatase. *Curr Genet* **53**, (5) 275
- Somolinos M, Garcia D, Condon S, Manas P, Pagan R*. (2008). *Univ Zaragoza, Dept Prod Anim & Ciencia Alimentos, C/ Miguel Servet 177, ES-50013 Zaragoza, Spain. Biosynthetic requirements for the repair of sublethally injured *Saccharomyces cerevisiae* cells after pulsed electric fields. *J Appl Microbiol* **105**, (1) 166
- Stolworthy TS, Korkegian AM, Willmon CL, Ardiani A, Cundiff J, Stoddard BL, Black ME*. (2008). *Washington State Univ, Dept Pharmaceut Sci, Pullman, Wa 99164, USA. Yeast cytosine deaminase mutants with increased thermostability impart sensitivity to 5-fluorocytosine. *J Mol Biol* **377**, (3) 854
- Sun ZJ, Li SY, Lv GJ, Zhu J, Yu WT, Wang W, Ma XJ*. (2008). *CAS, Dalian Inst Chem Phys, Lab Biomed Materials Engn, 457 Zhongshan Rd, CN-116023 Dalian, Peoples Rep China. Metabolic response of different osmo-sensitive *Saccharomyces cerevisiae* to ACA microcapsule. *Enzyme Microb Technol* **42**, (7) 576
- Vaupotic T, Veranic P, Jenoe P, Plemenitas A*. (2008). *Univ Ljubljana, Inst Biochem, Vrazov trg 2, SI-1000 Ljubljana, Slovenia. Mitochondrial mediation of environmental osmolytes discrimination during osmoadaptation in the extremely halotolerant black yeast *Hortaea werneckii*. *Fungal Genet Biol* **45**, (6) 994

8 Physiology

- Bertilsson M, Andersson J, Liden G. (2008). Lund Univ, Dept Chem Engn, SE-22100 Lund, Sweden. Modeling simultaneous glucose and xylose uptake in *Saccharomyces cerevisiae* from kinetics and gene expression of sugar transporters. *Bioprocess Biosyst Eng* **31**, (4) 369

9 Medical mycology

- Al-Dhaheeri RS, Douglas LJ*. (2008). *University Glasgow, Division Infection & Immunity, Joseph Black Building, Glasgow G12 8QQ, Scotland. Absence of amphotericin B-tolerant persister cells in biofilms of some *Candida* species. *Antimicrob Agents Chemother* **52**, (5) 1884
- Bernardo SM, Khalique Z, Kot J, Jones JK, Lee SA*. (2008). *Univ New Mexico HSC, New Mexico Veterans Healthcare Syst, Div Infect Dis, Albuquerque, NM 87108, USA. *Candida albicans* VPS1 contributes to protease secretion, filamentation, and biofilm formation. *Fungal Genet Biol* **45**, (6) 861
- Gaitanis G, Magiatis P*, Stathopoulou K, Bassukas ID, Alexopoulos EC, Velegraki A, Skaltsounis AL. (2008). *Univ Athens, Dept Pharmacognosy & Chem Nat Prod, Athens, Greece. AhR ligands, malassezin, and indolo[3,2-b]carbazole are selectively produced by *Malassezia furfur* strains isolated from seborrheic dermatitis. *J Invest Dermatol* **128**, (7) 1620
- Garcia J, Shea J, Alvarez-Vasquez F, Qureshi A, Luberto C, Voit EO, Del Poeta M*. (2008). *Med Univ Sth Carolina, Dept Biochem & Mol Biol, 173 Ashley Ave, BSB 503, Charleston, SC 29425, USA. Mathematical modeling of pathogenicity of *Cryptococcus neoformans*. *Mol Syst Biol* **4**, 183
- Jiang B, Xu DM, Allocco J, Parish C, Davison J, Veillette K, Sillaots S, Hu WQ, Rodriguez-Suarez R, Trosok S, Zhang L, Li Y, Rakhkhodae F, Ransom T, Martel N, Wang H, Gauvin D, Wiltsie J, Wisniewski D, Salowe S, Kahn JN, Hsu MJ, Giacobbe R, Abruzzo G, Flattery A, Gill C, Youngman P, Wilson K, Bills G, Platas G, Pelaez F, Diez MT, Kauffman S, Becker J, Harris G, Liberator P, Roemer T*. (2008). *Merck Frosst Canada & Co, Ctr Fungal Genet, 225 President Kennedy West, Montreal, Quebec, Canada H2X 3Y8. PAP inhibitor with *in vivo* efficacy identified by *Candida albicans* genetic profiling of natural products. *Chem Biol* **15**, (4) 363
- Kusch H, Engelmann S, Bode R, Albrecht D, Morschhauser J, Hecker M. (2008). Ernst Moritz Arndt Universität, Institut Mikrobiol, Friedrich Ludwig Jahn Strasse 15, DE-17487 Greifswald, Germany. A proteomic view of *Candida albicans* yeast cell metabolism in exponential and stationary growth phases. *Int J Med Microbiol* **298**, (3-4) 291
- Leal AL, Faganello J, Bassanesi MC, Vainstein MH*. (2008). *UFRGS, Ctr Biotechnol, POB 15005, BR-91501-970 Porto Alegre, RS, Brazil. *Cryptococcus* species identification by multiplex PCR. *Med Mycol* **46**, (4) 377
- Lott TJ, Scarborough RT. (2008). CDC, Natl Ctr Zoonotic Vectorborne & Enteric Dis, Atlanta, Ga 30333, USA. Development of a MLST-biased SNP microarray for *Candida albicans*. *Fungal Genet Biol* **45**, (6) 803
- MacKenzie DA, Defernez M, Dunn WB, Brown M, Fuller LJ, De Herrera SRMS, Gunther A, James SA, Eagles J, Philo M, Goodacre R, Roberts IN. (2008). Institute Food Research, National Collection Yeast Cultures, Norwich Research Pk, Colney, Norwich NR4 7UA, England. Relatedness of medically important strains of *Saccharomyces cerevisiae* as revealed by phylogenetics and metabolomics. *Yeast* **25**, (7) 501
- Maidan MM, De Rop L, Relloso M, Diez-Orejas R, Thevelein JM, Van Dijck P*. (2008). *KUL, VIB Department Molecular Microbiol, Kasteelpark Arenberg 31, BE-3001 Heverlee, Belgium. Combined inactivation of the *Candida albicans* *GPR1* and *TPS2* genes results in avirulence in a mouse model for systemic infection. *Infect Immun* **76**, (4) 1686
- Oliver BG, Silver PM, White TC*. (2008). *Seattle Biomedical Research Inst, 307 Westlake Ave Nth, Seattle, Wa 98109, USA. Polyene susceptibility is dependent on nitrogen source in the opportunistic pathogen *Candida albicans*. *J Antimicrob Chemother* **61**, (6) 1302
- Putignani L, Paglia MG, Bordi E, Nebuloso E, Pucillo LP, Visca P*. (2008). *Univ Roma Tre, Dipt Biol, Viale Marconi 446, IT-00146 Rome, Italy. Identification of clinically relevant yeast species by DNA sequence analysis of the D2 variable region of the 25-28S rRNA gene. *Mycoses* **51**, (3) 209
- Reiss E, Lasker BA, Iqbal NJ, James MJ, Arthington-Skaggs BA. (2008). CDC, Mycot Dis Branch, Div Foodborne Bacterial & Mycot Dis, Atlanta, Ga 30333, USA. Molecular epidemiology of *Candida parapsilosis* sepsis from outbreak investigations in neonatal intensive care units. *Infect Genet Evol* **8**, (2) 103
- Rimek D, Fehse B, Gopel P. (2008). Thueringer Landesamt Lebensmittelsicherheit & Verbraucherschutz, Dezernat Med Mikrobiologie, Nordhaeuserstrasse 74, DE-99089 Erfurt, Germany. Evaluation of Mueller-Hinton-agar as a simple medium for the germ tube production of *Candida albicans* and *Candida dubliniensis*. *Mycoses* **51**, (3) 205
- Satishkumar K, Latha SM, Prasanna JS, Dhanalakshmi K, Sarma BJR. (2008). Sri Venkateswara Univ, Dept Veterinary Clin Med, IN-500030 Hyderabad, India. Cultural and electron microscopic studies of *Malassezia pachydermatis*. *Indian Vet J* **85**, (4) 29
- Tongchusak S, Leelayuwat C, Brusci V, Chaiyaroj SC*. (2008). *Mahidol Univ, Dept Microbiol, TH-10400 Bangkok, Thailand. *In silico* prediction and immunological validation of common HLA-DRB1-restricted T cell epitopes of *Candida albicans* secretory aspartyl proteinase 2. *Microbiol Immunol* **52**, (4) 231
- Torres FAG, Vilaca R, De Moraes LMP, Reis VCB, Felipe MSS. (2008). Univ Brasilia, Dept Biol Celular, Mol Biol Lab, BR-70910-900 Brasilia, DF, Brazil. Expression of a kexin-like gene from the human pathogenic fungus *Paracoccidioides brasiliensis* in *Saccharomyces cerevisiae*. *Med Mycol* **46**, (4) 385
- Walker LA, Munro CA, De Bruijn I, Lenardon MD, McKinnon A, Gow NAR. (2008). Univ Aberdeen, Sch Med Sci, Inst Med Sci, Aberdeen, Scotland. Stimulation of chitin synthesis rescues *Candida albicans* from echinocandins. *PLoS Pathog* **4**, (4) 40
- Weber K, Sohr R, Schulz B, Fleischhacker M, Ruhnke M*. (2008). *Charite Univ Med Berlin, Dept Med, Div Hematol & Oncol, Charitepl 1, DE-10117 Berlin, Germany. Secretion of *E,E*-farnesol and biofilm formation in eight different *Candida* species. *Antimicrob Agents Chemother* **52**, (5) 1859
- Zeng X, Xiong CR, Wang Z, Jiang L, Hou XH, Shen J, Zhou M, Chen QM. (2008). Sichuan Univ, Natl Key Lab Stomatol, Chengdu, Peoples Rep China. Genotypic profiles and virulence attributes of *Candida albicans* isolates from patients with oral lichen planus. *APMIS* **116**, (4) 284
- Zomorodain K, Mirhendi H, Tarazooie B, Kordbacheh P, Zeraati H, Nayeri F. (2008). Shiraz Univ Med Sci, Dept Med Mycol & Parasitol, Shiraz, Iran. Molecular analysis of *Malassezia* species isolated from hospitalized neonates. *Pediatr Dermatol* **25**, (3) 312

10 Recombinant DNA technology

- Chen J, Zhou J, Bae W, Sanders CK, Nolan JP, Cai H*. (2008). *Los Alamos Natl Lab, Natl Flow Cytometry Resource, MS M888, Los Alamos, NM 87545, USA. A yEGFP-based reporter system for high-throughput yeast two-hybrid assay by flow cytometry. *Cytometry Part A* **73A**, (4) 312
- Hribar G, Smilovic V, Zupan AL, Gaberc-Porekar V. (2008). Natl Inst Chem, Dept Biosynth & Biotransformation, Hajdrihova 19, SI-1001 Ljubljana, Slovenia. β -Lactamase reporter system for selecting high-producing yeast clones. *Biotechniques* **44**, (4) 477
- Kepler-Ross S, Noffz C, Dean N*. (2008). *SUNY Stony Brook, Dept Biochem & Cell Biol, Stony Brook, NY 11794, USA. A new purple fluorescent color marker for genetic studies in *Saccharomyces cerevisiae* and *Candida albicans*. *Genetics* **179**, (1) 705
- Miyashita S, Shirako Y*. (2008). *Univ Tokyo, Div Biol Resources Dev, Asian Nat Environm Sci Ctr, 1-1-1 Yayoi, Bunkyo ku, Tokyo 113 8657, Japan. Chromosomal integration of a binding domain:bait gene into yeast enhances detection in the two-hybrid system. *J Microbiol Methods* **73**, (2) 179
- Yamagishi K, Sugiyama M, Kaneko Y, Harashima S*. (2008). *Osaka Univ, Dept Biotechnol, 2-1 Suita, Osaka 565 0871, Japan. Conditional chromosome splitting in *Saccharomyces cerevisiae* using the homing endonuclease PI-SceI. *Appl Microbiol Biotechnol* **79**, (4) 699