

X-RAY SATELLITE INTENSITIES IN THE SUDDEN APPROXIMATION

T. ÅBERG

Research Institute for Theoretical Physics, University of Helsinki, Helsinki, Finland

Received 24 February 1968

The relative intensities of the $K\alpha_1\alpha_2$ satellite group $\alpha'\alpha_3\alpha_4$ are calculated for neon-like ions ($9 \leq Z \leq 20$), and compared with experimental intensities resulting from both electron and X-ray excitation.

A recent theory of X-ray satellites is based on the sudden approximation [1,2]. In this note we show that it accounts for relative intensities of $K\alpha$ satellites appearing both in electron and in X-ray excitation. The influence of the K Auger width on the relative intensity is considered. Previously it has been assumed that the Auger transition rates of the K single- and multi-hole states are equal.

We treat the most prominent $K\alpha$ satellite group of neon-like ions in LS coupling. The initial states of the satellite lines are identified as

KL_I and $KL_{II,III}$ double-hole states in accordance with energy calculations [3,4]. The relative intensity is obtained from

$$\frac{I(\alpha'\alpha_3\alpha_4)}{I(\alpha_1\alpha_2)} = k_{2s}W(2s) + \frac{5}{8}k_{2p}W(2p). \quad (1)$$

Here $W(nl)$ is the relative sudden approximation probability [1,2] that K-shell ionization is followed by excitation or ionization of an n_l electron. The correction factor $k_{nl} = \nu_{nl}^4 M_{nl}^2 \Gamma_O^A / \nu_O^4 M_O^2 \Gamma_{nl}^A$ takes into account that the frequency ν and the

Table 1
Relative integrated intensities (in percent) of the $K\alpha_1\alpha_2$ satellite group $\alpha'\alpha_3\alpha_4$.

Z Target		Experiment				Theory Eq.(1)	
		electron excitation		Ref.4 *	X-ray excitation		
		Ref.11	Ref.10			Ref.13	Ref.12
9	NaF	39.5 ± 1.5					42.5
10							26.0
11	NaF	20.3 ± 0.8			21.5 ± 1		18.2
	NaCl		22				
12	Mg	16.1 ± 0.5		12.6	13.8 ± 0.8		13.6
	MgO	15.9 ± 0.5		14.1	15.9 ± 0.4	15 ± 1.5	
13	Al	10.1 ± 0.4		10.7	10.7 ± 0.4		10.5
	Al ₂ O ₃	11.9 ± 0.4		11.9		9.3 ± 1.5	
14	Si	7.5 ± 0.4	11	9.0	7.7 ± 0.4		8.3
	SiO ₂	9.3 ± 0.4		10.0	8.9 ± 0.4		
15	P	6.2 ± 0.4					6.8
16	S, Fes	5.6 ± 0.4	4.8				5.7
17							4.8
18							4.1
19	K ₃ PO ₄		2.6				3.5
20	CaO		2.2				3.1

* Excitation mode not given

dipole matrix element M of the line as well as the Auger width Γ^A of the initial state change when the nl hole is formed besides the K hole. The radiative width Γ^R has been neglected in k_{nl} as $\Gamma^R \ll \Gamma^A$ for atoms with $Z \leq 20$ [5].

The probabilities $W(nl)$ were first calculated by using ground-state orbitals [6] with the atomic number $Z+1$ instead of K-state orbitals with atomic number Z . Only for some ions [2] has it been possible to obtain the probabilities directly by using K-state orbitals [7]. For the remaining ions the final values $W(nl)$ corresponding to K-state orbitals were then found by interpolating the slowly varying reduction factor. The ratio M_{2p}^2/M_0^2 has been calculated in the frozen structure approximation [8]. In the 2s case this ratio has been estimated to be 1.02 times the 2p ratio. Using Archard's relative values [9] of the KLL Auger probabilities $P(nl', nl)$ in LS coupling we have obtained the Auger-width correction from

$$\frac{\Gamma_o^A}{\Gamma_{nl}^A} = \frac{P(2s, 2s) + P(2s, 2p) + P(2p, 2p)}{a_{nl}P(2s, 2s) + b_{nl}P(2s, 2p) + c_{nl}P(2p, 2p)} \quad (2)$$

The constants a_{nl} , b_{nl} and c_{nl} take the nl hole into account ($a_{2s} = 0$, $b_{2s} = \frac{1}{2}$, $c_{2s} = 1$, $a_{2p} = 1$, $b_{2p} = \frac{5}{6}$ and $c_{2p} = \frac{2}{3}$). The ratio (2) which results in a correction of about 30% is rather insensitive to the choice of the relative Auger probabilities and nearly independent of Z .

The experimental relative intensities in table 1 do not reveal any differences between electron and X-ray excitation in accordance with the sudden approximation. The agreement between the measurements and the calculations, resting on the assumption of a free neon core, is good up to $Z = 16$. For heavier atoms exchange effects between the 2p and 3p shell must be taken into account.

1. V.P. Sachenko and V.F. Demekhin, Zh. Eksp. i. Teor. Fiz. 49 (1965) 765; Soviet Phys. JETP 22 (1966) 532.
2. T. Åberg, Phys. Rev. 156 (1967) 35.
3. Z. Horak, Proc. Phys. Soc. (London) A77 (1961) 980.
4. V.F. Demekhin and V.P. Sachenko, in Röntgenspektren und Chemische Bindung (VEB Repprocolor, Leipzig 1966) 58.
5. E.J. Callan, Phys. Rev. 124 (1961) 793.
6. E. Clementi, Tables of atomic functions (Supplement to IBM J. Res. Devel. 9 (1965) 2).
7. P.S. Bagus, Phys. Rev. 139 (1965) A619.
8. Equation (3.3) in ref. 2. Note, however, the misprint: r^2 should be replaced by r^3 .
9. G.D. Archard, private communication to W.N. Asaad (W. Mehlhorn and W.N. Asaad, Z. Physik 191 (1966) 231).
10. L.G. Parrat, Phys. Rev. 50 (1936) 1.
11. The author is indebted to Dr. Fischer for sending him the relative integrated intensities; see also D.W. Fischer and W.L. Baun, Spectrochim. Acta 21 (1965) 443, J. Appl. Phys. 36 (1965) 534, and D.W. Fischer, J. Chem. Phys. 42 (1965) 3814.
12. From photoelectron spectra; M.O. Krause, T.A. Carlson and R.D. Dismukes, to be published.
13. J. Utriainen, M. Linkoaho, E. Rantavuori, T. Åberg and G. Graeffe, to be published.

* * * * *

ERRATA

Influence of lattice defects on the hyperfine interaction: Mn^{2+} in NaCl, K.N. Schrivastava, Physics Letters 26A (1968) 251.

The correct name of the author is K. N. Shrivastava.

p.251, col. 1, line 8

read An interesting variation ... *instead of* An increasing variation...

p.251, col. 2, line 15

read corresponds to $\rho_{Mn} = 0.910$... *instead of* corresponds to $\rho_{Mn} = -0.910$...

In the caption of table 1

read NaCl *instead of* CaCl

The correct references 1, 7 and 13 are

1. J.S. Van Wieringen, Diss. Faraday Soc. 19 (1955) 118.
7. E. Clementi, IBM J. Res. Develop. 9 (1965) 2, and supplement; C. Fraose, Proc. Cambridge Phil. Soc. 53 (1957) 206.
13. M.I. Kornfel'd and Yu.N. Tolparov, Fiz. Tverd. Tela 9 (1967) 2047; Soviet Phys. Solid State 9 (1968) 1607.

* * *