
Perusing the Literature

Barbara S. Hawkins

BRADFORD HILL HONORED

The October–December 1982 issue of *Statistics in Medicine* (Volume 1, Number 4) is dedicated to Sir Austin Bradford Hill, one of the fathers of controlled medical trials. In addition to reminiscences of Hill and discussions of his work and its influence on other investigators, the journal contains a bibliography of his English language publications. Papers included in this special issue that may be of particular interest to readers are the following:

Schoolman HM: The clinician and the statistician. *Stat Med* 1(4):311–316, 1982

Doll R: Clinical trials: Retrospect and prospect. *Stat Med* 1(4):337–344, 1982

Armitage P: The role of randomization in clinical trials. *Stat Med* 1(4):345–352, 1982

Fleiss JL: Multicentre clinical trials: Bradford Hill's contributions and some subsequent developments. *Stat Med* 1(4):353–359, 1982

Holland WW, Breeze E, Swan AV: Clinical trials: Some reflections. *Stat Med* 1(4):361–368, 1982

IMPLICATIONS OF CLINICAL TRIAL FINDINGS

A group of investigators have reported a benefit–cost analysis of application of their findings from a controlled clinical trial. The references for the current paper, the original report of trial findings, and the editorial which accompanied publication of the current paper are as follows:

Shapiro M, Schoenbaum SC, Tager IB, Munoz A, Polk BF: Benefit-cost analysis of antimicrobial prophylaxis in abdominal and vaginal hysterectomy. *JAMA* 249(10):1290–1294, 1983

Polk BF, Tager IB, Shapiro M, Goren-White B, Goldstein P, Schoenbaum SC: Randomised clinical trial of perioperative cefazolin in preventing infection after hysterectomy. *Lancet* 1:437–441, 1980

Scheckler WE: The cost of prevention: Common sense and decision trees. *JAMA* 249(10):1328, 1983

The investigators used a decision tree to determine excess costs of in-hospital and post-hospital morbidity for both treated and untreated patients. The benefit–cost analysis demonstrated that cefazolin, the drug found to be efficacious in preventing such morbidity in the clinical trial, was also cost-effective. The authors commented:

The analysis has projected substantial savings were this prophylaxis regimen to be used universally. For other antimicrobials to be equally cost-effective, either

they must be more efficacious or less expensive . . . The analysis indicates the importance of examining not just efficacy of a prophylactic antibiotic regimen but cost. The new cephalosporins may be quite effective at preventing post-hysterectomy infections, but this analysis indicates that even with substantially greater effectiveness than cefazolin they might not be equally cost-effective. Similarly, ampicillin, which is less expensive than cefazolin, could be equally cost-effective even if it were somewhat less efficacious.

This current report is recommended reading for those interested in evaluating the implications of implementing findings from clinical trials.

DRUG DEVELOPMENT AND EVALUATION

A series of articles that deals with detection of adverse drug reactions has been published recently:

Venning GR: Identification of adverse reactions to new drugs. I: What have been the important adverse reactions since thalidomide? *Br Med J* 286:199–202, 1983

Venning GR: Identification of adverse reactions to new drugs. II—How were 18 important adverse reactions discovered and with what delays? *Br Med J* 286:289–292, 1983

Venning GR: Identification of adverse reactions to new drugs. II (continued): How were 18 important adverse reactions discovered and with what delays? *Br Med J* 286:365–368, 1983

Venning GR: Identification of adverse reactions to new drugs. III: Alerting processes and early warning systems. *Br Med J* 286:458–460, 1983

Venning GR: Identification of adverse reactions to new drugs. IV—Verification of suspected adverse reactions. *Br Med J* 286:544–547, 1983

Venning surveyed ten British physicians and ten heads of drug regulatory authorities for their opinions on the most important adverse drug reactions determined since thalidomide. From the lists provided by the 20 respondents, a master list was compiled. Venning then examined the published reports of adverse reactions to these drugs in order to identify the first, which alerted attention to the adverse effect, and the subsequent reports, which confirmed the initial warning (since the possibility of false alerts makes verification essential). The author reviews the designs of the reported studies and provides criteria for evaluating the evidence provided by them. In the summary Venning comments that the delays in publication of alerting reports are minor in comparison with delays before verification of adverse reactions and action by regulatory agencies.

Two other investigators concerned with adverse drug reactions have described a study of post-marketing drug evaluation:

Borden EK, Lee JG: A methodologic study of post-marketing drug evaluation using a pharmacy-based approach. *J Chronic Dis* 35:803–816, 1982

These investigators were interested in developing a model that could be used for concurrent data collection on large groups of patients. In addition, the model had to be appropriate for both established and new drugs; the population to which the data applied had to be definable in such a way that both study participants and nonparticipants could be characterized; and health

events had to be completely reported, classified, and coded. As the authors noted, "At present, faced with a credible question of drug-induced adversity there are frequently no data from which to judge the magnitude of the problem or investigate for causal associations."

Borden and Lee enrolled patients through participating pharmacists by identifying patients who filled prescriptions for antibacterial agents. Patients were followed for 30 days after the initial prescription using a mail questionnaire and telephone follow-up when necessary. The investigators achieved questionnaire follow-up of 92.9% of the participants. Life/death status was determined for 97.9% of the participants. The authors noted that the approach they used required a sophisticated organizational structure to assure that the investigators from the various disciplines involved worked "in harmony." This reader found the approach described in this article appealing, not as a replacement for clinical trials as the authors suggested, but as a method for evaluating the impact of controlled drug trials on future patients, after trial findings have been reported.

A special supplement to *Neurology*, edited by R.H. Mattson and entitled "The design of clinical studies to assess the efficacy and toxicity of antiepileptic drugs," has been published. The following papers are included:

Smith DB, Escueta AVD, Cramer JA, Mattson RH: Historical perspective on the choice of antiepileptic drugs for the treatment of seizures in adults. *Neurology* 33(suppl 1):2-7, 1983

Escueta AVD, Mattson RH, Smith DB, Cramer JA, Collins JF: Principles in designing clinical trials for antiepileptic drugs. *Neurology* 33(suppl 1):8-13, 1983

Mattson RH, Cramer JA, Escueta AVD, Smith DB, Collins JF, VA Epilepsy Cooperative Study Group (Browne TR, Crill WE, Homan RW, Mayersdorf A, McCutchen CB, McNamara JO, Rosenthal NP, Treiman DM, Wilder BJ, Williamson PD, Young LM): A design for the prospective evaluation of the efficacy and toxicity of antiepileptic drugs in adults. *Neurology* 33(suppl 1):14-25, 1983

Cramer JA, Smith DB, Mattson RH, Escueta AVD, Collins JF, VA Epilepsy Cooperative Study Group (Browne TR, Crill WE, Homan RW, Mayersdorf A, McCutchen CB, McNamara JO, Rosenthal NP, Treiman DM, Wilder BJ, Williamson PD): A method of quantification for the evaluation of antiepileptic drug therapy. *Neurology* 33(suppl 1):26-37, 1983

Finally, a recent report on trends in new drug development may be of interest to some readers:

Wardell WM, May MS, Trimble AG: New drug development by United States pharmaceutical firms, with analyses of trends in the acquisition and origin of drug candidates, 1963-1979. *Clin Pharmacol Ther* 32(4):407-417, 1982

The authors give particular attention to the decline in self-originated new chemical entities brought to the stage of clinical testing by U.S.-owned companies in the period 1975 through 1979.

NOTES TO THE READER

Regular readers of this column may observe that the listing of publications that appears at the end of the current column has been divided into more classifications than previously. The revised classification scheme is intended

to facilitate review of references by readers with specific interests. In particular, reports of findings from controlled trials have been subdivided on the basis of the *intervention strategy under evaluation*. Trials in which more than one type of intervention were evaluated (e.g., a drug regimen and a surgical procedure in addition to the control intervention) are included under "Other Trials." Publications included in the category "Reports from Related Studies" are often reports from investigators involved in a specific trial but deal with questions other than comparison of findings for the randomized intervention groups.

As noted in earlier columns, the identification of articles for citation depends upon this writer's awareness of the published reports. Only a few journals are screened on a regular basis [see *Controlled Clin Trials* 2(4):327-333, 1981]. The references included in this report were derived from screening 246 separate issues of 35 different journals. All readers are not only invited but *encouraged* to send either preprints or reprints of recent publications or a letter or card to provide references for inclusion in future reviews.

Announcements

M.R.C. trial of multivitamin prophylaxis in pregnancies at risk of neural tube defects. *Lancet* 2:1412, 1982

Discussions and Critiques of Specific Trials

Isoniazid prevention of tuberculosis. *Lancet* 1:395-396, 1983

Recruiting for multicentre trials. *Lancet* 1:78, 1983

Senile disciform macular degeneration. *Lancet* 1:278-279, 1983

Treatment of multiple sclerosis. *Lancet* 1:909-910, 1983

The trial of homoeopathy. *Lancet* 1:108, 1983

Adler MW, Belsey EM, McCutchan JA, Mindel A: Should homosexuals be vaccinated against hepatitis B virus? Cost and benefit assessment. *Br Med J* 286:1621-1624, 1983

Allen ED: Laser photocoagulation of senile macular degeneration. *Br Med J* 286:1357-1358, 1983

Angell M: Multiple sclerosis and the Ingelfinger Rule. *N Engl J Med* 308(4):217-218, 1983

Barclay WR: BCG: An effective immunizing agent. *JAMA* 249(17):2376, 1983

Barnes G: A nurse's experience in the MRC's hypertension trial. *Br Med J* 285:1625-1627, 1982

Bernstein C: Alerting SMD victims in time. *Sightsaving* 51(2):16-21, 1982

Bird AC: Laser photocoagulation of senile macular degeneration. *Br Med J* 286:1001, 1983

Burch PRJ, Seltzer CC, Oster KA, Gordon DB, Walker WJ, Bross IDJ, Cutler JA: The Multiple Risk Factor Intervention Trial. *JAMA* 249(11):1435-1437, 1983

Clemens JD, Chuong JJH, Feinstein AR: The BCG controversy. A methodological and statistical reappraisal. *JAMA* 249(17):2362-2369, 1983

Findley LJ, Whelan DM, Moser KM: Long-term oxygen therapy in COPD. *Chest* 83(4):671-674, 1983

Furberg CD, Friedewald WT, Eberlein KA, eds.: *Proceedings of the Workshop on Implications of Recent Beta-Blocker Trials for Post-Myocardial Infarction Patients*. *Circulation* 67(suppl 1):1-111, 1983 (American Heart Association monograph no. 96)

- Galton DJ, Winder AF: Surgical approach to the hyperlipidaemias. *J R Soc Med* 75:837–838, 1982
- Ghosh A, Roe PF, Shipley M, English JM: Homoeopathic treatment of osteoarthritis. *Lancet* 1:304–305, 1983
- Goldman KP, Valenzuela P, Galleguillos F, Empey DW, Festenstein F: Tuberculosis. *Lancet* 1:592–593, 1983
- Gordon RS, Sanders WE: Implications of recent B-blocker clinical trials for patients after myocardial infarction. *JAMA* 249(18):2482–2483, 1983
- Griffiss JM: Vaccination against meningococcal disease. *N Engl J Med* 308(23):1421, 1983
- Gross F, Strasser T, Buhler FR, Doyle AE, Epstein FH, Ganten D, Griffiths A, Hansson L, Julius S, Keil U, Klumbiene J, Labarthe DR, Langford HG, Li-Sheng L, Lund-Johansen PF, Smith WM, Menard J, Meyer P, Miall WE, Mugambi M, Pobee JOM, Reader R, Robertson JIS, Rosenfeld J, Shvacabaja IK, Hart JT, Tyroler A, Vallotton M, Wilhelmsen L, Yamori Y, Zanchetti A: Guidelines for the treatment of mild hypertension: Memorandum from a W.H.O./I.S.H. meeting. *Lancet* 1:457–458, 1983
- Hanckel FS, Duff P, Polk BF, Shapiro M, Munoz A, Tager IB, Schoenbaum SC: Risk factors for infection after hysterectomy. *N Engl J Med* 308(21):1290–1291, 1983
- Hart JT, Barley SL: Secondary prevention of myocardial infarction: An uncertain feeling. *J R Coll Gen Pract* 32:651–653, 1982
- Kaplan NM: Therapy for mild hypertension: Toward a more balanced view. *JAMA* 249(3):365–367, 1983
- Kennedy CO, Couzigou P, Richard-Mollard B, Fleury B, Amouretti A, Beraud C, Jenkins MD, Clover AM: Homoeopathy. *Lancet* 1:482, 1983
- Lorber J: Vitamins to prevent neural tube defects. *Lancet* 2:1458–1459, 1982
- McFarlin DE: Treatment of multiple sclerosis. *N Engl J Med* 308(4):215–217, 1983
- Nixon PGF, Pugh S: Coronary artery bypass surgery. *Lancet* 1:242, 1983
- Oslo Study Research Group, Leren P, Helgeland A, Hjermann I, Holme I: MRFIT and the Oslo Study. *JAMA* 249(7):893–894, 1983
- Stallones RA: Mortality and the Multiple Risk Factor Intervention Trial. *Am J Epidemiol* 117(6):647–650, 1983
- Thompson E, Parker B: Screening for diabetic eye disease—Mississippi. *JAMA* 249(16):2147, 1983
- Turi ZG, Braunwald E: The use of β -blockers after myocardial infarction. *JAMA* 249(18):2512–2516, 1983
- Winegrad AI, Simmons DA, Martin DB: Has one diabetic complication been explained? *N Engl J Med* 308(3):152–154, 1983
- Zumoff B, Weinblatt E, Ruberman W: Mortality after myocardial infarction. *JAMA* 249(7):885, 1983

Methods

- Code of practice for the clinical assessment of licensed medicinal products in general practice. Joint agreement between BMA, RCGP, and ABPI. *Br Med J* 286:1295–1297, 1983
- Altman DG, Gore SM, Gardner MJ, Pocock SJ: Statistical guidelines for contributors to medical journals. *Br Med J* 286:1489–1493, 1983
- Cease KB: Cooperative clinical trials and continuing medical education. *N Engl J Med* 308(4):225, 1983
- Diamond GA, Forrester JS: Clinical trials and statistical verdicts: Probable grounds for appeal. *Ann Intern Med* 98:385–394, 1983

- Feinstein AR, Horwitz RI: Double standards, scientific methods, and epidemiologic research. *N Engl J Med* 307(26):1611–1617, 1982
- Frisancho AR, Flegel PN: Elbow breadth as a measure of frame size for US males and females. *Am J Clin Nutr* 37:311–314, 1983
- Gardner MJ, Altman DG, Jones DR, Machin D: Is the statistical assessment of papers submitted to the “British Medical Journal” effective? *Br Med J* 286:1485–1488, 1983
- Hall JC: A method for the rapid assessment of sample size in dietary studies. *Am J Clin Nutrition* 37:473–477, 1983
- Herson J: Practical side of multistage clinical trials for screening new agents. *Cancer Treat Rep* 67(1):71–75, 1983
- Horn SD: Measuring severity of illness: Comparisons across institutions. *Am J Public Health* 73(1):25–31, 1983
- Howard J, Whittemore AS, Hoover JJ, Panos M, Aspirin Myocardial Infarction Study Research Group: How blind was the patient blind in AMIS? *Clin Pharmacol Ther* 32(5):543–553, 1982
- Kanep VV, Solomonov SL: A “passport” to monitor health status. *World Health Forum* 3(3):317–320, 1983
- Kraus JF, Conley A, Hardy R, Sexton M, Sweezy Z: Relationship of demographic characteristics of interviewers to blood pressure measurements. *J Comm Health* 8(1):3–12, 1982
- Lee YJ: Interim recruitment goals in clinical trials. *J Chron Dis* 36(5):379–389, 1983
- el Lozy M: Dietary variability and its impact on nutritional epidemiology. *J Chron Dis* 36(3):237–249, 1983
- Munro I, Fox R, Sharp D, DerSimonian R, Charette LJ, McPeck B, Mosteller F: Reporting on methods in clinical trials. *N Engl J Med* 308(10):596–597, 1983
- Sriwatanakul K, Kelvie W, Lasagna L: The quantification of pain: An analysis of words used to describe pain and analgesia in clinical trials. *Clin Pharmacol Ther* 32(2):143–148, 1982
- Stuff JE, Garza C, Smith EO, Nichols BL, Montandon CM: A comparison of dietary methods in nutritional studies. *Am J Clin Nutr* 37:300–306, 1983
- WHO Expert Committee: International nonproprietary names for pharmaceutical substances. *WHO Chron* 36(5, suppl):1–28, 1982

General Issues

- Prevention of insulin-dependent diabetes. *Lancet* 1:104–105, 1983
- Some tribulations of antibiotic trials. *Lancet* 1:337, 1983
- Block SH: Substantial evidence of effectiveness. *JAMA* 249(17):2329–2330, 1983
- Brophy JJ: Patients should be responsible for their own medical records. *N Engl J Med* 307:1530, 1982
- Cancer Research Campaign Working Party in Breast Conservation: Informed consent: Ethical, legal, and medical implications for doctors and patients who participate in randomised clinical trials. *Br Med J* 286:1117–1121, 1983
- Capron AM, Ackerman TF, Strong CM, Lipsett MB: Ethics of phase I clinical trials. *JAMA* 249(7):882–883, 1983
- Darsee J: A retraction of two papers on cardiomyopathy. *N Engl J Med* 308(23):1419, 1983
- Earlam R, Pollock AV, Evans M, Welsh J, Freed DLJ, Francis H: Informed consent. *Br Med J* 286:1436–1437, 1983
- Glass RL, Peterson JK, Bixler D: The effects of changing caries prevalence and diagnostic criteria on clinical caries trials. *Caries Res* 17(2):145–151, 1983

- Hall JC, Hill D, Watts JM: Misuse of statistical methods in the Australasian surgical literature. *Aust NZ J Surg* 52(5):541-543, 1982
- Hemminki E: Quality of clinical trials—A concern of three decades. *Meth Inform Med* 21:81-85, 1982
- Holle R: Clinical trials: Design—management—results—consequences. Report on the 26th Annual Meeting of the GMDS. *Meth Inform Med* 21:94-95, 1982
- Hutt PB: Investigations and reports respecting FDA regulation of new drugs (Part I). *Clin Pharmacol Ther* 33(4):537-548, 1983
- Hutt PB: Investigations and reports respecting FDA regulation of new drugs (Part II). *Clin Pharmacol Ther* 33(5):674-687, 1983
- Iliff RD: The significance of statistical significance. *N Engl J Med* 308(10):596, 1983
- Knox RA: Deeper problems for Darsee: Emory probe. *JAMA* 249(21):2867-2876, 1983
- Knox R: The Harvard fraud case: Where does the problem lie? *JAMA* 249(14):1797-1807, 1983
- Lasagna L: Discovering adverse drug reactions. *JAMA* 249(16):2224-2225, 1983
- Lionel W, Herxheimer A: Coherent policies on drugs. *World Health Forum* 3(3):285-296, 1982
- May MS, Wardell WM, Lasagna L: New drug development during and after a period of regulatory change: Clinical research activity of major United States pharmaceutical firms, 1958 to 1979. *Clin Pharmacol Ther* 33(6):691-700, 1983
- Mischelevich DJ: Lessons and challenges from the demise of the Department of Medical Computer Science at the University of Texas Health Science Center at Dallas. *J Med Syst* 7(1):1-9, 1983
- Mishkin B: Dealing with alleged fraud in medical research. *JAMA* 249(9):1150, 1983
- Newell FW: Junior author—senior name. *Am J Ophthalmol* 95(6):833-834, 1983
- Posner HS: Unambiguous reporting of the sex of human beings. *Lancet* 1:175-176, 1983
- Relman AS: Lessons from the Darsee affair. *N Engl J Med* 308(23):1415-1417, 1983
- Rossi AC, Knapp DE, Anello C, O'Neill RT, Graham CF, Mendelis PS, Stanley GR: Discovery of adverse drug reactions. A comparison of selected phase IV studies with spontaneous reporting methods. *JAMA* 249(16):2226-2228, 1983
- Silverman WA: The public character of scientific medicine. *Perspectives Biol Med* 26(3):343-353, 1983
- Singer J, Sacks HS, Lucente F, Chalmers TC: Physician attitudes toward applications of computer data base systems. *JAMA* 249(12):1610-1614, 1983
- Spodick DH, Grylack LJ, Sade RM, Miller MC, Appelbaum PS, Roth LH, Lidz CW, Schafer A: Ethics of the randomized clinical trial. *N Engl J Med* 308(6):343-345, 1983
- Spodick DH, Gould KL: Controlled clinical trials. *JAMA* 249(11): 1434-1435, 1983
- Wikler D, Pattullo EL: Institutional review boards and the freedom to take risks. *N Engl J Med* 308(10):596, 1983

Statistical Methods

- Aitkin M, Laird N, Francis B: A reanalysis of the Stanford heart transplant data. (With discussion) *J Am Stat Assoc* 78:264-292, 1983
- Anbar D: On estimating the difference between two probabilities, with special reference to clinical trials. *Biometrics* 39:257-262, 1983
- Aranda-Ordaz FJ: An extension of the proportional-hazards model for grouped data. *Biometrics* 39:109-117, 1983

- Atkinson AC: Optimum biased coin designs for sequential clinical trials with prognostic factors. *Biometrika* 69(1):61–67, 1982
- Benedetti JK, Liu P-Y, Sather HN, Seinfeld J, Epton MA: Effective sample size for test of censored survival data. *Biometrika* 69(2):343–349, 1982
- Chernick MR, Downing DJ, Pike DH: Detecting outliers in time series data. *J Am Stat Assoc* 77:743–747, 1982
- Cox DR: On the role of data of possibly lowered sensitivity. *Biometrika* 69(1):215–219, 1982
- Dempster AP, Selwyn MR, Weeks BJ: Combining historical and randomized controls for assessing trends in proportions. *J Am Stat Assoc* 78:221–227, 1983
- DuMouchel WH, Harris JE: Bayes methods for combining the results of cancer studies in humans and other species. *J Am Stat Assoc* 78:293–315, 1983
- Gould AL, Pecore VJ: Group sequential methods for clinical trials allowing early acceptance of H_0 and incorporating costs. *Biometrika* 69(1):75–80, 1982
- Hanley JA, Parnes MN: Nonparametric estimation of a multivariate distribution in the presence of censoring. *Biometrics* 39:129–139, 1983
- Harrington DP, Fleming TR: A class of rank test procedures for censored survival data. *Biometrika* 69(3):553–566, 1982
- Hearne EM, Clark GM, Hatch JP: A test for serial correlation in univariate repeated-measures analysis. *Biometrics* 39:237–243, 1983
- Hsieh FY, Crowley J, Tormey DC: Some test statistics for use in multistate survival analysis. *Biometrika* 70(1):111–119, 1983
- Kay R, Schumacher M: Unbiased assessment of treatment effects on disease recurrence and survival in clinical trials. *Statistics Med* 2(1):41–58, 1983
- Laird NM, Ware JH: Random-effects models for longitudinal data. *Biometrics* 38(4):963–974, 1982
- McHugh R, Matts J: Post-stratification in the randomized clinical trial. *Biometrics* 39:217–225, 1983
- Slud E, Wei LJ: Two-sample repeated significance tests based on the modified Wilcoxon statistic. *J Am Stat Assoc* 77:862–868, 1982
- Stewart WH, Pierce DA: Efficiency of Cox's model in estimating regression parameters with grouped survival data. *Biometrika* 69(3):539–545, 1982
- Tibshirani RJ, Ciampi A: A family of proportional- and additive-hazards models for survival data. *Biometrics* 39:141–147, 1983
- Tsiatis AA: Repeated significance testing for a general class of statistics used in censored survival analysis. *J Am Stat Assoc* 77:855–861, 1982
- Victor N: Exploratory data analysis and clinical research. *Meth Inform Med* 21:53–54, 1982
- Wahrendorf J: Routine statistics and clinical trials. *Biometrics* 39:269–273, 1983
- Whitehead J, Stratton I: Group sequential clinical trials with triangular continuation regions. *Biometrics* 39:227–236, 1983
- Willan AR: Relative efficiency in survival analysis. *J Chronic Dis* 36(2):183–192, 1983

Computer Programs and Methods

- Aickin M: A program for balancing the allocation of subjects to treatment in a clinical trial. *Comp Biomed Res* 15:519–524, 1982
- Atkinson EN, Brown BW, Herson J: KSTAGE: An interactive computer program for designing phase II clinical trials using predictive probability. *Comp Biomed Res* 15:220–227, 1982
- Stead WW, Hammond WE, Straube MJ: A chartless record—Is it adequate? *J Med Syst* 7(2):103–109, 1983

Reports of Findings from Drug Trials

- Aberg-Wistedt A: A double-blind study of zimelidine, a serotonin uptake inhibitor, and desipramine, a noradrenaline uptake inhibitor, in endogenous depression. I. Clinical findings. *Acta Psychiatr Scand* 66:50–65, 1982
- Aberg-Wistedt A, Ross SB, Jostell K-G, Sjoquist B: A double-blind study of zimelidine, a serotonin uptake inhibitor, and desipramine, a noradrenaline uptake inhibitor, in endogenous depression. II. Biochemical findings. *Acta Psychiatr Scand* 66:66–82, 1982
- Allen GS, Ahn HS, Preziosi TJ, Battye R, Boone SC, Chou SN, Kelly DL, Weir BK, Crabbe RA, Lavik PJ, Rosenbloom SB, Dorsey FC, Ingram CR, Mellits DE, Bertsch LA, Boisvert DPJ, Hundley MB, Johnson RK, Strom JA, Transou CR: Cerebral arterial spasm—A controlled trial of nimodipine in patients with subarachnoid hemorrhage. *N Engl J Med* 308(11):619–624, 1983
- Anabwani GM, Ngira JA, Dimiti G, Bryceson ADM: Comparison of two dosage schedules of sodium stibogluconate in the treatment of visceral leishmaniasis in Kenya. *Lancet* 1:210–213, 1983
- Anderson JL, Marshall HW, Bray BE, Lutz JR, Frederick PR, Yanowitz FG, Datz FL, Klausner SC, Hagan AD: A randomized trial of intracoronary streptokinase in the treatment of acute myocardial infarction. *N Engl J Med* 308(22):1312–1318, 1983
- Anderson JR, Wilson JF, Jenkin RDT, Meadows AT, Kersey J, Chilcote RR, Coccia P, Exelby P, Kushner J, Siegel S, Hammond D: Childhood non-Hodgkin's lymphoma—The results of a randomized therapeutic trial comparing a 4-drug regimen (COMP) with a 10-drug regimen (LSA₂-L₂). *N Engl J Med* 308(10):559–565, 1983
- Balfour HH, Bean B, Laskin OL, Ambinder RF, Meyers JD, Wade JC, Zaia JA, Aeppli D, Kirk LE, Segreti AC, Keeney RE, and the Burroughs Wellcome Collaborative Acyclovir Study Group: Acyclovir halts progression of herpes zoster in immunocompromised patients. *N Engl J Med* 308(24):1448–1453, 1983
- Bjertnaes A, Block JM, Hafstad PE, Holte M, Ottemo I, Larsen T, Pinder RM, Steffensen K, Stulemeijer SM: A multicentre placebo-controlled trial comparing the efficacy of mianserin and chlordiazepoxide in general practice patients with primary anxiety. *Acta Psychiatr Scand* 66:199–207, 1982
- Cantekin EI, Mandel EM, Bluestone CD, Rockette HE, Paradise JL, Stool SE, Fria TJ, Rogers KD: Lack of efficacy of a decongestant-antihistamine combination for otitis media with effusion ("secretory" otitis media) in children. Results of a double-blind, randomized trial. *N Engl J Med* 308(6):297–301, 1983
- Cavalli F, Beer M, Martz G, Jungi WF, Alberto P, Obrecht JP, Mermillod B, Brunner KW: Concurrent or sequential use of cytotoxic chemotherapy and hormone treatment in advanced breast cancer: Report of the Swiss Group for Clinical Cancer Research. *Br Med J* 286:5–8, 1983
- Cook PJ, Huggett A, Graham-Pole R, Savage IT, James IM: Hypnotic accumulation and hangover in elderly inpatients: A controlled double-blind study of temazepam and nitrazepam. *Br Med J* 286:100–102, 1983
- Coutinho RA, Lelie N, Albrecht-Van Lent P, Reerink-Brongers EE, Stoutjesdijk L, Dees P, Nivard J, Huisman J, Reesink HW: Efficacy of heat inactivated hepatitis B vaccine in male homosexuals: Outcome of a placebo controlled double blind trial. *Br Med J* 286:1305–1308, 1983
- Dahl L-E, Lundin L, Honoré PL, Dencker SJ: Antidepressant effect of femoxetine and desipramine and relationship to the concentration of amine metabolites in cerebrospinal fluid: A double-blind evaluation. *Acta Psychiatr Scand* 66:9–17, 1982
- Del Rio M, Chrane D, Shelton S, McCracken GH, Nelson JD: Ceftriaxone versus ampicillin and chloroamphenicol for treatment of bacterial meningitis in children. *Lancet* 1:1241–1244, 1983
- Douglas JG, Gough J, Preston PG, Frazer I, Haslett C, Chalmers SR, Munro JF: Long-term efficacy of fenfluramine in treatment of obesity. *Lancet* 1:384–386, 1983

- DuPont HL, Galindo E, Evans DG, Cabada FJ, Sullivan P, Evans DJ: Prevention of travelers' diarrhea with trimethoprim-sulfamethoxazole and trimethoprim alone. *Gastroenterology* 84:75-80, 1983
- Einhorn LH, Williams SD, Stevens EE, Bond WH, Chenoweth L: Random prospective study of cyclophosphamide, doxorubicin, and methotrexate (CAM) combination chemotherapy versus single-agent sequential chemotherapy in non-small cell lung cancer. *Cancer Treat Rep* 66(12):2005-2011, 1982
- Elomaa I, Blomqvist C, Grohn P, Porkka L, Kairento A-L, Selander K, Lamberg-Allardt C, Holmstrom T: Long-term controlled trial with diphosphonate in patients with osteolytic bone metastases. *Lancet* 1:146-149, 1983
- Ericsson CD, DuPont HL, Sullivan P, Galindo E, Evans DG, Evans DJ: Bicozamycin, a poorly absorbable antibiotic, effectively treats travelers' diarrhea. *Ann Intern Med* 98:20-25, 1983
- Feighner JP, Meridith CH, Dutt JE, Hendrickson GG: A double blind comparison of lofepramine, imipramine and placebo in patients with primary depression. *Acta Psychiatr Scand* 66:100-108, 1982
- Freeman AI, Weinberg V, Brecher ML, Jones B, Glicksman AS, Sinks LF, Weil M, Pleuss H, Hananian J, Burgert EO, Gilchrist GS, Necheles T, Harris M, Kung F, Patterson RB, Maurer H, Leventhal B, Chevalier L, Forman E, Holland JF: Comparison of intermediate-dose methotrexate with cranial irradiation for the post-induction treatment of acute lymphocytic leukemia in children. *N Engl J Med* 308(9):477-484, 1983
- Freeman LD, Hooper DR, Lathen DF, Nelson DP, Harrison WO, Anderson DS: Brief prophylaxis with doxycycline for the prevention of traveler's diarrhea. *Gastroenterology* 84:276-280, 1983
- Green SB, Byar DP, Walker MD, Pistenmaa DA, Alexander E, Batzdorf U, Brooks WH, Hunt WE, Mealey J, Odom GL, Paoletti P, Ransohoff J, Robertson JT, Selker RG, Shapiro WR, Smith KR, Wilson CB, Strike TA: Comparisons of carmustine, procarbazine, and high-dose methylprednisolone as additions to surgery and radiotherapy for the treatment of malignant glioma. *Cancer Treat Rep* 67(2):121-132, 1983
- Hagen C, Christensen MS, Christiansen C, Stocklund K-E, Transbol I: Effects of two years' estrogen-gestagen replacement on climacteric symptoms and gonadotropins in the early post-menopausal period. *Acta Obstetr Gynecol Scand* 61:237-241, 1982
- Harmer M, Slattery PJ, Rosen M, Vickers MD: Intramuscular on demand analgesia: Double blind controlled trial of pethidine, buprenorphine, morphine, and mepazinol. *Br Med J* 286:680-682, 1983
- Holmes DG, Bogers WAJL, Wideroe T-E, Huunan-Seppala A, Wideroe B: Endralazine, a new peripheral vasodilator: Absence of effect of acetylator status on anti-hypertensive effect—Results of a multicentre clinical study. *Lancet* 1:670-671, 1983
- Isenberg JI, Peterson WL, Elashoff JD, Sandersfeld MA, Reedy TJ, Ippoliti AF, Van Deventer GM, Frankl H, Longstreth GF, Anderson DS: Healing of benign gastric ulcer with low-dose antacid or cimetidine. A double-blind, randomized, placebo-controlled trial. *N Engl J Med* 308(22):1319-1324, 1983
- Javid MJ, Nordby EJ, Ford LT, Hejna WJ, Whisler WW, Burton C, Millett DK, Wiltse LL, Widell EH, Boyd RJ, Newton S-E, Thisted R: Safety and efficacy of chymopapain (Chymodiactin) in herniated nucleus pulposus with sciatica. Results of a randomized, double-blind study. *JAMA* 249(18):2489-2494, 1983
- Judzewitsch RG, Jaspan JB, Polonsky KS, Weinberg CR, Halter JB, Halar E, Pfeifer MA, Vukadinovic C, Bernstein L, Schneider M, Liang K-Y, Gabbay KH, Rubenstein AH, Porte D: Aldose reductase inhibition improves nerve conduction velocity in diabetic patients. *N Engl J Med* 308(3):119-125, 1983

- Knatterud GL, Klimt CR, Goldner MG, Hawkins BS, Weisenfeld S, Kreines K, Haddock L: The University Group Diabetes Program. Effects of hypoglycemic agents on vascular complications in patients with adult-onset diabetes. VIII. Evaluation of insulin therapy: Final report. *Diabetes* 31(suppl 5):1-81, 1982
- Leppik IE, Derivan AT, Homan RW, Walker J, Ramsay RE, Patrick B: Double-blind study of lorazepam and diazepam in status epilepticus. *JAMA* 249(11):1452-1454, 1983
- Lerner AM, Reyes MP, Cone LA, Blair DC, Jansen W, Wright GE, Lorber RR: Randomised, controlled trial of the comparative efficacy, auditory toxicity, and nephrotoxicity of tobramycin and netilmicin. *Lancet* 1:1123-1126, 1983
- Levy R, Little A, Chuaqui P, Reith M: Early results from double blind, placebo controlled trial of high dose phosphatidylcholine in Alzheimer's disease. *Lancet* 1:987-988, 1983
- Lingjaerde O, Bratlid T, Westby OC, Gordeladze JO: Effect of midazolam, flunitrazepam, and placebo against midwinter insomnia in northern Norway. *Acta Psychiatr Scand* 67:118-129, 1983
- Manthorpe T, Justesen T: Metronidazole prophylaxis in abdominal hysterectomy—A double-blind controlled trial. *Acta Obstetr Gynecol Scand* 61:243-246, 1982
- Messinis IE, Nillius SJ: Comparison between tamoxifen and clomiphene for induction of ovulation. *Acta Obstetr Gynecol Scand* 61:377-379, 1982
- Mills KB: Blind randomised non-crossover long-term trial comparing topical timolol 0.25% with timolol 0.5% in the treatment of simple chronic glaucoma. *Br J Ophthalmol* 67:216-219, 1983
- Nolvadex Adjuvant Trial Organisation: Controlled trial of tamoxifen as adjuvant agent in management of early breast cancer—Interim analysis at four years. *Lancet* 1:257-261, 1983
- Palmblad J, Lonnqvist B: Combination of amikacin and either ampicillin or cephalotin as initial treatment of febrile neutropenic patients. *Acta Med Scand* 212:379-384, 1982
- Papadakis J, Brown CB, Cameron JS, Adu D, Bewick M, Donaghey R, Ogg CS, Rudge C, Williams DG, Taube D: High versus "low" dose corticosteroids in recipients of cadaveric kidneys: Prospective controlled trial. *Br Med J* 286:1097-1100, 1983
- Partamian LG, Kass MA, Gordon M: A dose-response study of the effect of levobunolol on ocular hypertension. *Am J Ophthalmol* 95:229-232, 1983
- Pinichpongse S, Doberstyn EB, Cullen JR, Yisunsri L, Thongsombun Y, Thimasarn K: An evaluation of five regimens for the outpatient therapy of falciparum malaria in Thailand 1980-81. *Bull WHO* 60(6):907-912, 1982
- Pugh RNH, Teesdale CH: Single dose oral treatment in urinary schistosomiasis: A double blind trial. *Br Med J* 286:429-432, 1983
- Pyati SP, Pildes RS, Jacobs NM, Ramamurthy RS, Yeh TF, Raval DS, Lilien LD, Amma P, Metzger WI: Penicillin in infants weighing two kilograms or less with early-onset group B streptococcal disease. *N Engl J Med* 308(23):1383-1389, 1983
- Ribeiro G, Palmer MK: Adjuvant tamoxifen for operable carcinoma of the breast: Report of clinical trial by the Christie Hospital and Holt Radium Institute. *Br Med J* 286:827-830, 1983
- Rickels K, Weise CC, Zal HM, Csanalosi I, Werblowsky J: Lofepamine and imipramine in unipolar depressed outpatients. *Acta Psychiatr Scand* 66:109-120, 1982
- Rubens RD, Hayward JL, Knight RK, Bulbrook RD, Fentiman IS, Chaudary M, Howell A, Bush H, Crowther D, Sellwood RA, George WD, Howat JMT: Controlled trial of adjuvant chemotherapy with melphalan for breast cancer. *Lancet* 1:839-843, 1983
- Rubin PC, Butters L, Clark DM, Reynolds B, Sumner DJ, Steedman D, Low RA, Reid JL: Placebo-controlled trial of atenolol in treatment of pregnancy-associated hypertension. *Lancet* 1:431-434, 1983

- Rydén L, Ariniego R, Arnman K, Herlitz J, Hjalmarson A, Holmberg S, Reyes C, Smedgard P, Svedberg K, Vedin A, Waagstein F, Waldenstrom A, Wilhelmsson C, Wedel H, Yamamoto M: A double-blind trial of metoprolol in acute myocardial infarction—Effects on ventricular tachyarrhythmias. *N Engl J Med* 308(11):614–618, 1983
- Shildt RA, Kennedy PS, Chen TT, Athens JW, O'Bryan RM, Balcerzak SP: Management of patients with metastatic adenocarcinoma of unknown origin: A Southwest Oncology Group study. *Cancer Treat Rep* 67(1):77–79, 1983
- Sponsel WE, Dallas NL, Burbridge L: Visual field survival: The response to timolol therapy in open-angle glaucoma. *Br J Ophthalmol* 67:220–227, 1983
- Thuesen L, Jorgensen JR, Kvistgaard HJ, Sorensen JA, Vaeth M, Jensen EB, Jensen JJ, Hagerup L: Effect of verapamil on enzyme release after early intravenous administration in acute myocardial infarction: Double blind randomised trial. *Br Med J* 286:1107–1108, 1983
- Tormey DC, Lippman ME, Edwards BK, Cassidy JG: Evaluation of tamoxifen doses with and without fluoxymesterone in advanced breast cancer. *Ann Intern Med* 98:139–144, 1983
- Toss G, Andersson R, Diffey BL, Fall PA, Larko O, Larsson L: Oral vitamin D and ultraviolet radiation for the prevention of vitamin D deficiency in the elderly. *Acta Med Scand* 212:157–161, 1982

Reports of Findings from Surgical Trials

- Gear MWL: Proximal gastric vagotomy versus long-term maintenance treatment with cimetidine for chronic duodenal ulcer: A prospective randomised trial. *Br Med J* 286:98–99, 1983
- MacLeod IA, Mills PR, MacKenzie JF, Joffe SN, Russell RI, Carter DC: Neodymium yttrium aluminium garnet laser photocoagulation for major haemorrhage from peptic ulcers and single vessels: A single blind controlled study. *Br Med J* 286:345–348, 1983
- Robertson DM, Ilstrup D: Direct, indirect, and sham laser photocoagulation in the management of central serous chorioretinopathy. *Am J Ophthalmol* 95:457–466, 1983

Reports of Findings from Diet Trials

- Harries AD, Jones LA, Danis V, Fifield R, Heatley RV, Newcombe RG, Rhodes J: Controlled trial of supplemented oral nutrition in Crohn's disease. *Lancet* 1:887–890, 1983
- Mellies MJ, Jandacek RJ, Taulbee JD, Tewksbury MB, Lamkin G, Baehler L, King P, Boggs D, Goldman S, Gouge A, Tsang R, Glueck CJ: A double-blind, placebo-controlled study of sucrose polyester in hypercholesterolemic outpatients. *Am J Clin Nutrition* 37:339–346, 1983
- Miettinen M, Turpeinen O, Karvonen MJ, Pekkarinen M, Paavilainen E, Elosuo R: Dietary prevention of coronary heart disease in women: The Finnish Mental Hospital Study. *Int J Epidemiol* 12(1):17–25, 1983
- Puska P, Iacono JM, Nissinen A, Korhonen HJ, Vartiainen E, Pietinen P, Dougherty R, Leino U, Mutanen M, Moisio S, Huttunen J: Controlled, randomised trial of the effect of dietary fat on blood pressure. *Lancet* 1:1–5, 1983
- Rouse IL, Beilin LJ, Armstrong BK, Vandongen R: Blood-pressure-lowering effect of a vegetarian diet: Controlled trial in normotensive subjects. *Lancet* 1:5–10, 1983

Reports of Findings from Other Trials

- Ambrose NS, Hares MM, Alexander-Williams J, Keighley MRB: Prospective randomised comparison of photocoagulation and rubber band ligation in treatment of haemorrhoids. *Br Med J* 286:1389–1391, 1983

- Avorn J, Soumerai SB: Improving drug-therapy decisions through educational outreach. A randomized controlled trial of academically based "detailing." *N Engl J Med* 308(24):1457-1463, 1983
- Bentley JM, Cormier P, Oler J: The Rural Dental Health Program: The effect of a school-based, dental health education program on children's utilization of dental services. *Am J Pub Health* 73(5):500-505, 1983
- Bulpitt CJ, Rowntree RK, Semmence A: A randomised controlled trial of the effects of screening for ulcer-type dyspepsia. *J Epidemiol Comm Health* 36:172-175, 1982
- Carpenter WT, Sadler JH, Light PD, Hanlon TE, Kurland AA, Penna MW, Reed WP, Wilkinson EH, Bartko JJ: The therapeutic efficacy of hemodialysis in schizophrenia. *N Engl J Med* 308(12):669-675, 1983
- Champlin R, Ho W, Gale RP: Antithymocyte globulin treatment in patients with aplastic anemia. A prospective randomized trial. *N Engl J Med* 308(3):113-118, 1983
- Dwosh IL, Giles AR, Ford PM, Pater JL, Anastassiades TP, Queen's University Plasmapheresis Study Group: Plasmapheresis therapy in rheumatoid arthritis. A controlled, double-blind, crossover trial. *N Engl J Med* 308(19):1124-1129, 1983
- Fischer BH, Marks M, Reich T: Hyperbaric-oxygen treatment of multiple sclerosis—A randomized, placebo-controlled, double-blind study. *N Engl J Med* 308(4):181-186, 1983
- Foot A, Erfurt JC: Hypertension control at the work site. Comparison of screening and referral alone, referral and follow-up, and on-site treatment. *N Engl J Med* 308(14):809-813, 1983
- Gillespie WA, Simpson RA, Jones JE, Nashef L, Teasdale C, Speller DCE: Does the addition of disinfectant to urine drainage bags prevent infection in catheterised patients? *Lancet* 1:1037-1039, 1983
- Gross SJ: Growth and biochemical response of preterm infants fed human milk or modified infant formula. *N Engl J Med* 308(5):237-241, 1983
- Hauser SL, Dawson DM, Leirich JR, Beal MF, Kevy SV, Propper RD, Mills JA, Weiner HL: Intensive immunosuppression in progressive multiple sclerosis—A randomized, three-arm study of high-dose intravenous cyclophosphamide, plasma exchange, and ACTH. *N Engl J Med* 308(4):173-180, 1983
- Holman RR, Dornan TL, Mayon-White V, Howard-Williams J, Orde-Peckar C, Jenkins L, Steemson J, Rolfe R, Smith B, Barbour D, McPherson K, Poon P, Rizza C, Mann JJ, Knight AH, Bron AJ, Turner RC: Prevention of deterioration of renal and sensory-nerve function by more intensive management of insulin-dependent diabetic patients—A two-year randomised prospective study. *Lancet* 1:204-208, 1983
- Janakiramaiah N, Channabasavanna SM, Murthy NSN: ECT/chlorpromazine combination versus chlorpromazine alone in acutely schizophrenic patients. *Acta Psychiatr Scand* 66:464-470, 1982
- Jeffery RW, Gillum R, Gerber WM, Jacobs D, Elmer PJ, Prineas RJ: Weight and sodium reduction for the prevention of hypertension: A comparison of group treatment and individual counseling. *Am J Pub Health* 73(6):691-693, 1983
- Keeler EB, Rolph JE: How cost sharing reduced medical spending of participants in the Health Insurance Experiment. *JAMA* 249(16):2220-2222, 1983
- Lauritzen T, Frost-Larsen K, Larsen H-W, Deckert T, Steno Study Group: Effect of 1 year of near-normal blood glucose levels on retinopathy in insulin-dependent diabetics. *Lancet* 1:200-204, 1983
- Levine DM: Health education for behavioral change—Clinical trial to public health program. *Johns Hopkins Med J* 151:215-219, 1982
- Newhouse JP, Spolsky V, Feldman B, Messing S, Black A, Goldberg G, Brook R, Bailit HL: Controlled trial in dental and medical insurance. *J Am Dental Assoc* 108:173-177, 1983

- Ornish D, Scherwitz LW, Doody RS, Kesten D, McLanahan SM, Brown SE, DePuey EG, Sonnemaker R, Haynes C, Lester J, McAllister GK, Hall RJ, Burdine JA, Gotto AM: Effects of stress management training and dietary changes in treating ischemic heart disease. *JAMA* 249(1):54–59, 1983
- Platt R, Polk BF, Murdock B, Rosner B: Reduction of mortality associated with nosocomial urinary tract infection. *Lancet* 1:893–897, 1983
- Shipley M, Berry H, Broster G, Jenkins M, Clover A, Williams I: Controlled trial of homoeopathic treatment of osteoarthritis. *Lancet* 1:97–98, 1983
- Stewart P, Hillan E, Calder AA: A randomised trial to evaluate the use of a birth chair for delivery. *Lancet* 1:1296–1298, 1983
- Templeton JL, Spence RAJ, Kennedy TL, Parks TG, MacKenzie G, Hanna WA: Comparison of infrared coagulation and rubber band ligation for first and second degree haemorrhoids: A randomised prospective clinical trial. *Br Med J* 286:1387–1389, 1983
- Wei N, Klippel JH, Huston DP, Hall RP, Lawley TJ, Balow JE, Steinberg AD, Decker JL: Randomised trial of plasma exchange in mild systemic lupus erythematosus. *Lancet* 1:17–22, 1983

Reports from Related Studies

- Chaitman BR, Alderman EL, Sheffield LT, Tong T, Fisher L, Mock MB, Weins RD, Kaiser GC, Roitman D, Berger R, Gersh B, Schaff H, Bourassa MG, Killip T, Participating CASS Medical Centers: Use of survival analysis to determine the clinical significance of new Q waves after coronary bypass surgery. *Circulation* 67(2):302–309, 1983
- Dorros G, Cowley MJ, Simpson J, Bentivoglio LG, Block PC, Bourassa M, Detre K, Gosselin AJ, Gruntzig AR, Kelsey SF, Kent KM, Mock MB, Mullin SM, Myler RK, Passamani ER, Stertz SH, Williams DO: Percutaneous transluminal coronary angioplasty: Report of complications from the National Heart, Lung, and Blood Institute PTCA Registry. *Circulation* 67(4):723–729, 1983
- Gersh BJ, Kronmal RA, Frye RL, Schaff HV, Ryan TJ, Gosselin AJ, Kaiser GC, Killip T, Participants in the Coronary Artery Surgery Study: Coronary arteriography and coronary artery bypass surgery: Morbidity and mortality in patients ages 65 years or older—A report from the Coronary Artery Surgery Study. *Circulation* 67(3):483–491, 1983
- Hardy RJ, Hawkins CM: The impact of selected indices of antihypertensive treatment on all-cause mortality. *Am J Epidemiol* 117(5):566–574, 1983
- Jacobs DR, Barrett-Connor E: Retest reliability of plasma cholesterol and triglyceride. The Lipid Research Clinics Prevalence Study. *Am J Epidemiol* 116(6):878–885, 1982
- Kornitzer M, De Backer G, Dramaix M, Kittel F, Thilly C, Graffar M, Vuylsteek K: Belgian Heart Disease Prevention Project: Incidence and mortality results. *Lancet* 1:1066–1070, 1983
- Nash S, Tilley BC, Kurland LT, Gundersen J, Barnes AB, Labarthe D, Donohew PS, Kovacs L: Identifying and tracing a population at risk: The DESAD Project experience. *Am J Pub Health* 73(3):253–259, 1983
- Rose G, Tunstall-Pedoe HD, Heller RF: UK Heart Disease Prevention Project: Incidence and mortality results. *Lancet* 1:1062–1066, 1983
- Rosner B, Polk BF: Predictive values of routine blood pressure measurements in screening for hypertension. *Am J Epidemiol* 117(4):429–442, 1983
- Ruchlin HS, Morris JN: Pennsylvania's domiciliary care experiment: II. Cost-benefit implications. *Am J Pub Health* 73(6):654–660, 1983
- Ryan TJ, Bailey KR, McCabe CH, Luk S, Fisher LD, Mock MB, Killip T: The effects of digitalis on survival in high-risk patients with coronary artery disease. The Coronary Artery Surgery Study (CASS). *Circulation* 67(4):735–742, 1983

- Sherwood S, Morris JN: The Pennsylvania domiciliary care experiment: I. Impact on Quality of life. *Am J Pub Health* 73(6):646–653, 1983
- Southall DP, Richards JM, de Swiet M, Arrowsmith WA, Cree JE, Fleming PJ, Franklin AJ, L'E Orme R, Radford MJ, Wilson AJ, Shannon DC, Alexander JR, Brown NJ, Shinebourne EA: Identification of infants destined to die unexpectedly during infancy: Evaluation of predictive importance of prolonged apnoea and disorders of cardiac rhythm or conduction. *Br Med J* 286:1092–1096, 1983
- Thelle DS, Arnesen E, Forde OH: The Tromsø Heart Study: Does coffee raise serum cholesterol? *N Engl J Med* 308(24):1454–1457, 1983
- UK-TIA Study Group: Variation in the use of angiography and carotid endarterectomy by neurologists in the UK-TIA Aspirin Trial. *Br Med J* 286:514–517, 1983