

Execution of nonsense-mediated mRNA decay: what defines a substrate?

Indrani Rebbapragada and Jens Lykke-Andersen

The nonsense-mediated mRNA decay (NMD) pathway targets mRNAs with premature termination codons as well as a subset of normal mRNAs for rapid decay. Emerging evidence suggests that mRNAs become NMD substrates based on the composition of the mRNP downstream of the translation termination event, which either stimulates or antagonizes recruitment of the NMD machinery. The NMD mRNP subsequently undergoes several remodeling events, which involve hydrolysis of ATP by the NMD factor Upf1 and in metazoans, a phosphorylation/dephosphorylation cycle of Upf1 mediated by Smg proteins. This leads to mRNA decay following translational repression. Recent evidence suggests that in *Drosophila* and human cells, decay is initiated by the endonuclease Smg6.

Address

Department of Molecular, Cellular and Developmental Biology,
University of Colorado, Boulder, CO 80309, USA

Corresponding author: Lykke-Andersen, Jens (jens.lykke-andersen@colorado.edu)

Current Opinion in Cell Biology 2009, **21**:394–402

This review comes from a themed issue on
Nucleus and gene expression
Edited by Elisa Izaurralde and Phillip Zamore

Available online 7th April 2009

0955-0674/\$ – see front matter
© 2009 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.ceb.2009.02.007](https://doi.org/10.1016/j.ceb.2009.02.007)

Introduction

The expression of protein-coding genes in eukaryotes involves the orchestration of transcriptional and post-transcriptional processes. To ensure the fidelity of these processes, the eukaryotic cell has evolved several quality-control mechanisms. One such mechanism is the nonsense-mediated mRNA decay (NMD) pathway. NMD rids the cell of aberrant mRNAs that have acquired premature translation termination codons (PTCs) [1–7]. In addition, the NMD pathway serves to destabilize a subset of apparently normal mRNAs [8,9]. Several *trans*-acting factors critical for the NMD pathway have been identified. These include Upf1, a DEAD-box ATPase, Upf2, and Upf3, and in metazoans, Smg1, Smg5, Smg6, and Smg7. In this review we discuss recent insights into the mechanisms by which Upf and Smg proteins are

specifically recruited to NMD substrates, and how they subsequently activate translational repression and mRNA decay.


Which mRNAs are substrates of NMD?

Recent evidence suggests that the NMD pathway targets not only mRNAs that have acquired premature termination codons (PTCs), but also a subset of endogenous transcripts lacking PTCs (Figure 1). Global gene-expression profiling of several eukaryotes, where components of the NMD pathway were either transiently or stably inactivated, revealed an upregulation of about 1–10% of cellular transcripts [8,10–12,13^{••},14^{••},15^{••},16]. Although a subset of these transcripts may not be direct targets of NMD [16], some have been subsequently confirmed experimentally. Many of these endogenous NMD targets do not contain PTCs, although some have recognizable features such as upstream open reading frames (uORFs) that would make translation termination events appear as premature [8,9].

Although some mRNAs lacking PTCs are targets of NMD, mRNAs that have acquired PTCs likely constitute a major fraction of NMD substrates in the cell (Figure 1). PTC-containing mRNAs can arise from erroneous gene expression such as errors in transcription, faulty mRNA processing, somatic mutations, or nonproductive programmed DNA rearrangements, for example those that occur in the T-cell receptor (TCR) and immunoglobulin (Ig) genes [4–6,17]. PTCs can also be inheritable as in the case of genetic mutations in disease genes, expressed pseudogenes and small nucleolar RNA (snoRNA) host genes that have lost their protein-coding potential [17]. However, not all PTCs trigger NMD. For example, those PTCs positioned close to the normal termination or initiation codons often fail to trigger NMD [18–24].

A major contributor of PTC-containing mRNAs is likely faulty or alternative splicing. For example, a recent study reported a strong bias for in-frame termination codons in introns in the ciliate *P. tetraurelia*, suggesting a reliance on NMD to eliminate inefficiently spliced transcripts [25^{••}]. In mammals, the analyses of databases of expressed sequence tags suggested that one-third of alternative splicing events have the potential to generate PTC-containing mRNAs [26–28]. However, quantitative microarray splice variant profiling experiments revealed that many PTC-containing alternatively spliced transcripts are not significantly upregulated upon Upf1 knockdown [29]. Interestingly, several mRNAs encoding

Figure 1



Nonsense-mediated mRNA decay (NMD) can target both mRNAs that have acquired PTCs and normal mRNAs. NMD targets a majority of mRNAs that contain premature termination codons (PTCs) for rapid decay (top; red arrow), although some PTC-containing mRNAs escape NMD and are stable (top; green arrow). NMD also targets a subset of apparently normal mRNAs for NMD (bottom; red arrow).


splicing-activator proteins are targets of NMD because of alternative splicing events that cause the inclusion of conserved stop codon-containing exons [27^{••},30–32]. This may provide a negative feedback loop that uses alternative splicing to autoregulate the levels of splicing factor proteins, while the NMD pathway depletes the cell of the PTC-containing alternatively spliced products that would otherwise produce truncated proteins.

How is a translation termination event that induces NMD discriminated from normal termination?

Evidence from multiple studies suggests that the composition of the messenger ribonucleoprotein (mRNP) downstream of the terminating ribosome is a key determinant for NMD. Whether the downstream mRNP triggers NMD may be decided by the arrangement of components of the mRNP that stimulate or antagonize NMD (Figure 2) [33^{••},34^{••},35^{••},36^{••},37,38^{••}]. To allow for the observation that both normal termination codons and PTCs can elicit NMD we will here avoid the often used term 'PTC recognition' and instead refer to translation termination events that trigger NMD as NMD-inducing termination events.

Factors that stimulate NMD when part of the mRNP downstream of a termination event have been identified in both *Saccharomyces cerevisiae* and mammals. Early studies identified *S. cerevisiae* Hrp1p as a protein that coimmunoprecipitates (co-IPs) with Upf1, and associates with a downstream sequence element important for NMD of PTC-containing PGK1 mRNA [39]. Similarly, in mam-

Figure 2



Model for the mechanism of discrimination between a normal and an NMD-inducing translation termination event. The composition of the mRNP downstream of the terminating ribosome may dictate whether NMD occurs through a competition between stimulators and antagonists of Upf complex recruitment. NMD-stimulating factors, such as Hrp1p (red) in *S. cerevisiae* and the EJC and RNPS1 (brown) in human cells, stimulate NMD, likely by recruiting the Upf proteins to the mRNP downstream of the termination event. By contrast, PABPC (light green), and likely other factors including *S. cerevisiae* Pub1p (dark green), antagonizes NMD when in close downstream proximity of the termination event.

malian cells the exon-junction complex (EJC), which is deposited upstream of exon–exon junctions after pre-mRNA splicing, and the EJC-associated splicing-activator RNPS1, both interact directly with the NMD factor Upf3 and stimulate NMD of a number of tested substrates when positioned downstream of a translation termination event [40–48,49^{••}]. Interestingly, several splicing-activator SR proteins have been shown to stimulate NMD when overexpressed in human cells, although the mechanism is unknown [50,51]. Thus, factors such as Hrp1p, the EJC, and RNPS1 stimulate NMD, possibly by increasing the affinity of the Upf complex to the mRNP downstream of a termination event and would be predicted to be absent from normal 3'-UTRs. Consistently, introns are rarely found in the 3'-UTRs of mammalian genes [52]. Some EJC and SR proteins have also been observed to activate translation when located upstream of a termination event [53,54], but the relation between this activity and NMD remains unclear.

Recent evidence has identified cytoplasmic poly(A)-binding protein (PABPC) as an NMD antagonist in *S. cerevisiae*, *D. melanogaster*, and human cells. This includes observations that mRNAs with artificially extended

3'-UTRs, which positions the poly(A)-tail and PABPC distal to the translation termination event, are targeted for NMD, while artificial recruitment of PABPC in downstream proximity of the termination codon, by tethering or 3'-UTR looping, impairs NMD [33^{••},34^{••},35^{••},36^{••},38^{••},55^{••}]. The mechanism by which PABPC antagonizes NMD remains unclear. However, direct interactions have been reported of both Upf1 and PABPC with eukaryotic release factor 3 (eRF3) [36^{••},56–61], and human PABPC was observed to out-compete the interaction between Upf1 and eRF3 *in vitro* [38^{••}]. Additionally, overexpression of PABPC suppresses a yeast temperature-sensitive allele of eRF3 [56], while depletion of PABPC in human cell lines results in increased read-through of a dual luciferase reporter mRNA [36^{••}], indicating that PABPC stimulates translation termination. Together these observations implicate PABPC as an NMD-antagonizing factor that when present in downstream proximity of a termination codon stimulates normal termination and prevents the recruitment of the Upf complex and NMD.

Factors other than PABPC that antagonize NMD likely exist. For example, evidence suggests that the NMD machinery of *S. cerevisiae* is capable of identifying NMD substrates even in the absence of PABPC or a poly(A)-tail [62^{••}]. Pub1p has been identified as an *S. cerevisiae* factor that associates with a stabilizer element downstream of the uORF of GCN4 mRNA and antagonizes NMD when tethered at the same position [63]. Similarly, the 3'-UTR of the Rous sarcoma virus (RSV) gag gene contains a stability element that antagonizes NMD [64]. Future studies should reveal the relative importance of NMD-stimulating versus NMD-antagonizing factors in defining NMD substrates, and the molecular mechanisms by which these factors act. Moreover, an important question for future studies is whether the recruitment of NMD-modulating factors can be a regulated event, thereby using the NMD pathway to control gene expression upon cellular cues.

Does NMD take place during a pioneer round of translation?

Observations that the yeast nuclear cap-binding protein cbc1p co-IPs with eIF4G and is able to support translation *in vitro* led to the hypothesis that a newly exported mRNA can engage in translation before the exchange of the nuclear cap binding complex (CBC) for cytoplasmic translation initiation factor eIF4E [65]. The pioneer round of translation hypothesis for NMD posits that translation termination events that trigger NMD in mammalian cells can occur only during a pioneer round of translation, defined as the translation that takes place while the mRNA is associated with CBC. By contrast, according to this hypothesis, once eIF4E replaces the CBC the mRNA is immune to NMD [66].

Much of the evidence for the pioneer round of translation model comes from interaction studies. First, the ratio of an NMD substrate to its cognate wild-type mRNA that co-purifies from human cell extracts with CBP80 (the large subunit of CBC) or eIF4E was indistinguishable by RT-PCR assays [67]. Second, NMD factors, including Upf proteins, Smg1, and EJC components, were observed to co-IP in the absence of RNase with CBP80 but not with eIF4E. By contrast, components of the general translation machinery, co-IP with both CBP80 and eIF4E [68,69]. Third, knockdown of CBP80 was reported to impair the co-IP of Upf1 with Upf2, while far-western and pull down assays suggest that recombinant CBP80 interacts with Upf1 and stimulates Upf1–Upf2 interaction [69]. Functional evidence for the pioneer round of translation model comes from the observation that overexpression of eIF4E-binding protein 1 (4E-BP1) at levels that cause translational repression of reporter mRNAs in human cells does not show a corresponding increase in the steady-state levels of tested NMD substrates as measured by RT-PCR [68]. These observations suggest that eIF4E-initiated translation is not limiting for NMD of tested reporters, and that at steady state, NMD factors are primarily found on CBC-associated mRNA. This is consistent with the idea that NMD primarily takes place on CBC-associated mRNA. However, these observations do not provide evidence that eIF4E-associated mRNA is immune to NMD. The most direct evidence that NMD may be restricted to CBC-associated mRNA comes from reports that knockdown of endogenous CBP80 by RNAi causes increased steady-state levels of tested NMD reporters, which is rescued upon expression of exogenous CBP80 [69]. However, this observation leaves open the possibility that pleiotropic effects could interfere with NMD in an indirect manner. Interestingly, a PTC-containing mRNA in which translation is driven by an EMCV internal ribosome entry site was targeted for NMD, suggesting that cap-independent translation initiation can sustain NMD [70]. However knockdown of CBC components was reported to cause upregulation of this mRNA [71].

All evidence suggests that NMD in *S. cerevisiae* can take place on eIF4E-associated mRNA. First, NMD substrates expressed and translated under conditions of Upf protein depletion, are rapidly degraded upon restoration of the depleted Upf proteins at a time when CBC is presumed to have been replaced by eIF4E [72]. Second, an NMD reporter kept stably in polysomal fractions by drug-induced inhibition of translation elongation is rapidly degraded upon removal of the drug [73]. Third, the CBC is not limiting for *S. cerevisiae* NMD, as the depletion of cbc1p does not affect the abundance of tested NMD substrates [74]. Finally, analysis of the decay rates of a reporter mRNA engineered with programmed -1 ribosomal frame-shift signals' suggested that mRNAs


were available for NMD after the first ribosome has translated the mRNA [75]. While these lines of evidence suggest that eIF4E-associated mRNA is susceptible to NMD in *S. cerevisiae*, it has not been directly tested whether CBC-associated mRNA can also be targeted for NMD.

Several predictions from the pioneer round hypothesis of mammalian NMD remain to be tested. For example, the prediction that a fraction of NMD substrates should escape NMD by undergoing replacement of CBP80 with eIF4E before recognition by the NMD pathway, and thus should result in two-component decay kinetics has not been carefully tested. In previous experiments that indicated two-component NMD kinetics [76–78], it was not ruled out that the apparent slow phase of decay was not a result of incomplete transcriptional repression. Moreover, experiments monitoring decay of NMD substrates expressed under conditions of NMD repression followed by reactivation of NMD have not been performed in mammalian cells.

What happens to the NMD mRNP after translation termination?

Although the exact order of events downstream of an NMD-inducing termination event remains unknown, recent evidence suggests that at least two transitions in the NMD mRNP occur after the recruitment of the Upf proteins and before the degradation of the NMD substrate. These involve ATP hydrolysis by Upf1 and, at least in metazoans, a Upf1 phosphorylation/dephosphorylation cycle (Figure 3). Upf1 is a DEAD-box ATPase that shows RNA-dependent ATPase activity and ATP-dependent 5'–3' RNA helicase activity *in vitro* [3–5]. Mutant Upf1 proteins that fail to bind or hydrolyze ATP are inactive in NMD and cause the accumulation of the NMD mRNP in P-bodies [79,80^{••}]. This suggests that the ATPase activity of Upf1 is important for steps in NMD downstream of translation termination. Consistent with this, Upf2 and Upf3 stimulate ATP hydrolysis by Upf1 [49^{••}] and NMD substrates accumulate in P-bodies upon depletion of Upf2 or Upf3 in *S. cerevisiae* [80^{••}]. While it is thus clear that the ATPase activity of Upf1 is critical for

Figure 3



Transitions in the NMD mRNP leading to translation repression and decay. **(a)** For an mRNA to be targeted for NMD, it needs to undergo at least one round of translation, which results in the recruitment of Upf proteins to the terminating ribosome, followed by translational repression and degradation by both exonucleolytic and endonucleolytic processes. **(b)** After recruitment to the NMD mRNP and before decay, Upf1 undergoes Smg1-mediated phosphorylation and hydrolyzes ATP, followed by Smg5/Smg7-stimulated dephosphorylation by PP2A. Phosphorylation and ATP hydrolysis by Upf1 is stimulated by Upf2 and Upf3. See text for details.

NMD, the specific transition in the NMD mRNP that occurs upon ATP hydrolysis remains unknown.

Another critical transition in the metazoan NMD mRNP is a Upf1 phosphorylation/dephosphorylation cycle (Figure 3). The Upf1 kinase in metazoans is the phosphoinositol 3-kinase (PIK)-related protein kinase Smg1. Depletion of Upf2 or Upf3 inhibits the accumulation of phosphorylated Upf1, suggesting that Upf2 and Upf3 stimulate Smg1 activity [81^{••},82]. Biochemical and structural studies suggest that phospho-Upf1 interacts with Smg5 via the Upf1 N-terminus, and with Smg7 through interactions between the phosphorylated SQ-motifs of Upf1 and a 14-3-3 like domain of Smg7 [83–88]. Smg5 and Smg6 also contain predicted 14-3-3 like domains, suggesting that these proteins may also associate with Upf1 through the phospho-SQ-motifs [89]. Smg5 and Smg7 co-IP with protein phosphatase 2A (PP2A) and their depletion causes accumulation of phospho-Upf1, suggesting that Smg5 and Smg7 mediate dephosphorylation of Upf1 through the recruitment of PP2A. Phosphorylation and dephosphorylation of Upf1 is critical for metazoan NMD as evidenced by the impairment of NMD under conditions that inactivate these events [82,83,87,90^{••},91,92]. It is not clear whether phosphorylation of Upf1 is important for NMD in *S. cerevisiae*, although Upf1 has been observed to be a phospho-protein [93], and the protein Ebs1p has been identified as a putative *S. cerevisiae* ortholog of human Smg7 that interacts with Upf1 and inhibits NMD when overexpressed or depleted [94]. An important goal for future studies will be to understand the significance of the mRNP transitions that take place in the NMD pathway downstream of translation termination. It has been speculated that they may serve as a second step after translation termination to discriminate true NMD substrates from normal mRNAs [95].

How are NMD substrates translationally silenced and degraded?

The transitions in the NMD mRNP described above lead to decay of the NMD substrate, which is likely preceded by the inhibition of translation. Early studies in *S. cerevisiae* provided evidence that the translation of NMD substrates was downregulated several-fold more than their stability, suggesting that the recognition of an NMD substrate results in translational repression in addition to mRNA decay [96]. In mammalian cells, reduced expression of proteins from PTC-containing mRNAs has also been observed. However, this may at least in some cases be attributed to intrinsic instability of the truncated protein rather than that of translational repression [97,98]. Two-hybrid analyses identified translation initiation factor eIF4A1 as a weak interaction partner of eIF4G-homology (4GH) domains in *S. pombe* Upf2, and it was speculated that those interactions might contribute to translational repression [99]. More recent

observations suggest that phospho-Upf1 interacts with translation initiation factor eIF3 and interferes with recruitment of the 60S ribosomal subunit to the 40S preinitiation complex [100,101^{••}].

Several nuclease activities have been implicated in the decay of NMD substrates. Early observations in *S. cerevisiae* suggested that decapping is a rate-limiting step in NMD [102]. More recently, additional activation of deadenylation and 5'–3' and 3'–5' exonucleolytic activities by the NMD machinery has been observed in both *S. cerevisiae* and mammalian cells [3,4,6,66]. Interestingly, recent studies have implicated the NMD factor Smg6 as an endonuclease in NMD in *D. melanogaster* and humans [103^{••},104,105^{••}]. The C-terminal PilT N-terminus (PIN) domain of Smg6 shows endonuclease activity *in vitro* [106] and catalytically inactive Smg6 fails to support NMD in human and *Drosophila* cells suggesting that endonucleolytic activity of Smg6 is important for NMD [103^{••},105^{••}]. Thus both exonucleolytic and endonucleolytic activities are implicated in NMD. It remains to be determined which decay mechanism is predominant in various organisms.

Perspectives

The NMD pathway plays an important role in modulating gene expression by downregulating not only aberrant mRNAs that contain PTCs but also a subset of normal cellular mRNAs. Whether a translation termination event induces NMD seems to be dictated by the arrangement of the mRNP downstream of the termination event. Future studies should reveal the mechanisms and relative contributions to NMD in various organisms of NMD-stimulating and NMD-antagonizing mRNP factors. It will also be important to learn the specific mechanisms by which transitions in the NMD mRNP lead to translational repression and subsequent decay. Moreover, an important question for future studies is whether the NMD pathway can be regulated upon specific cellular cues to control expression of normal mRNAs. Given the rapid pace of the NMD field, the answers to many of these questions may be coming in the near future.

Acknowledgements

We thank Drs Tom Blumenthal, Sebastien Durand, and Toby Franks for critical comments on the manuscript. Research on NMD in the Lykke-Andersen laboratory is supported by funding from the National Science Foundation (MCB-0645798) to JL.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

•• of outstanding interest

1. Akimitsu N: **Messenger RNA surveillance systems monitoring proper translation termination.** *J Biochem* 2008, **143**:1–8.
2. Baker KE, Parker R: **Nonsense-mediated mRNA decay: terminating erroneous gene expression.** *Curr Opin Cell Biol* 2004, **16**:293–299.

3. Behm-Ansmant I, Kashima I, Rehwinkel J, Sauliere J, Wittkopp N, Izaurralde E: **mRNA quality control: an ancient machinery recognizes and degrades mRNAs with nonsense codons.** *FEBS Lett* 2007, **581**:2845-2853.
4. Chang YF, Imam JS, Wilkinson MF: **The nonsense-mediated decay RNA surveillance pathway.** *Annu Rev Biochem* 2007, **76**:51-74.
5. Isken O, Maquat LE: **The multiple lives of NMD factors: balancing roles in gene and genome regulation.** *Nat Rev Genet* 2008, **9**:699-712.
6. Muhlemann O, Eberle AB, Stalder L, Zamudio Orozco R: **Recognition and elimination of nonsense mRNA.** *Biochim Biophys Acta* 2008, **1779**:538-549.
7. Shyu AB, Wilkinson MF, van Hoof A: **Messenger RNA regulation: to translate or to degrade.** *EMBO J* 2008, **27**:471-481.
8. Mendell JT, Sharifi NA, Meyers JL, Martinez-Murillo F, Dietz HC: **Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutes genomic noise.** *Nat Genet* 2004, **36**:1073-1078.
9. He F, Li X, Spatrick P, Casillo R, Dong S, Jacobson A: **Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5' to 3' mRNA decay pathways in yeast.** *Mol Cell* 2003, **12**:1439-1452.
10. Rehwinkel J, Letunic I, Raes J, Bork P, Izaurralde E: **Nonsense-mediated mRNA decay factors act in concert to regulate common mRNA targets.** *RNA* 2005, **11**:1530-1544.
11. Wittmann J, Hol EM, Jack HM: **hUPF2 silencing identifies physiologic substrates of mammalian nonsense-mediated mRNA decay.** *Mol Cell Biol* 2006, **26**:1272-1287.
12. Guan Q, Zheng W, Tang S, Liu X, Zinkel RA, Tsui KW, Yandell BS, Culbertson MR: **Impact of nonsense-mediated mRNA decay on the global expression profile of budding yeast.** *PLoS Genet* 2006, **2**:e203.
13. Johns L, Grimson A, Kuchma SL, Newman CL, Anderson P: **Caenorhabditis elegans SMG-2 selectively marks mRNAs containing premature translation termination codons.** *Mol Cell Biol* 2007, **27**:5630-5638.
- The authors show evidence that *C. elegans* Smg2 (Upf1) preferentially associates with mRNAs that contain premature termination codons and that this association is stimulated by Smg3 (Upf2) and Smg4 (Upf3).
14. Johansson MJ, He F, Spatrick P, Li C, Jacobson A: **Association of yeast Upf1p with direct substrates of the NMD pathway.** *Proc Natl Acad Sci U S A* 2007, **104**:20872-20877.
- The authors identify direct targets of the NMD pathway in *S. cerevisiae*, by comparing microarray data sets from mRNAs isolated by immunoprecipitation of Upf1, mRNAs upregulated in Upf1 depletion conditions, and mRNAs downregulated upon reactivation of Upf1.
15. Weischenfeldt J, Damgaard I, Bryder D, Theilgaard-Monch K, Thoren LA, Nielsen FC, Jacobsen SE, Nerlov C, Porse BT: **NMD is essential for hematopoietic stem and progenitor cells and for eliminating by-products of programmed DNA rearrangements.** *Genes Dev* 2008, **22**:1381-1396.
- This study shows that conditional depletion of Upf2 in mouse hematopoietic cells resulted in dramatic loss of hematopoietic stem and progenitor populations, while Upf2 depletion in more differentiated myeloid lineage cells did not have as strong an effect, suggesting that NMD is important in proliferating cells. Moreover, Upf2 depletion was shown to result in the upregulation of aberrant T-cell receptor mRNA, pseudogenes and many mRNAs encoding splice factors.
16. Viegas MH, Gehring NH, Breit S, Hentze MW, Kulozik AE: **The abundance of RNPS1, a protein component of the exon junction complex, can determine the variability in efficiency of the nonsense mediated decay pathway.** *Nucleic Acids Res* 2007, **35**:4542-4551.
17. Weischenfeldt J, Lykke-Andersen J, Porse B: **Messenger RNA surveillance: neutralizing natural nonsense.** *Curr Biol* 2005, **15**:R559-R562.
18. Inacio A, Silva AL, Pinto J, Ji X, Morgado A, Almeida F, Faustino P, Lavinha J, Liebhaver SA, Romao L: **Nonsense mutations in close proximity to the initiation codon fail to trigger full nonsense-mediated mRNA decay.** *J Biol Chem* 2004, **279**:32170-32180.
19. Denecke J, Kranz C, Kemming D, Koch HG, Marquardt T: **An activated 5' cryptic splice site in the human ALG3 gene generates a premature termination codon insensitive to nonsense-mediated mRNA decay in a new case of congenital disorder of glycosylation type Id (CDG-Id).** *Hum Mutat* 2004, **23**:477-486.
20. Wang J, Gudikote JP, Olivas OR, Wilkinson MF: **Boundary-independent polar nonsense-mediated decay.** *EMBO Rep* 2002, **3**:274-279.
21. Cao D, Parker R: **Computational modeling and experimental analysis of nonsense-mediated decay in yeast.** *Cell* 2003, **113**:533-545.
22. Thermann R, Neu-Yilik G, Deters A, Frede U, Wehr K, Hagemeyer C, Hentze MW, Kulozik AE: **Binary specification of nonsense codons by splicing and cytoplasmic translation.** *EMBO J* 1998, **17**:3484-3494.
23. Zhang J, Sun X, Qian Y, Maquat LE: **Intron function in the nonsense-mediated decay of beta-globin mRNA: indications that pre-mRNA splicing in the nucleus can influence mRNA translation in the cytoplasm.** *RNA* 1998, **4**:801-815.
24. Zhang J, Maquat LE: **Evidence that translation reinitiation abrogates nonsense-mediated mRNA decay in mammalian cells.** *EMBO J* 1997, **16**:826-833.
25. Jaillon O, Bouhouche K, Gout JF, Aury JM, Noel B, Saudemont B, Nowacki M, Serrano V, Porcel BM, Segurens B *et al.*: **Translational control of intron splicing in eukaryotes.** *Nature* 2008, **451**:359-362.
- This paper shows evidence from several eukaryotes that the occurrence of introns that generate a frame-shift or PTCs in the event of intron retention is significantly higher than that would be expected from random, suggesting an evolutionary bias for introns that trigger NMD when retained. The authors also show that the depletion of Upf1 by RNAi results in the accumulation of unspliced mRNAs in *P. tetraurelia* suggesting that NMD compensates for inefficient splicing events.
26. Lewis BP, Green RE, Brenner SE: **Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans.** *Proc Natl Acad Sci U S A* 2003, **100**:189-192.
27. Saltzman AL, Kim YK, Pan Q, Fagnani MM, Maquat LE, Blencowe BJ: **Regulation of multiple core spliceosomal proteins by alternative splicing-coupled nonsense-mediated mRNA decay.** *Mol Cell Biol* 2008, **28**:4320-4330.
- This paper identifies several genes encoding spliceosomal proteins that contain alternative PTC-containing exons. Overexpression of the splicing factors results in increased inclusion of the PTC-containing exons suggesting that alternative splicing followed by NMD is employed as a feedback mechanism for regulating expression of splicing factors.
28. Sorek R, Shamir R, Ast G: **How prevalent is functional alternative splicing in the human genome?** *Trends Genet* 2004, **20**:68-71.
29. Pan Q, Saltzman AL, Kim YK, Misquitta C, Shai O, Maquat LE, Frey BJ, Blencowe BJ: **Quantitative microarray profiling provides evidence against widespread coupling of alternative splicing with nonsense-mediated mRNA decay to control gene expression.** *Genes Dev* 2006, **20**:153-158.
30. Barberan-Soler S, Zahler AM: **Alternative splicing regulation during C. elegans development: splicing factors as regulated targets.** *PLoS Genet* 2008, **4**:e1000001.
31. Lareau LF, Inada M, Green RE, Wengrod JC, Brenner SE: **Unproductive splicing of SR genes associated with highly conserved and ultraconserved DNA elements.** *Nature* 2007, **446**:926-929.
32. Ni JZ, Grate L, Donohue JP, Preston C, Nobida N, O'Brien G, Shiue L, Clark TA, Blume JE, Ares M Jr: **Ultraconserved elements are associated with homeostatic control of splicing regulators by alternative splicing and nonsense-mediated decay.** *Genes Dev* 2007, **21**:708-718.
33. Amrani N, Ganesan R, Kervestin S, Mangus DA, Ghosh S, Jacobson A: **A faux 3'-UTR promotes aberrant termination and triggers nonsense-mediated mRNA decay.** *Nature* 2004, **432**:112-118.

This study shows evidence using *in vitro* toe-printing assays that ribosomes fail to be released from the mRNA when terminating at premature termination codons and that this is dependent on Upf1. Normal termination is restored and NMD prevented when the termination codon is flanked by a normal 3'-UTR or by tethered PABPC providing evidence that the mRNP downstream of the termination codon influences translation termination and NMD.

34. Behm-Ansmant I, Gatfield D, Rehwinkel J, Hilgers V, Izaurralde E:
 •• **A conserved role for cytoplasmic poly(A)-binding protein 1 (PABPC1) in nonsense-mediated mRNA decay.** *EMBO J* 2007, **26**:1591-1601.

This study provides evidence that short 3'-UTRs and PABPC tethered in downstream proximity of a termination codon in *D. melanogaster* inhibits NMD suggesting that that PABPC serves to antagonize NMD in *D. melanogaster*.

35. Eberle AB, Stalder L, Mathys H, Orozco RZ, Muhlemann O:
 •• **Posttranscriptional gene regulation by spatial rearrangement of the 3' untranslated region.** *PLoS Biol* 2008, **6**:e92.

This study provides evidence that decreasing the distance between the termination codon and PABPC, by tethering or through an engineered 3'-UTR that folds back upon itself, impairs NMD in human cells, suggesting that PABPC antagonizes NMD when in spatial proximity to the termination event.

36. Ivanov PV, Gehring NH, Kunz JB, Hentze MW, Kulozik AE:
 •• **Interactions between UPF1, eRFs, PABP and the exon junction complex suggest an integrated model for mammalian NMD pathways.** *EMBO J* 2008, **27**:736-747.

The authors provide evidence that tethering PABPC downstream of a premature termination codon inhibits NMD in human cells and that depletion of PABPC by RNAi increases translational read-through of a dual luciferase reporter mRNA suggesting that PABPC not only antagonizes NMD but also stimulates the efficiency of translation termination.

37. Kerenyi Z, Merai Z, Hiripi L, Benkovics A, Gyula P, Lacomme C, Barta E, Nagy F, Silhavy D: **Inter-kingdom conservation of mechanism of nonsense-mediated mRNA decay.** *EMBO J* 2008, **27**:1585-1595.

38. Singh G, Rebbapragada I, Lykke-Andersen J: **A competition between stimulators and antagonists of Upf complex recruitment governs human nonsense-mediated mRNA decay.** *PLoS Biol* 2008, **6**:e111.

This paper shows evidence that PABPC antagonizes NMD in human cells when positioned in proximity of the termination codon through shortening the 3'-UTR or PABPC tethering, even when an intron is present in the 3'-UTR. Moreover evidence is shown that PABPC can outcompete eRF3-Upf1 interaction *in vitro*, suggesting that PABPC may antagonize NMD in part by preventing Upf1 from being recruited to the terminating ribosome.

39. Gonzalez CI, Ruiz-Echevarria MJ, Vasudevan S, Henry MF, Peltz SW: **The yeast hnRNP-like protein Hrp1/Nab4 marks a transcript for nonsense-mediated mRNA decay.** *Mol Cell* 2000, **5**:489-499.

40. Singh G, Jakob S, Kleedehn MG, Lykke-Andersen J: **Communication with the exon-junction complex and activation of nonsense-mediated decay by human Upf proteins occur in the cytoplasm.** *Mol Cell* 2007, **27**:780-792.

41. Shibuya T, Tange TO, Sonenberg N, Moore MJ: **eIF4AIII binds spliced mRNA in the exon junction complex and is essential for nonsense-mediated decay.** *Nat Struct Mol Biol* 2004, **11**:346-351.

42. Palacios IM, Gatfield D, St Johnston D, Izaurralde E: **An eIF4AIII-containing complex required for mRNA localization and nonsense-mediated mRNA decay.** *Nature* 2004, **427**:753-757.

43. Lykke-Andersen J, Shu MD, Steitz JA: **Communication of the position of exon-exon junctions to the mRNA surveillance machinery by the protein RNPS1.** *Science* 2001, **293**:1836-1839.

44. Le Hir H, Izaurralde E, Maquat LE, Moore MJ: **The spliceosome deposits multiple proteins 20-24 nucleotides upstream of mRNA exon-exon junctions.** *EMBO J* 2000, **19**:6860-6869.

45. Le Hir H, Gatfield D, Izaurralde E, Moore MJ: **The exon-exon junction complex provides a binding platform for factors involved in mRNA export and nonsense-mediated mRNA decay.** *EMBO J* 2001, **20**:4987-4997.

46. Kim VN, Kataoka N, Dreyfuss G: **Role of the nonsense-mediated decay factor hUpf3 in the splicing-dependent exon-exon junction complex.** *Science* 2001, **293**:1832-1836.

47. Kataoka N, Yong J, Kim VN, Velazquez F, Perkinson RA, Wang F, Dreyfuss G: **Pre-mRNA splicing imprints mRNA in the nucleus with a novel RNA-binding protein that persists in the cytoplasm.** *Mol Cell* 2000, **6**:673-682.

48. Gehring NH, Neu-Yilik G, Schell T, Hentze MW, Kulozik AE: **Y14 and hUpf3b form an NMD-activating complex.** *Mol Cell* 2003, **11**:939-949.

49. Chamieh H, Ballut L, Bonneau F, Le Hir H: **NMD factors UPF2 and UPF3 bridge UPF1 to the exon junction complex and stimulate its RNA helicase activity.** *Nat Struct Mol Biol* 2008, **15**:85-93.

This study provides evidence that Upf2 and Upf3 stimulate ATPase and RNA helicase activities of Upf1 *in vitro* and that the EJC and Upf proteins can form a stable complex on RNA, consistent with the model that the EJC stimulates NMD by the recruitment of the Upf complex.

50. Sato H, Hosoda N, Maquat LE: **Efficiency of the pioneer round of translation affects the cellular site of nonsense-mediated mRNA decay.** *Mol Cell* 2008, **29**:255-262.

51. Zhang Z, Krainer AR: **Involvement of SR proteins in mRNA surveillance.** *Mol Cell* 2004, **16**:597-607.

52. Nagy E, Maquat LE: **A rule for termination-codon position within intron-containing genes: when nonsense affects RNA abundance.** *Trends Biochem Sci* 1998, **23**:198-199.

53. Le Hir H, Seraphin B: **EJCs at the heart of translational control.** *Cell* 2008, **133**:213-216.

54. Sanford JR, Gray NK, Beckmann K, Caceres JF: **A novel role for shuttling SR proteins in mRNA translation.** *Genes Dev* 2004, **18**:755-768.

55. Silva AL, Ribeiro P, Inacio A, Liebhaber SA, Romao L: **Proximity of the poly(A)-binding protein to a premature termination codon inhibits mammalian nonsense-mediated mRNA decay.** *RNA* 2008, **14**:563-576.

This study presents evidence that tethering PABPC downstream to the premature termination event can inhibit NMD in human cells, suggesting that PABPC can antagonize NMD. In addition, the authors report that IRES-mediated translation of an NMD-resistant AUG-proximal PTC-containing mRNA rendered the mRNA sensitive to NMD.

56. Cosson B, Couturier A, Chabelskaya S, Kiktev D, Inge-Vechtomov S, Philippe M, Zhouravleva G: **Poly(A)-binding protein acts in translation termination via eukaryotic release factor 3 interaction and does not influence [PSI(+)] propagation.** *Mol Cell Biol* 2002, **22**:3301-3315.

57. Czaplinski K, Ruiz-Echevarria MJ, Paushkin SV, Han X, Weng Y, Perlick HA, Dietz HC, Ter-Avanesyan MD, Peltz SW: **The surveillance complex interacts with the translation release factors to enhance termination and degrade aberrant mRNAs.** *Genes Dev* 1998, **12**:1665-1677.

58. Hoshino S, Hosoda N, Araki Y, Kobayashi T, Uchida N, Funakoshi Y, Katada T: **Novel function of the eukaryotic polypeptide-chain releasing factor 3 (eRF3/GSPT) in the mRNA degradation pathway.** *Biochemistry (Mosc)* 1999, **64**:1367-1372.

59. Hoshino S, Imai M, Kobayashi T, Uchida N, Katada T: **The eukaryotic polypeptide chain releasing factor (eRF3/GSPT) carrying the translation termination signal to the 3'-poly(A) tail of mRNA. Direct association of eRF3/GSPT with polyadenylate-binding protein.** *J Biol Chem* 1999, **274**:16677-16680.

60. Uchida N, Hoshino S, Imataka H, Sonenberg N, Katada T: **A novel role of the mammalian GSPT/eRF3 associating with poly(A)-binding protein in Cap/poly(A)-dependent translation.** *J Biol Chem* 2002, **277**:50286-50292.

61. Wang W, Czaplinski K, Rao Y, Peltz SW: **The role of Upf proteins in modulating the translation read-through of nonsense-containing transcripts.** *EMBO J* 2001, **20**:880-890.

62. Meaux S, van Hoof A, Baker KE: **Nonsense-mediated mRNA decay in yeast does not require PAB1 or a poly(A) tail.** *Mol Cell* 2008, **29**:134-140.

This paper reports that in yeast, the NMD machinery can discriminate PTC-containing mRNA from its cognate normal mRNA, even when the mRNA lacks a poly(A)-tail or when PABPC is depleted. This suggests that the poly(A)-tail and PABPC are not required for distinguishing NMD substrates from normal mRNA in yeast.

63. Ruiz-Echevarria MJ, Peltz SW: **The RNA binding protein Pub1 modulates the stability of transcripts containing upstream open reading frames.** *Cell* 2000, **101**:741-751.
64. Weil JE, Beemon KL: **A 3' UTR sequence stabilizes termination codons in the unspliced RNA of Rous sarcoma virus.** *RNA* 2006, **12**:102-110.
65. Fortes P, Inada T, Preiss T, Hentze MW, Mattaj JW, Sachs AB: **The yeast nuclear cap binding complex can interact with translation factor eIF4G and mediate translation initiation.** *Mol Cell* 2000, **6**:191-196.
66. Maquat LE: **Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics.** *Nat Rev Mol Cell Biol* 2004, **5**:89-99.
67. Ishigaki Y, Li X, Serin G, Maquat LE: **Evidence for a pioneer round of mRNA translation: mRNAs subject to nonsense-mediated decay in mammalian cells are bound by CBP80 and CBP20.** *Cell* 2001, **106**:607-617.
68. Chiu SY, Lejeune F, Ranganathan AC, Maquat LE: **The pioneer translation initiation complex is functionally distinct from but structurally overlaps with the steady-state translation initiation complex.** *Genes Dev* 2004, **18**:745-754.
69. Hosoda N, Kim YK, Lejeune F, Maquat LE: **CBP80 promotes interaction of Upf1 with Upf2 during nonsense-mediated mRNA decay in mammalian cells.** *Nat Struct Mol Biol* 2005, **12**:893-901.
70. Holbrook JA, Neu-Yilik G, Gehring NH, Kulozik AE, Hentze MW: **Internal ribosome entry sequence-mediated translation initiation triggers nonsense-mediated decay.** *EMBO Rep* 2006, **7**:722-726.
71. Woeller CF, Gaspari M, Isken O, Maquat LE: **NMD resulting from encephalomyocarditis virus IRES-directed translation initiation seems to be restricted to CBP80/20-bound mRNA.** *EMBO Rep* 2008, **9**:446-451.
72. Maderazo AB, Belk JP, He F, Jacobson A: **Nonsense-containing mRNAs that accumulate in the absence of a functional nonsense-mediated mRNA decay pathway are destabilized rapidly upon its restitution.** *Mol Cell Biol* 2003, **23**:842-851.
73. Zhang S, Welch EM, Hogan K, Brown AH, Peltz SW, Jacobson A: **Polysome-associated mRNAs are substrates for the nonsense-mediated mRNA decay pathway in *Saccharomyces cerevisiae*.** *RNA* 1997, **3**:234-244.
74. Gao Q, Das B, Sherman F, Maquat LE: **Cap-binding protein 1-mediated and eukaryotic translation initiation factor 4E-mediated pioneer rounds of translation in yeast.** *Proc Natl Acad Sci U S A* 2005, **102**:4258-4263.
75. Plant EP, Wang P, Jacobs JL, Dinman JD: **A programmed-1 ribosomal frameshift signal can function as a cis-acting mRNA destabilizing element.** *Nucleic Acids Res* 2004, **32**:784-790.
76. Baserga SJ, Benz EJ Jr: **Beta-globin nonsense mutation: deficient accumulation of mRNA occurs despite normal cytoplasmic stability.** *Proc Natl Acad Sci U S A* 1992, **89**:2935-2939.
77. Cheng J, Maquat LE: **Nonsense codons can reduce the abundance of nuclear mRNA without affecting the abundance of pre-mRNA or the half-life of cytoplasmic mRNA.** *Mol Cell Biol* 1993, **13**:1892-1902.
78. Lykke-Andersen J, Shu MD, Steitz JA: **Human Upf proteins target an mRNA for nonsense-mediated decay when bound downstream of a termination codon.** *Cell* 2000, **103**:1121-1131.
79. Cheng Z, Muhlrud D, Lim MK, Parker R, Song H: **Structural and functional insights into the human Upf1 helicase core.** *EMBO J* 2007, **26**:253-264.
80. Sheth U, Parker R: **Targeting of aberrant mRNAs to cytoplasmic processing bodies.** *Cell* 2006, **125**:1095-1109.
81. Kashima I, Yamashita A, Izumi N, Kataoka N, Morishita R, Hoshino S, Ohno M, Dreyfuss G, Ohno S: **Binding of a novel SMG-1-Upf1-eRF1-eRF3 complex (SURF) to the exon junction complex triggers Upf1 phosphorylation and nonsense-mediated mRNA decay.** *Genes Dev* 2006, **20**:355-367.
- This paper presents evidence that human Smg1 exists as part of a complex with Upf1 and translation release factors eRF1 and eRF3. Moreover, Smg1 interacts directly with Upf2, and Upf2 and Upf3 stimulate Smg1-mediated phosphorylation of Upf1.
82. Grimson A, O'Connor S, Newman CL, Anderson P: **SMG-1 is a phosphatidylinositol kinase-related protein kinase required for nonsense-mediated mRNA decay in *Caenorhabditis elegans*.** *Mol Cell Biol* 2004, **24**:7483-7490.
83. Anders KR, Grimson A, Anderson P: **SMG-5, required for *C. elegans* nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A.** *EMBO J* 2003, **22**:641-650.
84. Cali BM, Kuchma SL, Latham J, Anderson P: **smg-7 is required for mRNA surveillance in *Caenorhabditis elegans*.** *Genetics* 1999, **151**:605-616.
85. Chiu SY, Serin G, Ohara O, Maquat LE: **Characterization of human Smg5/7a: a protein with similarities to *Caenorhabditis elegans* SMG5 and SMG7 that functions in the dephosphorylation of Upf1.** *RNA* 2003, **9**:77-87.
86. Fukuhara N, Ebert J, Unterholzner L, Lindner D, Izaurralde E, Conti E: **SMG7 is a 14-3-3-like adaptor in the nonsense-mediated mRNA decay pathway.** *Mol Cell* 2005, **17**:537-547.
87. Ohnishi T, Yamashita A, Kashima I, Schell T, Anders KR, Grimson A, Hachiya T, Hentze MW, Anderson P, Ohno S: **Phosphorylation of hUPF1 induces formation of mRNA surveillance complexes containing hSMG-5 and hSMG-7.** *Mol Cell* 2003, **12**:1187-1200.
88. Page MF, Carr B, Anders KR, Grimson A, Anderson P: **SMG-2 is a phosphorylated protein required for mRNA surveillance in *Caenorhabditis elegans* and related to Upf1p of yeast.** *Mol Cell Biol* 1999, **19**:5943-5951.
89. Conti E, Izaurralde E: **Nonsense-mediated mRNA decay: molecular insights and mechanistic variations across species.** *Curr Opin Cell Biol* 2005, **17**:316-325.
90. Durand S, Cougot N, Mahuteau-Betzer F, Nguyen CH, Grierson DS, Bertrand E, Tazi J, Lejeune F: **Inhibition of nonsense-mediated mRNA decay (NMD) by a new chemical molecule reveals the dynamic of NMD factors in P-bodies.** *J Cell Biol* 2007, **178**:1145-1160.
- This study identifies a small hydrophobic indole derivative that leads to the accumulation of phospho-Upf1 and exclusion of Smg5 from P-bodies suggesting that the molecule inhibits NMD by interfering Upf1-Smg5 interactions in human cells.
91. Yamashita A, Ohnishi T, Kashima I, Taya Y, Ohno S: **Human SMG-1, a novel phosphatidylinositol 3-kinase-related protein kinase, associates with components of the mRNA surveillance complex and is involved in the regulation of nonsense-mediated mRNA decay.** *Genes Dev* 2001, **15**:2215-2228.
92. Gatfield D, Unterholzner L, Ciccarelli FD, Bork P, Izaurralde E: **Nonsense-mediated mRNA decay in *Drosophila*: at the intersection of the yeast and mammalian pathways.** *EMBO J* 2003, **22**:3960-3970.
93. Wang W, Cajigas LJ, Peltz SW, Wilkinson MF, Gonzalez CI: **Role for Upf2p phosphorylation in *Saccharomyces cerevisiae* nonsense-mediated mRNA decay.** *Mol Cell Biol* 2006, **26**:3390-3400.
94. Luke B, Azzalin CM, Hug N, Deplazes A, Peter M, Lingner J: ***Saccharomyces cerevisiae* Ebs1p is a putative ortholog of human Smg7 and promotes nonsense-mediated mRNA decay.** *Nucleic Acids Res* 2007, **35**:7688-7697.

95. Parker R, Sheth U: **P bodies and the control of mRNA translation and degradation.** *Mol Cell* 2007, **25**:635-646.
96. Muhlrads D, Parker R: **Recognition of yeast mRNAs as 'nonsense containing' leads to both inhibition of mRNA translation and mRNA degradation: implications for the control of mRNA decapping.** *Mol Biol Cell* 1999, **10**:3971-3978.
97. Anczukow O, Ware MD, Buisson M, Zetoune AB, Stoppa-Lyonnet D, Sinilnikova OM, Mazoyer S: **Does the nonsense-mediated mRNA decay mechanism prevent the synthesis of truncated BRCA1, CHK2, and p53 proteins?** *Hum Mutat* 2008, **29**:65-73.
98. You KT, Li LS, Kim NG, Kang HJ, Koh KH, Chwae YJ, Kim KM, Kim YK, Park SM, Jang SK *et al.*: **Selective translational repression of truncated proteins from frameshift mutation-derived mRNAs in tumors.** *PLoS Biol* 2007, **5**:e109.
99. Mendell JT, Medghalchi SM, Lake RG, Noensie EN, Dietz HC: **Novel Upf2p orthologues suggest a functional link between translation initiation and nonsense surveillance complexes.** *Mol Cell Biol* 2000, **20**:8944-8957.
100. Morris C, Wittmann J, Jack HM, Jalinot P: **Human INT6/eIF3e is required for nonsense-mediated mRNA decay.** *EMBO Rep* 2007, **8**:596-602.
101. Isken O, Kim YK, Hosoda N, Mayeur GL, Hershey JW, Maquat LE: **Upf1 phosphorylation triggers translational repression during nonsense-mediated mRNA decay.** *Cell* 2008, **133**:314-327.
The authors show that phosphorylated Upf1 interacts with subunits of eIF3, represses the translation of a reporter mRNA and inhibits the joining of 60S ribosomal subunit in rabbit reticulocyte lysates, suggesting a role for phospho-Upf1 in repressing translation. In addition, phospho-Upf1 preferentially coimmunoprecipitates with several mRNA decay factors suggesting that the phosphorylation status of Upf1 is important for the recruitment of mRNA decay factors to the NMD substrate.
102. Muhlrads D, Parker R: **Premature translational termination triggers mRNA decapping.** *Nature* 1994, **370**:578-581.
103. Eberle AB, Lykke-Andersen S, Muhlemann O, Jensen TH: **SMG6 promotes endonucleolytic cleavage of nonsense mRNA in human cells.** *Nat Struct Mol Biol* 2008, **16**(1):49-55.
This study shows evidence that endonucleolytic cleavage takes place in human NMD substrates. Moreover, human Smg6 shows endonuclease activity *in vitro* and when mutated in critical catalytic residues, Smg6 fails to support NMD, suggesting that Smg6 is an endonuclease in human NMD.
104. Gatfield D, Izaurralde E: **Nonsense-mediated messenger RNA decay is initiated by endonucleolytic cleavage in *Drosophila*.** *Nature* 2004, **429**:575-578.
105. Huntzinger E, Kashima I, Fauser M, Sauliere J, Izaurralde E: **SMG6 is the catalytic endonuclease that cleaves mRNAs containing nonsense codons in metazoan.** *RNA* 2008, **14**:2609-2617.
This study identifies Smg6 as the protein that mediates endonucleolytic cleavage of NMD substrates in *D. melanogaster*. The catalytic activity of Smg6 was shown to be important for NMD in both *D. melanogaster* and human cells.
106. Glavan F, Behm-Ansmant I, Izaurralde E, Conti E: **Structures of the PIN domains of SMG6 and SMG5 reveal a nuclease within the mRNA surveillance complex.** *EMBO J* 2006, **25**:5117-5125.