

Italy); Dr. R. Capildeo, Romford; Dr. W.J. Cumming, Manchester; Dr. M.L. Dattachaudhuri, Shawheath; Dr. C.J.K. Ellis, Poole; Dr. N. Fletcher, Liverpool; Dr. S.K. Gupta, Carmarthen; Dr. B.N. McLean, Truro.

## REFERENCES

- Hubble JP, Koller WC, Cutler NR, et al. Pramipexole in patients with early Parkinson's disease. *Clin Neuropharmacol* 1995;18:338–347.
- Molho ES, Factor SA, Weiner WJ, et al. The use of pramipexole, a novel dopamine (DA) agonist, in advanced Parkinson's disease. *J Neural Transm Suppl* 1995;45:225–230.
- Lieberman A, Ranhosky A, Korts D. Clinical evaluation of pramipexole in advanced Parkinson's disease: results of a double-blind, placebo-controlled, parallel-group study. *Neurology* 1997;49:162–168.
- Guttman M. Double-blind comparison of pramipexole and bromocriptine treatment with placebo in advanced Parkinson's disease. *International Pramipexole-Bromocriptine Study Group. Neurology* 1997;49:1060–1065.
- Parkinson Study Group. Safety and efficacy of pramipexole in early Parkinson disease. A randomized dose-ranging study. *JAMA* 1997;278:125–130.
- Shannon KM, Bennett JP, Friedman JH, et al. Efficacy of pramipexole, a novel dopamine agonist, as monotherapy in mild to moderate Parkinson's disease. *Neurology* 1997;49:724–728.
- Wermuth L. A double-blind, placebo-controlled, randomized, multi-center study of pramipexole in advanced Parkinson's disease. *Eur J Neurol* 1998;5:235–242.
- Künig G, Pogarell O, Möller JC, et al. Pramipexole, a nonergot dopamine agonist, is effective against rest tremor in intermediate to advanced Parkinson's disease. *Clin Neuropharmacol* 1999;22:301–305.
- Pinter MM, Pogarell O, Oertel WH. Efficacy, safety, and tolerance of the non-ergoline dopamine agonist pramipexole in the treatment of advanced Parkinson's disease: a double blind, placebo controlled, randomised, multicentre study. *J Neurol Neurosurg Psychiatry* 1999;66:436–441.
- Parkinson Study Group. Pramipexole vs. levodopa as initial treatment for Parkinson disease. *JAMA* 2000;284:1931–1938.
- Mizuno Y, Yanagisawa N, Kuno S, et al. Randomized, double-blind study of pramipexole with placebo and bromocriptine in advanced Parkinson's disease. *Mov Disord* 2003;18:1149–1156.
- Frucht S, Rogers JD, Greene PE, et al. Falling asleep at the wheel: motor vehicle mishaps in persons taking pramipexole and ropinirole. *Neurology* 1999;52:1908–1910.
- Pogarell O, Gasser T, van Hilten JJ, et al. Pramipexole in patients with Parkinson's disease and marked drug resistant tremor: a randomised, double blind, placebo controlled multicentre study. *J Neurol Neurosurg Psychiatry* 2002;72:713–720.
- Pinter MM, Rutgers AW, Hebenstreit E. An open-label, multicentre clinical trial to determine the levodopa dose-sparing capacity of pramipexole in patients with idiopathic Parkinson's disease. *J Neural Transm* 2000;107:1307–1323.
- Navan P, Findley LJ, Jeffs JA, et al. Double-blind, single-dose, cross-over study of the effects of pramipexole, pergolide, and placebo on rest tremor and UPDRS part III in Parkinson's disease. *Mov Disord* 2003;18:176–180.
- Corrigan MH, Denahan AQ, Wright CE, et al. Comparison of pramipexole, fluoxetine, and placebo in patients with major depression. *Depress Anxiety* 2000;11:58–65.
- Rektorova I, Rektor I, Bares M, et al. Pramipexole and pergolide in the treatment of depression in Parkinson's disease: a national multicentre prospective randomized study. *Eur J Neurol* 2003;10:399–406.
- Ondo WG, Dat Vuong K, Khan H, et al. Daytime sleepiness and other sleep disorders in Parkinson's disease. *Neurology* 2001;57:1392–1396.
- Hobson DE, Lang AE, Martin WR, et al. Excessive daytime sleepiness and sudden-onset sleep in Parkinson disease: a survey by the Canadian Movement Disorders Group. *JAMA* 2002;287:455–463.

## HIV Encephalitis Simulating Huntington's Disease

Jeffrey J. Sevigny, MD,<sup>1,2\*</sup>  
Steven S.M. Chin, MD, PhD,<sup>3</sup>

Yvonne Milewski, MD,<sup>3</sup>

Mark W. Albers, MD, PhD,<sup>1,4</sup> Marc L. Gordon, MD,<sup>5</sup>  
and Karen Marder, MD, MPH<sup>1,2,4,6</sup>

<sup>1</sup>Department of Neurology, Columbia University College of Physician and Surgeons, New York, New York, USA

<sup>2</sup>Gertrude H. Sergievsky Center, Columbia University College of Physician and Surgeons, New York, New York, USA

<sup>3</sup>Department of Pathology, Columbia University College of Physicians and Surgeons, New York, New York, USA

<sup>4</sup>Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University College of Physician and Surgeons, New York, New York, USA

<sup>5</sup>Departments of Neurology and Psychiatry, Long Island Jewish Medical Center, New Hyde Park, New York, USA

<sup>6</sup>Psychiatric Institute of New York, Columbia University College of Physician and Surgeons, New York, New York, USA

**Abstract:** Complications from human immunodeficiency virus (HIV)/acquired immune deficiency syndrome are notorious for mimicking other neurological diseases. We describe a case of HIV encephalitis presenting with the classic clinical features of Huntington's disease in a woman without known HIV risk factors or other clinical stigmata suggestive of immunosuppression. This case reminds us that HIV should be part of the differential diagnosis in unexplainable neurological diseases. © 2005 Movement Disorder Society

**Key words:** HIV; encephalitis; dementia; chorea

Pathological hyperkinetic movements in those infected with human immunodeficiency virus (HIV) are rare. Unilateral chorea or ballismus are the most reported forms, usually betraying a toxoplasmosis abscess in a contralateral basal ganglia.<sup>1–6</sup> We report a case of slowly progressive generalized chorea, behavioral changes, and

\*Correspondence to: Dr. Jeffrey J. Sevigny, The Neurological Institute, 710 W 168th Street, Box 160, New York, NY 10032.  
E-mail: js1112@columbia.edu

Received 9 June 2004; Revised 19 August 2004; Accepted 21 September 2004

Published online 8 February 2005 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/mds.20379

dementia in an otherwise "healthy" woman without any "known" risk factors, signs, or symptoms of HIV in whom the differential diagnosis included Huntington's disease (HD), prion disease, an autoimmune disorder, or a paraneoplastic syndrome, but an autopsy revealed HIV encephalitis (HIVE).

### CASE REPORT

A 49-year-old woman was referred to our Huntington's disease center with progressive dementia, chorea, and dystonia. She was an adopted child having no knowledge of her biological family's history. Her first husband had died 8 years earlier, and she was remarried. Her symptoms had started 6 years ago with insidious behavioral changes that included social withdrawal and emotional lability. Three years later, she underwent a multi-specialty evaluation for the subacute onset of fatigue; paresthesias, pain, and weakness in her arms and shoulders; and gait impairment. Based on a cervical spine magnetic resonance imaging (MRI) scan and electromyographic/nerve conduction velocity studies, she was diagnosed with cervical arthritis, C-7 radiculopathy, and median and ulnar neuropathies. One year later, she developed chorea of her arms and head as well as short-term memory loss. Further work-up revealed an iron-deficiency anemia; a monoclonal gammopathy (immunoglobulin [Ig] G, 2220; IgM, 1190); an erythrocyte sedimentation rate of 122; normal white blood cell count and ceruloplasmin, B12, rapid plasma reagin, Lyme antibody values; a normal electroencephalogram; cortical and central atrophy on MRI scan; and an unremarkable bone marrow biopsy.


On examination at the HD center (6 years after symptom onset), she could no longer function independently, was disoriented to place, and could not perform serial 7's, spell "WORLD" backward, add change, recall two of three items, or copy figures (Mini-Mental State Examination score, 22/30). She made paraphasic errors. Her ocular pursuits were slow and saccades were interrupted; strength and sensation were intact; tone was normal; rapid alternating movements were slow throughout; and, except for absent ankle jerks, reflexes were 1+ with absent Babinski signs. Chorea and intermittent dystonic posturing of her head and extremities were present. Ambulation required assistance. Cerebrospinal fluid (CSF) analyses revealed the following: white blood cell count, 18 (96% lymphocytes and 4 monocytes); red blood cell count, 156; protein, 50; glucose, 50; elevated 14-3-3 protein; and negative Venereal Disease Research Laboratory test. Lupus anticoagulant was present and anticardiolipin IgG was elevated. Cerebral angiogram findings were normal. Genetic testing for HD, Huntington's dis-

ease-like (HDL-2), and dentatorubropallidolusian atrophy was negative.

Over the ensuing year, she became nonambulatory and severely demented. A diagnostic right temporal lobe biopsy was performed. It revealed a chronic meningoencephalitis with microglial nodules. Granulomata, multinucleated giant cells, viral cytopathic inclusions, and spongiform changes were absent. Further CSF analyses revealed two oligoclonal bands; IgG synthesis rate of 163; negative polymerase chain reaction findings for Jakob-Creutzfeldt (JC) virus, human T-cell leukemia virus I/II, Herpes family, Arbo, Coxsackie, and enteroviruses; and absent measles and Lyme antibodies. Serum anti-neuronal nuclear antibody types 1 and 2 and Purkinje cell cytoplasmic antibody type 1 were absent. A repeat bone marrow biopsy revealed a mild erythroid and megakaryocytic hyperplasia and plasmacytosis consistent with a myelodysplastic syndrome or monoclonal gammopathy. Computed tomography of the chest, abdomen, and pelvis revealed multiple enlarged axillary and retroperitoneal lymph nodes. An MRI scan of her brain revealed significant atrophy and subtle periventricular white matter changes. An HIV test was contemplated, but the patient was transferred to hospice before her husband could be deemed her legal proxy to consent for the test.

She died 1 month later. Autopsy of the brain revealed frequent microglial nodules, many associated with multinucleated giant cells, predominately involving white matter and gray-white junction regions. Virtually all sampled cortical and subcortical brain regions were affected. An immunohistochemical stain for HIV p24 antigen positively stained the microglial nodules and associated multinucleated giant cells (Fig. 1). Other central nervous system findings included a marked diffuse astrogliosis of the caudate nucleus, putamen, claustrum, amygdala, and thalamus; a diffuse microgliosis; a marked loss of Purkinje cells; and marked vacuolar myelopathy. The rest of the autopsy showed bilateral bronchopneumonia, retroperitoneal lymphadenopathy with reactive plasmacytosis, and wasting of distal and proximal muscles. A CSF HIV RNA level obtained on a stored postmortem sample was log<sub>10</sub> 5.8. Plasma viral level and CD4 lymphocyte count were not obtained.

The results from the autopsy were discussed with her husband, who subsequently discovered from a friend that her previous husband had passed away from an unknown HIV-related complication. Our patient never shared this information with her family or her medical providers. Her husband tested HIV-negative, despite unprotected sex.



**FIG. 1.** Microglial nodule with multinucleated giant cells in internal capsule immunohistochemically stained for HIV p24 antigen. Scale bar = 50  $\mu$ m. [Color figure can be viewed in the online issue, which is available at [www.interscience.wiley.com](http://www.interscience.wiley.com).]

## DISCUSSION

This report is an unusual case of chronic, slowly progressive HIVE simulating the cardinal features of HD in a woman not known to be HIV infected. The clinical correlate of HIVE is HIV-associated dementia (HIVD),<sup>7</sup> which, like HD, is classified as a “subcortical” dementia and typically manifests with behavioral changes, cognitive impairment (e.g., slowed processing and executive function), and motor abnormalities. Phenotypic differences in the motor abnormalities can usually enable one to discriminate between the two diseases.

One commonly cited study suggests that movement disorders among acquired immune deficiency syndrome (AIDS) patients are rare.<sup>1</sup> However, extrapyramidal signs such as bradykinesia and rigidity are common neurological abnormalities associated with AIDS, particularly among those with HIVD.<sup>8</sup> What is uncommon among those with AIDS are hyperkinetic movements such as chorea. When present, it is usually unilateral and associated with an underlying space-occupying lesion such as toxoplasmosis.<sup>1–6</sup>

Three cases of generalized chorea attributed to HIVE have been reported. In the only biopsy proven case, a subject with known HIV and a CD4 count of 383 cells/mm<sup>3</sup> presented with rapidly progressive encephalopathy and generalized chorea.<sup>9</sup> In the two presumed cases of HIVE, 1 subject with AIDS and a CD4 cell count of 50 cells/mm<sup>3</sup>, developed rapid onset of chorea that improved with zidovudine (AZT),<sup>10</sup> whereas the other, who was previously undiagnosed with HIV and had a CD4 cell count of 42 cells/mm<sup>3</sup>, presented with a 6-month

progressive course of chorea, personality changes, and cognitive impairment and also improved with AZT.<sup>11</sup>

HIV is considered “the great mimicker” in contemporary medicine. Despite advances in antiretroviral therapy, the central nervous system is affected, directly or indirectly, by HIV in over 70% of AIDS patients, and the hallmark pathological features of HIVE, namely microglial nodules with multinucleated giant cells, perivascular infiltrates, and astrogliosis, can be found in up to 60% in autopsy series.<sup>12</sup> HIVE can affect the entire brain, but for reasons not yet fully elucidated, it has a predilection for the subcortical structures, particularly the globus pallidus.<sup>13</sup> In our patient, the basal ganglia, including caudate, putamen, globus pallidus, and claustrum, showed a diffuse astrogliosis and numerous microglial nodules. In untreated patients, the dementia rapidly progresses and death usually ensues within months.

The ultimate diagnosis of HIVE was overlooked for several reasons: its unusual presentation and the long duration of illness; the lack of identifiable or reported HIV risk factors; and the absence of laboratory (e.g., leukopenia), clinical (opportunistic infections), or imaging (e.g., HIV-associated leukopathy) abnormalities suggestive of HIV infection. Therefore, the leading diagnoses considered reflected those likely to manifest with generalized chorea and dementia, including Huntington’s disease, dentatorubropallidoluysian syndrome, prion disease, autoimmune disorders, or paraneoplastic syndromes. Further complicating the picture were the presence of “red herrings” on diagnostic testing. The lupus anticoagulant and elevated anticardiolipin IgG antibodies suggested an autoimmune disorder, while the elevated CSF 14-3-3 protein suggested Jakob–Creutzfeldt disease, although antiphospholipid antibody syndrome and elevated 14-3-3 protein have been reported in HIV.<sup>14–16</sup>

Diagnosis and treatment with antiretroviral therapy might have prevented progression and indeed reversed the symptoms, as reported in the two presumed cases noted above and further suggested by recent studies on HIVD.<sup>17,18</sup> This case reminds us that HIV testing is warranted in unexplained neurological disease, irrespective of seemingly mitigating historical or clinical factors.

## REFERENCES

1. Nath A, Jankovic J, Pettigrew LC. Movement disorders and AIDS. *Neurology* 1987;37:37–41.
2. Nath A, Hobson DE, Russell A. Movement disorders with cerebral toxoplasmosis and AIDS. *Mov Disord* 1993;8:107–112.
3. Navia BA, Petito CK, Gold JW, Cho ES, Jordan BD, Price RW. Cerebral toxoplasmosis complicating the acquired immune deficiency syndrome: clinical and neuropathological findings in 27 patients. *Ann Neurol* 1986;19:224–238.
4. Sanchez-Ramos JR, Factor SA, Weiner WJ, Marquez J. Hemichorea-hemiballismus associated with acquired immune deficiency



syndrome and cerebral toxoplasmosis. *Mov Disord* 1989;4:266–273.

5. Pestre P, Milandre L, Farnarier P, Gallais H. [Hemichorea in acquired immunodeficiency syndrome. Toxoplasmosis abscess in the striatum]. *Rev Neurol (Paris)* 1991;147:833–837.
6. Maggi P, de Mari M, De Blasi R, et al. Chorea in acquired immune deficiency syndrome patients with cerebral toxoplasmosis. *Mov Disord* 1996;11:434–436.
7. Wiley CA, Achim C. Human immunodeficiency virus encephalitis is the pathological correlate of dementia in acquired immunodeficiency syndrome. *Ann Neurol* 1994;36:673–676.
8. Marder K, Liu X, Stern Y, et al. Neurologic signs and symptoms in a cohort of homosexual men followed for 4.5 years. *Neurology* 1995;4:261–267.
9. Gallo BV, Shulman LM, Weiner WJ, Petit CK, Berger JR. HIV encephalitis presenting with severe generalized chorea. *Neurology* 1996;46:1163–1165.
10. Piccolo I, Causarano R, Sterzi R, et al. Chorea in patients with AIDS. *Acta Neurol Scand* 1999;100:332–336.
11. Pardo J, Marcos A, Bhathal H, Castro M, Varea de Seijas E. Chorea as a form of presentation of human immunodeficiency virus-associated dementia complex. *Neurology* 1998;50:568–569.
12. Neuenburg JK, Brodt HR, Herndier BG, et al. HIV-related neuropathology, 1985 to 1999: rising prevalence of HIV encephalopathy in the era of highly active antiretroviral therapy. *J Acquir Immune Defic Syndr* 2002;31:171–177.
13. Brew BJ, Rosenblum M, Cronin K, Price RW. AIDS dementia complex and HIV-1 brain infection: clinical-virological correlations. *Ann Neurol* 1995;38:563–570.
14. Asherson RA, Shoenfeld Y. Human immunodeficiency virus infection, antiphospholipid antibodies, and the antiphospholipid syndrome. *J Rheumatol* 2003;30:214–219.
15. Leder AN, Flansbaum B, Zandman-Goddard G, Asherson R, Shoenfeld Y. Antiphospholipid syndrome induced by HIV. *Lupus* 2001;10:370–374.
16. Wakabayashi H, Yano M, Tachikawa N, Oka S, Maeda M, Kido H. Increased concentrations of 14-3-3 epsilon, gamma and zeta isoforms in cerebrospinal fluid of AIDS patients with neuronal destruction. *Clin Chim Acta* 2001;312:97–105.
17. Sacktor NC, Lyles RH, Skolasky RL, et al. Combination antiretroviral therapy improves psychomotor speed performance in HIV-seropositive homosexual men. Multicenter AIDS Cohort Study (MACS). *Neurology* 1999;52:1640–1647.
18. Tozzi V, Balestra P, Galgani S, et al. Positive and sustained effects of highly active antiretroviral therapy on HIV-1-associated neurocognitive impairment. *Aids* 1999;13:1889–1897.

## Parkinsonism in the Course of HTLV-I-Associated Myelopathy

Marzia Puccioni-Sohler, MD, PhD<sup>1–3\*</sup>

Regina Papais-Alvarenga, MD, PhD<sup>1</sup>

Priscila Mesquita de Souza,<sup>1</sup> Silvia Curi de França,<sup>1</sup>

Reizer Reis Gonçalves,<sup>2,3</sup>

and Steven Jacobson, PhD<sup>4</sup>

<sup>1</sup>Neurology Service, ESP/Hospital Universitário Gaffree Guinle (HUGG/UniRio), University of Rio de Janeiro, Rio de Janeiro, Brazil; <sup>2</sup>Cerebrospinal Fluid Laboratory/Hospital Universitário Clementino Fraga Filho (HUCFF/UFRJ), Federal University of Rio de Janeiro, Rio de Janeiro, Brazil; <sup>3</sup>Cerebrospinal Fluid Neurolife Laboratory, Rio de Janeiro, Brazil; <sup>4</sup>Neuroimmunology Branch, National Institutes of Health, Bethesda, Maryland, USA



**Abstract:** Parkinsonian syndromes may represent a complication of viral infection. Human T cell lymphotropic virus I (HTLV-I) is a cause of a chronic myelopathy in which encephalic involvement has been also found. We report on the case of a 60-year-old man with HTLV-I-associated myelopathy, complicated with bradykinesia, resting tremor, and cogwheel rigidity. These findings suggest that parkinsonian features may represent a neurological disorder associated with HTLV-I infection. © 2005 Movement Disorder Society

**Key words:** Parkinson syndrome; HAM/TSP; HTLV-I; CSF

Parkinson's disease (PD) is a progressive disease of the extrapyramidal system, characterized by degeneration of pigmented neurons in substantia nigra and other pigmented brainstem nuclei with dopamine deficiency transmitter in basal ganglia.<sup>1</sup> The syndrome includes bradykinesia, cogwheel rigidity, postural instability, and resting tremor. The most important problem is to differentiate PD from other parkinsonian syndromes related to other causes. Postencephalitic parkinsonism has been reported after viral encephalitis.<sup>2</sup>

This work was presented at the 11th International Conference on Human Retrovirology: HTLV and Related Viruses, June 9–12, 2003, San Francisco, CA.

This article includes Supplementary Video, available online at <http://interscience.wiley.com/jpages/0885-3185/suppmat>.

\*Correspondence to: Dr. M. Puccioni-Sohler, Rua de Zenove de Fevereiro 185/705, 2280-030 Rio de Janeiro, RJ, Brazil.

E-mail: [mpuccioni@hucff.ufrj.br](mailto:mpuccioni@hucff.ufrj.br)

Received 25 March 2004; Revised 24 August 2004; Accepted 22 September 2004

Published online 23 February 2005 in Wiley InterScience ([www.interscience.wiley.com](http://www.interscience.wiley.com)). DOI: 10.1002/mds.20402