



Effect of vitamin D and β -sitosterol on immune function of macrophages

Lini Alappat, Michael Valerio, Atif B. Awad *

Department of Exercise and Nutrition Sciences, University at Buffalo, Buffalo, NY 14214, United States

ARTICLE INFO

Article history:

Received 27 January 2010

Received in revised form 28 June 2010

Accepted 5 August 2010

Keywords:

Cytokines

Cell proliferation

Chemokines

Nitric oxide

NF- κ B

p65

ABSTRACT

Among the many functions of vitamin D (VD) is its role in the immunomodulation of macrophage. As VD deficiency is a wide-spread nutritional problem, there is a tendency for the public to overdose with vitamin D supplementation which can result in hypercalcemia and several associated disorders. The present study was designed to investigate the possibility that combining low doses of vitamin D with β -sitosterol (SIT), a common phytosterol in the diet without toxicity, enhances the efficacy of the vitamin. Murine macrophages were stimulated with LPS and supplemented with VD3 (80 nM) and SIT (8 μ M) for 24 hr and examined for cell proliferation, release of nitric oxide (NO) and cytokines and the activation of NF κ B. SIT (8 μ M) was found to reduce cell proliferation by 62% while VD3 was found to be not effective. In combination, SIT and VD3 reduced cell proliferation by 75%. The amount of NO released, as influenced by 8 μ M SIT or 80 nM VD3 treatments, was not significantly different from control. Combining SIT and VD3, resulted in a 220% greater increase in NO release compared to control. The SIT + VD3 treatment brought about significant increase in all the cytokine release, regardless of whether they were pro- or anti-inflammatory. The effects were either additive or synergistic. We conclude that SIT enhances the action of VD3 on the immune function of macrophages which could be beneficial to vitamin D deficient individuals and to those with autoimmune diseases such as multiple sclerosis.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

The most well established role of vitamin D (VD) is its role in calcium absorption. It is essential for preserving the calcium-phosphate balance in the body and thus it is vital for bone health [1–3]. Besides calcium homeostasis, VD has other functions in the human body, including its role in immune [4] and muscle function [5]. In addition, many regulatory proteins involved in cell proliferation, cell differentiation, and apoptosis are modulated by VD [1,6,7].

Nutrient deficiency can result from inadequate intake, poor absorption, added need, or increased excretion [8]. Usually vitamin D deficiency can occur when there is lower than recommended intake levels for prolonged time, limited sunlight exposure, inability of the kidneys to produce the active form of VD, or impaired absorption of VD from the digestive tract [8]. Consumption of natural food sources alone may not be enough to meet the daily dietary requirements of VD. Including VD fortified foods in their diet and exposure to sunlight are effective ways to get sufficient VD [8]. However, some people like those with disabilities, lactose intolerance, vegetarians, breast feeding mothers, or the elderly have to resort to dietary supplements to meet daily requirements [8].

Vitamin D toxicity is not likely to result from too much sunlight exposure or consumption of fortified foods [8], but excessive intake of supplements can present with many health problems [8,9]. Excessive intake of vitamin D supplements can increase serum calcium levels resulting in kidney stones, heart diseases or impaired mental status [10–12].

Although enough scientific evidence is not available to recommend the exact upper level of toxicity for vitamin D, there is no difference in opinion about the toxicity of vitamin D in excess amounts. Efforts are focused on finding a safe alternative that can augment the effects of VD. A lot of interest in this approach is being currently investigated by several drug manufacturers. Alternatively, the combination of low level of vitamin with other dietary molecules to increase its efficacy without the hypercalcemic side-effect offers an attractive solution.

One such molecule tested here is the β -sitosterol (SIT), the most common plant sterol or phytosterol, in the diet. Beside structural similarities between VD and SIT they each play role in immunomodulation [13,14]. Phytosterols are naturally found in a wide variety of plants and in good amounts in legumes such as peanuts [15] and oil seeds such as sesame seeds and soybeans [16]. Consumption of gram quantities of phytosterols has no side-effects [17]. Hence, it would be of great interest to see if SIT would act additively or synergistically with vitamin D to enhance its effect in lower doses on macrophage immune function. Macrophages produce nitric oxide (NO) and cytokines that play major roles in inflammation.

* Corresponding author. 15 Farber Hall, University at Buffalo, Buffalo, NY 14214, United States. Tel.: +1 716 829 5660; fax: +1 716 829 3700.

E-mail address: awad@buffalo.edu (A.B. Awad).

2. Materials and methods

2.1. Materials

Dulbecco's modified Eagle medium (DMEM) without phenol red, sulfanilamide, N-1-naphthylethylenediamine dehydrochloride, lipopolysaccharide (LPS) and β -sitosterol (SIT) were purchased from Sigma (St. Louis, MO). Vitamin D3 (VD3) or calcitriol was generously donated by SAFC Pharma, Inc. (Madison, WI); WST-8 reagent was obtained from BioVision (Mountain View, CA). ELISA^{MAX} deluxe kits for mouse interleukin-10 (IL-10), mouse monocyte chemoattractant protein (MCP-1), mouse interleukin-6 (IL-6) and mouse tumor necrosis factor – alpha (TNF- α) were purchased from BioLegend (San Diego, CA). NF κ B p65 profiler Kit was purchased from Active Motif (Carlsbad, CA). Sodium pyruvate, L-glutamine, Penicillin – Streptomycin and phosphate saline buffer (D-PBS) were from GIBCO (Carlsbad, CA). Other chemicals were obtained either from VWR or Fischer Scientific.

2.2. Methods

2.2.1. Cell culture

Murine J774A.1 macrophage cells (ATCC) were maintained in DMEM supplemented with 4 g/L glucose, 1.5 g/L sodium bicarbonate, 10 ml/L Pen-Strep, 10 ml/L sodium pyruvate and 10% fetal bovine serum.

2.2.2. Cell proliferation measurement

J774A.1 cells (5×10^5 cells/250 μ l) were seeded in a 96-well tissue culture plate and stimulated with LPS to a final concentration of 100 ng/ml [18]. To examine the effect of VD3 and SIT on cell proliferation, cells were incubated with different concentrations of these agents at the same time cells were stimulated with LPS. After 24 hours of incubation, the conditioned media were removed and 10 μ l of WST-8 reagent in 90 μ l media was added to all wells and incubated for 30 minutes to 4 hours under standard culture conditions according to the manufacture instructions. During that time the tetrazolium salt of WST-8 is cleaved to form formazan by mitochondrial dehydrogenase. The amount of the formazan produced was quantified by measuring the absorbance at 450 nm which is directly proportional to the number of viable cells. The reference wavelength used was 650 nm.

2.2.3. Nitric oxide measurement

LPS-stimulated cells were incubated with SIT, VD3 or both and incubated for at 37 °C in a 5% CO₂ – 95% air humidified atmosphere for 24 hours. Untreated cells were used as controls. Nitrite accumulation in the conditioned media was measured using the Griess assay [19].

2.2.4. Cytokine release measurement

Cells were treated either with SIT, VD3 or both and the experiment was performed as above. After 24 h incubation, conditioned media were collected from the 96-well plates and stored at -80 °C until cytokine assays were performed. ELISA^{MAX} Deluxe kits were used for measuring TNF- α , IFN- γ , IL-6, IL-10 and MCP-1 released from the treated cells. Standard curves were generated as per the instructions of the manufacturer to read the concentration of each cytokine released. The amount of cytokine released was expressed as percentage of control.

2.2.5. Measurement of phosphorylated and total NF κ B

Cells were seeded in 96-well plates and grown to about 80% confluence. Then they were treated with LPS and the test agents, SIT, VD3, either alone or in combination. After 24 h of incubation, the cells were fixed by replacing the growth medium with formaldehyde in PBS. The amount of phosphorylated NF κ B at Serine 468, Serine 536

and total NF κ B in the experimental samples were determined using NF κ B p65 profiler kit. The relative number of cells in each well was then determined according to the manufacturer's instructions.

2.2.6. Statistical analysis

The values are expressed as means \pm SEM and presented as percentages of control. Analysis of variance (ANOVA) followed by the student Newman-Keuls post-hoc test were utilized to examine the significance between groups. Differences between the means were considered to be statistically significant at a p-value <0.05. ProStat software (Pearl River, NY) was used to analyze the data.

3. Results

3.1. Effect of VD3 and SIT on cell proliferation of macrophages

The LPS-treated macrophages were incubated with respective treatments for 24 hours at 37 °C and 5% CO₂. Fig. 1 demonstrates the results of the first set of experiments conducted to identify the appropriate concentrations of the sterols used for subsequent experiments. The values are expressed as percentage of control.

VD3 treatments had no pronounced effects on cell proliferation. Here also, higher concentration of VD3 (160 nM) showed decrease in cell proliferation (by 45%) although not significantly different from control. Fig. 1 illustrates that with VD3 (80 nM), there was only a smaller reduction in cell proliferation (20%). Neither of the VD3 treatments was significantly different from each other.

SIT (1 μ M) treatment resulted in 59% decrease in cell proliferation while SIT (8 μ M) brought it down to 63% (Fig. 1). The decrease in cell proliferation with both the SIT treatments was found to be significantly different from control. The two SIT treatments did not significantly different from each other.

Since the VD3 at two concentrations studied were not different, we decided to use VD3 (80 nM) for subsequent experiments based on published literature [13,20]. In the case of SIT, although insignificant, SIT at 8 μ M was able to induce more pronounced decrease in cell proliferation compared to 1 μ M. In addition, SIT (8 μ M) corresponds to serum SIT level associated with the consumption of the western mixed diet. Accordingly, SIT (8 μ M) was used in further experiments.

Fig. 1 demonstrates the results of a set of experiments performed to examine the effect of the combined treatment (SIT + VD3) on cell proliferation. The treatment groups included control (no SIT or VD3),

Effect of VD3 and SIT on cell proliferation of J774a.1 cells




Fig. 1. Effect of VD3 and SIT treatments on cell proliferation of LPS-stimulated J774A.1 cells. J774A.1 cells (5×10^5 cells/250 μ l) were seeded in a 96-well tissue culture plate. LPS (100 ng/ml) was added to each well and incubated with SIT, VD3, both or none (control). After 24 h incubation, conditioned media was removed from each well and cells were subjected to WST-8 cell proliferation assay. Values are expressed as percentage of control. Columns represent average of three observations and bars represent SEM. Bars with different letters indicate significant differences ($p < 0.05$) as evidenced by student Newman-Keuls post-hoc test. Abbreviations: SIT = β -sitosterol, VD3 = vitamin D3.

SIT (8 μ M), VD3 (80 nM) and SIT + VD3. Cell proliferation is expressed as the percentage of the control, with the control representing 100%. VD3 treatment did not reveal any significant change in proliferation from control, while SIT treatment decreased it by 63%. The SIT treated cells showed significant inhibition of cell proliferation compared to VD3 treated cells.

Fig. 1 also illustrates that the SIT + VD3 treatment exhibited the greatest effect, as it led to a significant 76% decrease in cell proliferation. SIT treated cells showed no significant difference in cell proliferation from SIT + VD3 treated cells. The reverse effect was noticed between VD3 and SIT + VD3 treated cells. There was significant difference between these two groups. Overall, the treatments with SIT, alone or in combination with VD3, proved to be effective in reducing cell proliferation.

3.2. Effect of VD3 and SIT on NO release from J774A.1 macrophages

Using similar design to that used in cell proliferation experiments, NO release from treated macrophages was examined (Fig. 2). VD3 (80 nM) treatment had an effect on NO released from macrophages. The amount of NO released, as influenced by SIT treatment, was significantly different from control. Combining these two treatments also yielded significant results. SIT + VD3 treatment resulted in more than 220% increase in NO release above the control. This effect noticed with the combined treatment was the summation of the individual effects i.e. the effect was additive.

3.3. Effect of VD3 and SIT on cytokine release

The analyzed cytokines included IL-10, IL-6, TNF- α , MCP-1 and IFN- γ in the conditioned media of treated cells:

IL-10

IL-10 is an anti-inflammatory cytokine. Fig. 2 illustrates the release of IL-10 from the macrophages as influenced by treatments. VD3 treatment increased IL-10 production from J774A.1 cells by 32%. SIT treatment was found to significantly




Fig. 2. Effect of VD3 and SIT treatments on nitric oxide and cytokine release from LPS-stimulated J774A.1 cells. J774A.1 cells (5×10^5 cells/250 μ l) were seeded in a 96-well tissue culture plate. LPS (100 ng/ml) was added to each well and incubated with SIT, VD3, both or none (control). After 24 h incubation, conditioned media was collected from each well and quantified for NO by Griess assay. The conditioned media collected after 24 h incubation was also quantified for the respective cytokines by ELISA kit. Values are expressed as percentage of control. Columns represent average of three observations and bars represent SEM. Bars with different letters within an inflammatory marker indicate significant differences ($p < 0.05$) as evidenced by student Newman-Keuls post-hoc test. Abbreviations: SIT = β -sitosterol, VD3 = vitamin D3.

increase the IL-10 release by 114% from macrophages. The SIT + VD3 treatment resulted in a 230% increase in IL-10 production compared to control. The amount of IL-10 produced on treating with SIT + VD3 was also significantly higher than the summation of individual treatment groups suggesting a synergistic effect.

IL-6

With regards to IL-6 (Fig. 2), VD3 treatment did not produce any noticeable change in IL-6 release from macrophages. On the other hand, SIT treatment produced significantly higher amounts of this Th2 cytokine compared to the control. The SIT + VD3 treatment significantly increase IL-6 production compared to control and other treatment groups. Compared to the control, the increase in IL-6 for SIT + VD3 group was almost 3 times higher, suggesting a synergistic effect.

TNF- α

TNF- α is a pro-inflammatory cytokine produced by macrophages upon stimulation by foreign substances. TNF- α released from macrophages treated with VD3 was not significantly different from control (Fig. 2). It can be observed that the treatment with SIT produced significant increase (by 150%) in TNF- α production from J774A.1 cells. The SIT + VD3 treatment led to a 4 times increase in TNF- α production compared to control. With SIT + VD3 treatment the increase in TNF- α production was 2.5 times higher than SIT alone and 4 times more than VD3 alone.

MCP-1

MCP-1 is a chemokine that recruits monocytes to the site of injury or infection. Fig. 2 illustrates that the effect of SIT and VD3 on MCP-1 release followed the same pattern as in the case of TNF- α . VD3 produced significant rise in MCP-1 release from the macrophages. Treatment with SIT led to a noticeable increase in MCP-1 production. The combination treatment produced a substantial increase in MCP-1 release. Like TNF- α , the SIT + VD3 treatment produced a 4-fold increase in MCP-1 production compared to control, suggesting synergistic effect.

IFN- γ

IFN- γ is a cytokine needed for innate and adaptive immunity. Fig. 2 shows that IFN- γ release was markedly different for all the treatments. This followed the same pattern as that for MCP-1 and TNF- α . The VD3 treatment had no effect on IFN- γ release from the macrophages. IFN- γ was increased significantly with SIT treatment by 80%. The SIT + VD3 produced a sizeable increase (4 times) in IFN- γ release from macrophages with respect to control. The combined effect was also significantly different from those of SIT and VD3 treatment cells.

3.4. Effect of VD3 and SIT on total and phosphorylated NF κ B

By determining the phosphorylation of NF κ B on Ser536, one would be able to examine the role of NF κ B pathway in the observed results. Since the kit we used can also measure Ser 468 phosphorylation by activating NF κ B by another pathway with cytokines we decided to examine Ser 468 phosphorylation, as well. The experiment was repeated with similar results. Fig. 3 demonstrates the results in one of these experiments. There was a decrease in Ser 536 phosphorylation with both SIT and VD3 + SIT treatments. VD3 treatment also showed a slight decrease in Ser 536 phosphorylation. On the other hand, a noticeable increase in Ser 468 phosphorylation was found with SIT and SIT + VD3 treatments, while VD3 decreased it (Fig. 4). In Fig. 5, the total cellular NF κ B remained relatively unchanged except for SIT + VD3 treatment where a slight increase was noticed. Similar results were obtained when the experiment was repeated.




Fig. 3. Effect of VD3 and SIT treatments on Ser 536 phosphorylation. J774A.1 cells (3×10^5 cells/100 μ l) were seeded in a 96-well tissue culture plate. LPS (100 ng/ml) was added to each well and incubated with SIT, VD3, both or none (control). After 24 h incubation, the cells were subjected to FACE assay and values shown have been corrected for cell number. Abbreviations: SIT = β -sitosterol, VD3 = vitamin D3.




Fig. 5. Effect of VD3 and SIT treatments on total NF κ B. J774A.1 cells (3×10^5 cells/100 μ l) were seeded in a 96-well tissue culture plate. LPS (100 ng/ml) was added to each well and incubated with SIT, VD3, both or none (control). After 24 h incubation, the cells were subjected to FACE assay and values shown have been corrected for cell number. Abbreviations: SIT = β -sitosterol, VD3 = vitamin D3.

4. Discussion

The aim of the present study was conducted to examine the potential modulation of vitamin D3 action on macrophages with phytosterols. Phytosterols and VD3 are both sterols and thus they have structural similarity. We have demonstrated that the SIT, the main phytosterol in our diet, augments the action of VD3 on the immune function of macrophages. The markers of immune function we used in the study were cell proliferation, cytokine and nitric oxide released from LPS-challenged macrophages. To the best of our knowledge this is the first study that has examined the combined effect of SIT and VD3 on the immune function of macrophages.

The present study used a different protocol to activate the cells with LPS. Most studies pre-incubate the macrophages with the test agents and then treat them with LPS [13,21]. In this study, we incubated cells simultaneously with LPS and sterols for 24 hours to simulate the *in vivo* setting.

VD3 is an immuno-modulator [22,23] and could be produced by peritoneal macrophages during inflammation [13,24]. VD3 has been shown to down regulate the production of pro-inflammatory cytokines and up-regulate the anti-inflammatory ones [23,25–33]. VD3 exerts its effects on immune function by targeting NF κ B [32,34]. VD3 has been shown to decrease the expression of TNF- α mRNA and cytokine levels in LPS-stimulated murine macrophages [35,36]. The underlying mechanism behind this down regulation is thought to be due to the stabilization of I κ B mRNA and reducing the I κ B phosphorylation. This reduces the nuclear NF κ B-p65 complex content and increases the cytoplasmic fraction.




Fig. 4. Effect of VD3 and SIT treatments on Ser 468 phosphorylation. J774A.1 cells (3×10^5 cells/100 μ l) were seeded in a 96-well tissue culture plate. LPS (100 ng/ml) was added to each well and incubated with SIT, VD3, both or none (control). After 24 h incubation, the cells were subjected to FACE assay and values shown have been corrected for cell number. Abbreviations: SIT = β -sitosterol, VD3 = vitamin D3.

On the other hand, Zemel *et al* showed that calcitriol up-regulated inflammatory cytokine expression in RAW 264.7 cells cultured alone and co-cultured with differentiated adipocytes [37]. The cytokines so up-regulated include TNF- α , IL-6 and MCP-1 and is thought to be mediated by calcitriol stimulated Ca^{2+} signaling and diminished mitochondrial uncoupling [37]. Hakim *et al* also showed that VD3 treatment up regulated TNF- α and cytokine response both *in vitro* and *in vivo* [38]. Results from another study showed that VD3 stimulation did not affect the TNF- α production while increased IFN- γ , IL-10 and IL-6 production from PBMC of Crohn's disease patients [39]. Numerous studies have shown VD3 to be a potential anti-inflammatory agent by suppressing NF κ B activity and other transcription factors [28,40–42]. Contrary to these findings, our study did not find a change in cytokine release from the macrophages with the VD3 treatment. Both the pro- and anti-inflammatory cytokines were found to be unchanged compared to vehicle control. Only MCP-1 and IL-10 were found to be significantly increased. Such a sustained response is especially useful in bacterial and viral infections where a prolonged Th1 cell response is needed to destroy the pathogens. Increased MCP-1 response is ideal for mass recruitment of monocytes to the area of inflammation. Increased IL-10 is beneficial for multiple sclerosis patients where they experience a decreased IL-10 production [43]. Our data demonstrate that calcitriol sustains the cytokine response as opposed to down regulation. The unaltered response we noticed might be due to the different cell type [37], incubation period [39], VD3 dosage [13,21,37] and differentiation status of cells [38] used in our experiment.

NO is an important defense mechanism in the body against the invading micro-organisms [44,45]. Its synthesis is mediated by nitric oxide synthases (NOS). Among the NOS, inducible NOS (iNOS) can be induced in macrophages by stimulation with LPS resulting in large amounts of NO being produced [46,47]. The NO released can decrease cell proliferation and affect cytokine release [46,47].

Calcitriol has been shown to inhibit the iNOS mRNA expression and thereby decrease NO release [21,48,49]. On the contrary, our study did not show the VD3 treatment to decrease the NO release. The sustained NO release as reported in our study is thought to be due to prolonged iNOS expression and resultant NO buildup. The unaltered NO response goes hand in hand with the pattern of IFN- γ and TNF- α release seen with VD3 treatment. Cytokines such as IFN- γ and TNF- α have been shown to activate the macrophages to produce reactive nitrogen species to combat bacterial infection [50] and our study further corroborate this evidence. Another study by Rockett *et al* also showed that VD3 could enhance iNOS mRNA expression leading to restrained bacterial growth [51].

SIT also has immuno-modulatory properties [52–56]. Several *in vitro* studies have shown SIT to reduce TNF- α and IL-6 production

[52,57,58]. On the other hand, one study has shown SIT to have no effect on IL-10, IL-4 and IFN- γ levels [59]. SIT has been shown to preferentially induce Th1 cytokines like IFN- γ compared to Th2 cytokines [60]. Th1-type cytokines mediated cellular response is important for the host to defend against viral and bacterial pathogens and curbing infection [52]. The Th2-type cytokines (IL-4, IL-6, and IL-10) are engaged in B-lymphocyte differentiation limits the damage caused by micro-organisms [52]. The inflammatory response is mediated by the intrinsic balance between Th1 and Th2 cytokines. The increased response of Th1 cytokines is responsible in eliminating pathogens [52] while increased Th2 response is beneficial in autoimmune diseases like multiple sclerosis where there is a reduced IL-10 production [43]. SIT and its glycoside have been shown to increase IFN- γ secretion in allergic individuals [61]. Our data also show SIT to significantly increase the Th1 cytokine release from macrophages. SIT is shown to have no effect or to inhibit the release of Th2 cytokines [52,61]. We found SIT to increase Th2 cytokines contrary to what has been reported. Although not tested, preliminary analysis showed SIT to have a no preference for Th1 or Th2 cytokine stimulation as earlier reported [52].

SIT has been shown to decrease iNOS levels in PMA-stimulated RAW 264.7 cells [62]. However, the same study also pointed out that NO₂ production increased from 6–24 h after PMA stimulation and a decrease in the NO production with SIT was noticed after 24 hours [62]. However the decrease in NO production was noticed only with unphysiological levels of SIT. Our study also found SIT to significantly increase NO production within 24 h of incubation with LPS. This increase in NO might be part of the initial immune response to combat the inflammation. More studies are needed to understand how physiological levels of SIT might affect the NO production after 24 h of incubation.

VD3 inhibits cell proliferation and promotes differentiation [29,63]. The anti-proliferative property of VD3 has been found to be beneficial in combating cancer [64–70]. VD3 stimulation of dendritic cells has been shown to induce the apoptosis in T-cells [71]. VD3 has been shown to decrease CD4+ cell proliferation in experimental autoimmune encephalitis model [72] and PBMC multiplication in inflammatory bowel disease [73]. Bartels et al also reported similar results whereby VD3 stimulation increased apoptosis and decreased cell proliferation of T-cells in Crohn's disease [39]. Our results showed a trend for VD3 to reduce cell proliferation although it did not reach significance. This could have been due to the small number of observations used in our experiment.

SIT is an anti-proliferative and apoptotic molecule to several tumor cells [74,75]. Previous studies from our lab have shown SIT to activate the sphingomyelin cycle [76,77], impede the cell cycle [78–80] and stimulate apoptotic cell death in tumor cells [81]. Others have demonstrated that SIT induces apoptosis in U937 cells by caspase-3 activation and Bax/Bcl-2 ratio up-regulation [74]. Our present results are also in accordance with previously published data and further substantiate the anti-proliferative properties of SIT in J774A.1 macrophages.

Combining SIT and VD3, an additive effect has been revealed compared to what has been noticed with their individual treatments. This suggests that the effect of these agents may be similar and in the same direction. This could also imply that both agents may influence the same signal transduction pathways. The combined treatment showed that cell proliferation decreased significantly from vehicle control. The decrease in cell proliferation, albeit insignificant, was even lower than the individual SIT treatment. The VD3, as already discussed, did not bring about much difference in the cytokine or NO release and cell proliferation. SIT, on the other hand, significantly increased the cytokine and NO response. The release of cytokines and NO from the macrophages subjected to the combined treatment was even higher. The SIT + VD3 amplified the cytokine and NO release and the effect was found to be additive and in some cases synergistic. Such

an amplified Th1 and Th2 response could be beneficial in counter-acting pathogens and controlling autoimmune diseases.

To understand the underlying mechanism behind this treatment response we decided to look at NF- κ B phosphorylation. NF- κ B-p65 phosphorylation at Ser 468 is primarily induced by phorbol esters while Ser 536 phosphorylation is mediated by LPS. Although we could not apply statistical significance to our results due to limited number of observations, we found that all our treatments decreased Ser536 phosphorylation. This is of importance because increased Ser536 phosphorylation is thought to inhibit the p53, mediator of apoptosis [82]. This might explain the decreased cell proliferation we observed with our treatments.

Surprisingly, increased Ser 468 phosphorylation was found to be associated with the SIT and SIT + VD3 treatments. This corresponds well with the increase in cytokine and NO release we noticed in our results. Since we observed a decrease in Ser536, this may suggest that p65 has not been activated to a greater extent. So there might be one or more alternative pathways being activated to observe such response we noticed. Since Ser 468 phosphorylation is not expected with LPS stimulation, one may consider the cross talk between the signaling pathways inside the cell a factor in this deviation. The autocrine and paracrine nature of released cytokines would make identifying the role for signal transduction pathways involved a major task.

Other mechanisms may include the possibility that SIT works as a carrier for VD3 by virtue of the similarity in structure between these molecules associated with the difference in their concentrations in the blood. SIT may also influence the binding properties of VD3 to its receptors. However, these proposed mechanisms have to wait further investigation.

In summary, although much more work is necessary to understand the underlying mechanism, we conclude that SIT promotes the immune function of VD3. This ability of SIT to enhance the immunomodulatory action of vitamin D could be beneficial to vitamin D deficient individuals. It may not be easy to overcome a vitamin D deficiency as for people living in temperate climate, with disabilities, old age or renal diseases. However, to include fresh vegetables in the diet and thus enhancing the action of vitamin D may be an option. This study has far reaching implications not only for people with vitamin D deficiency but also for people with autoimmune diseases like multiple sclerosis.

References

- [1] Cranney C HT, O'Donnell S, Weiler H, Ooi D, Atkinson S, et al. Effectiveness and safety of vitamin D. Evidence Report/Technology Assessment No. 158 prepared by the University of Ottawa Evidence-based Practice Center under Contract No. 290-02.0021. AHRQ Publication No. 07-E013. Rockville, MD: Agency for Healthcare Research and Quality. 2007.
- [2] Institute of Medicine FaNBW. DC Dietary Reference Intakes: Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. National Academy Press; 1997.
- [3] van den Berg H. Bioavailability of vitamin D. Eur J Clin Nutr 1997;51:S76–79.
- [4] DeLuca HF. Overview of general physiologic features and functions of vitamin D. Am J Clin Nutr 2004;80:1689S–96S.
- [5] Pfeifer M, Begerow B, Minne HW. Vitamin D and muscle function. Osteoporos Int 2002;13:187–94.
- [6] Hayes CE HF, Spach KM, Pederson LB. The immunological functions of the vitamin D endocrine system. Cell Mol Biol 2003;49:277–300.
- [7] Holick MF. Evolution and function of vitamin D. Recent results. Cancer Res 2003;164:3–28.
- [8] Office of Dietary Supplements NIOH, Bethesda, Maryland. Dietary Supplement Fact Sheet: Vitamin D 2009.
- [9] Chesney RW. Vitamin D: can an upper limit be defined? J Nutr 1989;119:1825–8.
- [10] Favus MJ CS. Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism. Lippincott-Raven3rd ed. ; 1996. Philadelphia.
- [11] Jackson RD LA, Gass M, Wallace RB, Robbins J, Lewis CE, et al. Calcium plus vitamin D supplementation and the risk of fractures. N Engl J Med 2006;354:669–83.
- [12] Biser-Rohrbaugh AH-MN. Vitamin D deficiency in breast-fed toddlers. J Pediatr Orthop 2001;21:508–11.
- [13] Cohen-Lahav M, Douvdevani A, Chaimovitz C, Shany S. The anti-inflammatory activity of 1, 25-dihydroxyvitamin D3 in macrophages. J Steroid Biochem Mol Biol 2007;103:558–62.
- [14] Desai F, Ramanathan M, Fink CS, Wilding GE, Weinstock-Guttman B, Awad AB. Comparison of the immunomodulatory effects of the plant sterol beta-sitosterol

- to simvastatin in peripheral blood cells from multiple sclerosis patients. *Int Immunopharmacol* 2009;9:153–7.
- [15] Awad AB, Chan KC, Downie AC, Fink CS. Peanuts as a source of beta-sitosterol, a sterol with anticancer properties. *Nutr Cancer* 2000;36:238–41.
 - [16] Kennedy AR. The evidence for soybean products as cancer preventive agents. *J Nutr* 1995;125:733S–43S.
 - [17] Katan MB, Grundy SM, Jones P, Law M, Miettinen T, Paoletti R. Efficacy and safety of plant stanols and sterols in the management of blood cholesterol levels. *Mayo Clin Proc* 2003;78:965–78.
 - [18] Pham TN, Brown BL, Dobson PR, Richardson VJ. Protein kinase C- η (PKC- η) is required for the development of inducible nitric oxide synthase (iNOS) positive phenotype in human monocytic cells. *Nitric Oxide* 2003;9:123–34.
 - [19] Smith KL. The science of the total environment. 2000;262:185–190.
 - [20] Cohen-Lahav M, Shany S, Tobvin D, Chaimovitz C, Douvdevani A. Vitamin D decreases NF- κ B activity by increasing IkappaBalpha levels. *Nephrol Dial Transplant* 2006;21:889–97.
 - [21] C Jer-Ming, Kuo Mei-Chuan, Kuo Hung-Tien, Hwang Shang-Jyh, Tsai Jer-Chia, Chen Hung-Chun, et al. 1- α ,25-Dihydroxyvitamin D₃ regulates inducible nitric oxide synthase messenger RNA expression and nitric oxide release in macrophage-like RAW 264.7 cells. *J Lab Clin Med* 2003;143.
 - [22] Ardizzone S, Cassinotti A, Trabattini D, Manzionna G, Rainone V, Bevilacqua M, et al. Immunomodulatory effects of 1, 25-dihydroxyvitamin D₃ on TH1/TH2 cytokines in inflammatory bowel disease: an in vitro study. *Int J Immunopathol Pharmacol* 2009;22:63–71.
 - [23] May E, Asadullah K, Zugel U. Immunoregulation through 1, 25-dihydroxyvitamin D₃ and its analogs. *Curr Drug Targets Inflamm Allergy* 2004;3:377–93.
 - [24] Shany S, Rapoport J, Zuili I, Gavriel A, Lavi N, Chaimovitz C. Metabolism of 25-OH-vitamin D₃ by peritoneal macrophages from CAPD patients. *Kidney Int* 1991;39:1005–11.
 - [25] Schleithoff SS, Zittermann A, Tenderich G, Berthold HK, Stehle P, Koerfer R. Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind, randomized, placebo-controlled trial. *Am J Clin Nutr* 2006;83:754–9.
 - [26] Booi-Vrieling HE, Tryfonidou MA, Riemers FM, Penning LC, Hazewinkel HA. Inflammatory cytokines and the nuclear vitamin D receptor are implicated in the pathophysiology of dental resorptive lesions in cats. *Vet Immunol Immunopathol* 2009.
 - [27] Lefebvre d'Hellencourt C, Montero-Menei CN, Bernard R, Couez D. Vitamin D₃ inhibits proinflammatory cytokines and nitric oxide production by the EOC13 microglial cell line. *J Neurosci Res* 2003;71:575–82.
 - [28] Giarratana N, Penna G, Amuchastegui S, Mariani R, Daniel KC, Adorini L. A vitamin D analog down-regulates proinflammatory chemokine production by pancreatic islets inhibiting T cell recruitment and type 1 diabetes development. *J Immunol* 2004;173:2280–7.
 - [29] Hansdottir S, Monick MM, Hinde SL, Lohan N, Look DC, Hunninghake GW. Respiratory epithelial cells convert inactive vitamin D to its active form: potential effects on host defense. *J Immunol* 2008;181:7090–9.
 - [30] Zhang Z, Yuan W, Sun L, Szeto FL, Wong KE, Li X, et al. 1, 25-Dihydroxyvitamin D₃ targeting of NF- κ B suppresses high glucose-induced MCP-1 expression in mesangial cells. *Kidney Int* 2007;72:193–201.
 - [31] Yu XP, Bellido T, Manolagas SC. Down-regulation of NF- κ B protein levels in activated human lymphocytes by 1, 25-dihydroxyvitamin D₃. *Proc Natl Acad Sci USA* 1995;92:10990–4.
 - [32] Riis JL, Johansen C, Gesser B, Moller K, Larsen CG, Kragballe K, et al. 1 α , 25(OH) (2)D₃ regulates NF- κ B DNA binding activity in cultured normal human keratinocytes through an increase in IkappaBalpha expression. *Arch Dermatol Res* 2004;296:195–202.
 - [33] Sun J, Kong J, Duan Y, Szeto FL, Liao A, Madara JL, et al. Increased NF- κ B activity in fibroblasts lacking the vitamin D receptor. *Am J Physiol Endocrinol Metab* 2006;291:E315–22.
 - [34] Suzuki Y, Ichijima T, Ohsaki A, Hasegawa S, Shiraishi M, Furukawa S. Anti-inflammatory effect of 1 α , 25-dihydroxyvitamin D₃ in human coronary arterial endothelial cells: Implication for the treatment of Kawasaki disease. *J Steroid Biochem Mol Biol* 2009;113:134–8.
 - [35] Peterson CA, Heffernan ME. Serum tumor necrosis factor- α concentrations are negatively correlated with serum 25(OH)D concentrations in healthy women. *J Inflamm (Lond)* 2008;5:10.
 - [36] Shany S, Levy Y, Lahav-Cohen M. The effects of 1 α , 24(S)-dihydroxyvitamin D (2) analog on cancer cell proliferation and cytokine expression. *Steroids* 2001;66:319–25.
 - [37] Sun X, Zemmel MB. Calcitriol and calcium regulate cytokine production and adipocyte-macrophage cross-talk. *J Nutr Biochem* 2008;19:392–9.
 - [38] Hakim I, Bar-Shavit Z. Modulation of TNF- α expression in bone marrow macrophages: involvement of vitamin D response element. *J Cell Biochem* 2003;88:986–98.
 - [39] Bartels LE, Jorgensen SP, Agnholt J, Kelsen J, Hvas CL, Dahlerup JF. 1, 25-dihydroxyvitamin D₃ and dexamethasone increase interleukin-10 production in CD4+ T cells from patients with Crohn's disease. *Int Immunopharmacol* 2007;7:1755–64.
 - [40] Harant H, Andrew PJ, Reddy GS, Foglar E, Lindley JJ. 1 α , 25-dihydroxyvitamin D₃ and a variety of its natural metabolites transcriptionally repress nuclear-factor- κ B-mediated interleukin-8 gene expression. *Eur J Biochem* 1997;250:63–71.
 - [41] D'Ambrosio D, Cipitelli M, Cocciolo MG, Mazzeo D, Di Lucia P, Lang R, et al. Ambition of IL-12 production by 1, 25-dihydroxyvitamin D₃. Involvement of NF- κ B downregulation in transcriptional repression of the p40 gene. *J Clin Invest* 1998;101:252–62.
 - [42] Alroy I, Towers TL, Freedman LP. Transcriptional repression of the interleukin-2 gene by vitamin D₃: direct inhibition of NFATp/AP-1 complex formation by a nuclear hormone receptor. *Mol Cell Biol* 1995;15:5789–99.
 - [43] Hayes CE, Donald Acheson E. A unifying multiple sclerosis etiology linking virus infection, sunlight, and vitamin D, through viral interleukin-10. *Med Hypotheses* 2008;71:85–90.
 - [44] Fang FC. Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related antimicrobial activity. *J Clin Invest* 1997;99:2818–25.
 - [45] Wheeler MA, Smith SD, Garcia-Cardena G, Nathan CF, Weiss RM, Sessa WC. Bacterial infection induces nitric oxide synthase in human neutrophils. *J Clin Invest* 1997;99:110–6.
 - [46] Eun JS, Suh YH, Kim DK, Jeon H. Regulation of cytokine production by exogenous nitric oxide in murine splenocyte and peritoneal macrophage. *Arch Pharm Res* 2000;23:531–4.
 - [47] MacMicking J, Xie QW, Nathan C. Nitric oxide and macrophage function. *Annu Rev Immunol* 1997;15:323–50.
 - [48] Zhou Y, Zhou X, Wang X. 1, 25-Dihydroxyvitamin D₃ prevented allergic asthma in a rat model by suppressing the expression of inducible nitric oxide synthase. *Allergy Asthma Proc* 2008;29:258–67.
 - [49] Jiao ZM, Fu YH, Fu J, Zhang F, Wang WZ. 1, 25-dihydroxyvitamin D₃ promotes the apoptosis of inflammatory cells in acute experimental autoimmune encephalomyelitis: experiment with rats. *Zhonghua Yi Xue Za Zhi* 2008;88:2350–4.
 - [50] Chan J, Xing Y, Magliozzo RS, Bloom BR. Killing of virulent Mycobacterium tuberculosis by reactive nitrogen intermediates produced by activated murine macrophages. *J Exp Med* 1992;175:1111–22.
 - [51] Rockett KA, Brookes R, Udaloa I, Vidal V, Hill AV, Kwiatkowski D. 1, 25-Dihydroxyvitamin D₃ induces nitric oxide synthase and suppresses growth of Mycobacterium tuberculosis in a human macrophage-like cell line. *Infect Immun* 1998;66:5314–21.
 - [52] Bouic PJ, Lamprecht JH. Plant sterols and sterolins: a review of their immunomodulating properties. *Altern Med Rev* 1999;4:170–7.
 - [53] Bouic PJ, Clark A, Lamprecht J, Freestone M, Pool EJ, Liebenberg RW, et al. The effects of B-sitosterol (BSS) and B-sitosterol glucoside (BSSG) mixture on selected immune parameters of marathon runners: inhibition of post marathon immune suppression and inflammation. *Int J Sports Med* 1999;20:258–62.
 - [54] Bouic PJ, Clark A, Brittle W, Lamprecht JH, Freestone M, Liebenberg RW. Plant sterol/sterolin supplement use in a cohort of South African HIV-infected patients—effects on immunological and virological surrogate markers. *S Afr Med J* 2001;91:848–50.
 - [55] Bouic PJ. The role of phytosterols and phytosterolins in immune modulation: a review of the past 10 years. *Curr Opin Clin Nutr Metab Care* 2001;4:471–5.
 - [56] Garcia MD, Saenz MT, Gomez MA, Fernandez MA. Topical antiinflammatory activity of phytosterols isolated from Eryngium foetidum on chronic and acute inflammation models. *Phytother Res* 1999;13:78–80.
 - [57] Ding Y, Nguyen HT, Kim SI, Kim HW, Kim YH. The regulation of inflammatory cytokine secretion in macrophage cell line by the chemical constituents of Rhus sylvestris. *Bioorg Med Chem Lett* 2009;19:3607–10.
 - [58] Fang SH, Rao YK, Tzeng YM. Anti-oxidant and inflammatory mediator's growth inhibitory effects of compounds isolated from Phyllanthus urinaria. *J Ethnopharmacol* 2008;116:333–40.
 - [59] Aherne SA, O'Brien NM. Modulation of cytokine production by plant sterols in stimulated human Jurkat T cells. *Mol Nutr Food Res* 2008;52:664–73.
 - [60] Lee JH, Lee JY, Park JH, Jung HS, Kim JS, Kang SS, et al. Immunoregulatory activity by daucosterol, a beta-sitosterol glycoside, induces protective Th1 immune response against disseminated Candidiasis in mice. *Vaccine* 2007;25:3834–40.
 - [61] Myers L, Bouic PJD. Flow cytometric analysis of the TH1-TH2 shift in allergic individuals using Moducare™ (sterols/sterolins). 26th Annual Congress of the Physiology Society of Southern Africa; 1998.
 - [62] Moreno JJ. Effect of olive oil minor components on oxidative stress and arachidonic acid mobilization and metabolism by macrophages RAW 264.7. *Free Radic Biol Med* 2003;35:1073–81.
 - [63] Verstuyf A, Segaeert S, Verlinden L, Casteels K, Bouillon R, Mathieu C. Recent developments in the use of vitamin D analogues. *Curr Opin Nephrol Hypertens* 1998;7:397–403.
 - [64] Wactawski-Wende J, Kotchen JM, Anderson GL, Assaf AR, Brunner RL, O'Sullivan MJ, et al. Calcium plus vitamin D supplementation and the risk of colorectal cancer. *N Engl J Med* 2006;354:684–96.
 - [65] Garland CF, Garland FC, Gorham ED, Lipkin M, Newmark H, Mohr SB, et al. The role of vitamin D in cancer prevention. *Am J Public Health* 2006;96:252–61.
 - [66] Garland CF, Garland FC. Do sunlight and vitamin D reduce the likelihood of colon cancer? *Int J Epidemiol* 2006;35:217–20.
 - [67] Giovannucci E, Liu Y, Rimm EB, Hollis BW, Fuchs CS, Stampfer MJ, et al. Prospective study of predictors of vitamin D status and cancer incidence and mortality in men. *J Natl Cancer Inst* 2006;98:451–9.
 - [68] Ahonen MH, Tenkanen L, Teppo L, Hakama M, Tuohimaa P. Prostate cancer risk and prediagnostic serum 25-hydroxyvitamin D levels (Finland). *Cancer Causes Control* 2000;11:847–52.
 - [69] Wu K, Feskani D, Fuchs CS, Willett WC, Hollis BW, Giovannucci EL. A nested case control study of plasma 25-hydroxyvitamin D concentrations and risk of colorectal cancer. *J Natl Cancer Inst* 2007;99:1120–9.
 - [70] Young MV, Schwartz GG, Wang L, Jamieson DP, Whitlatch LW, Flanagan JN, et al. The prostate 25-hydroxyvitamin D-1 α -hydroxylase is not influenced by parathyroid hormone and calcium: implications for prostate cancer chemoprevention by vitamin D. *Carcinogenesis* 2004;25:967–71.
 - [71] van Halteren AG, Tysma OM, van Etten E, Mathieu C, Roep BO. 1 α , 25-dihydroxyvitamin D₃ or analogue treated dendritic cells modulate human

- autoreactive T cells via the selective induction of apoptosis. *J Autoimmun* 2004;23: 233–9.
- [72] Pedersen LB, Nashold FE, Spach KM, Hayes CE. 1, 25-dihydroxyvitamin D3 reverses experimental autoimmune encephalomyelitis by inhibiting chemokine synthesis and monocyte trafficking. *J Neurosci Res* 2007;85:2480–90.
- [73] Stio M, Martinesi M, Bruni S, Treves C, d'Albasio G, Bagnoli S, et al. Interaction among vitamin D(3) analogue KH 1060, TNF-alpha, and vitamin D receptor protein in peripheral blood mononuclear cells of inflammatory bowel disease patients. *Int Immunopharmacol* 2006;6:1083–92.
- [74] Park C, Moon DO, Rhu CH, Choi BT, Lee WH, Kim GY, et al. Beta-sitosterol induces anti-proliferation and apoptosis in human leukemic U937 cells through activation of caspase-3 and induction of Bax/Bcl-2 ratio. *Biol Pharm Bull* 2007;30:1317–23.
- [75] Awad AB, Fink CS. Phytosterols as anticancer dietary components: evidence and mechanism of action. *J Nutr* 2000;130:2127–30.
- [76] von Holtz RL, Fink CS, Awad AB. beta-Sitosterol activates the sphingomyelin cycle and induces apoptosis in LNCaP human prostate cancer cells. *Nutr Cancer* 1998;32: 8–12.
- [77] Awad AB, von Holtz RL, Cone JP, Fink CS, Chen YC. beta-Sitosterol inhibits growth of HT-29 human colon cancer cells by activating the sphingomyelin cycle. *Anticancer Res* 1998;18:471–3.
- [78] Awad AB, Williams H, Fink CS. Phytosterols reduce in vitro metastatic ability of MDA-MB-231 human breast cancer cells. *Nutr Cancer* 2001;40:157–64.
- [79] Awad AB, Fink CS, Williams H, Kim U. In vitro and in vivo (SCID mice) effects of phytosterols on the growth and dissemination of human prostate cancer PC-3 cells. *Eur J Cancer Prev* 2001;10:507–13.
- [80] Awad AB, Smith AJ, Fink CS. Plant sterols regulate rat vascular smooth muscle cell growth and prostacyclin release in culture. *Prostaglandins Leukot Essent Fatty Acids* 2001;64:323–30.
- [81] Awad AB, Roy R, Fink CS. Beta-sitosterol, a plant sterol, induces apoptosis and activates key caspases in MDA-MB-231 human breast cancer cells. *Oncol Rep* 2003;10:497–500.
- [82] Jeong SJ, Pise-Masison CA, Radonovich MF, Park HU, Brady JN. A novel NF-kappaB pathway involving IKKbeta and p65/RelA Ser-536 phosphorylation results in p53 inhibition in the absence of NF-kappaB transcriptional activity. *J Biol Chem* 2005;280:10326–32.