

Endothelial Function and Cerebrovascular Disease: Implications for Diagnosis and Treatment

Ahmad A. Elesber, MD

Piero O. Bonetti, MD

*Amir Lerman, MD**

Address

*Division of Cardiovascular Diseases, Mayo Clinic, 200 First Street SW, Rochester, MN 55905, USA.

E-mail: Lerman.amir@mayo.edu

Current Treatment Options in Cardiovascular Medicine 2006, **8**:213–219

Current Science Inc. ISSN 1092–8464

Copyright © 2006 by Current Science Inc.

Opinion statement

Cerebrovascular disease remains one of the most common causes of morbidity and mortality in the United States. There is strong evidence to implicate endothelial dysfunction in the initiation and progression of atherosclerosis and its complications. It is now well known that endothelial dysfunction represents a systemic syndrome involving multiple vascular beds, including the cerebral vasculature. Currently, no gold standard treatment for endothelial dysfunction exists. Nonetheless, several treatment strategies have been found to be helpful in improving endothelial function. A few of these strategies have been implicated in stroke risk reduction as well, adding another line of evidence to the relationship between endothelial function and cerebrovascular disease.

Introduction

It is now well recognized that the endothelium is an active biologic organ strategically located between the vascular wall and the circulation and is indispensable for a healthy vascular homeostasis (Table 1). Endothelial dysfunction is functionally characterized by an impairment of endothelium-dependent vasodilation, which is mainly caused by a reduced bioavailability of endothelium-derived nitric oxide (NO). In addition, endothelial dysfunction is associated with an imbalance in favor of increased inflammation, oxidative stress, thrombosis, and proliferation of the vasculature contributing to the initiation and development of atherosclerosis [1].

Endothelial dysfunction represents a systemic disorder involving multiple vascular beds [2]. Accordingly, measurement of endothelial function in a particular vascular bed (eg, the coronary arteries or the brachial artery) mirrors the functional state of the entire endothelium, including that lining the cerebral vasculature. Endothelial function testing relies on assessing endothelium-depen-

dent vasodilation. Such tests range from the invasive coronary angiographic measurements coupled with Doppler flow assessment of coronary blood flow in response to intracoronary injection of endothelium-dependent vasodilators (eg, acetylcholine), to less invasive methods employing intra-arterial infusion of endothelium-dependent vasodilators coupled with measurements of forearm arterial response, and finally to noninvasive measurement of endothelium-dependent flow-mediated dilatation of the brachial or digital arteries [3]. Limitations of the different tests used have recently been nicely reviewed with a call for the development of a noninvasive and more reliable method for assessment of endothelial dysfunction and vascular health [4].

Various studies showed that the presence of endothelial dysfunction in peripheral arteries or the coronary circulation is independently associated with different cardiovascular combined end points, including cerebrovascular events [5–7]. Notably, the study by Targonski et al. [8•] showed

Table 1. Function of a normal endothelium

Control of vasomotor tone
Regulation of nutrient trafficking
Control of leukocyte adhesion and migration
Regulation of smooth muscle cell migration and multiplication
Regulation of platelet adhesion and aggregation
Balance between coagulation and anticoagulation
Balance between antifibrinolytic and profibrinolytic effects
Control of inflammation
Antioxidant effects
Neovascularization

that the presence of coronary endothelial dysfunction in patients without obstructive coronary atherosclerosis is associated with a fourfold increased risk of cerebrovascular events independent of known risk factors for stroke. These results suggest that a healthy endothelium represents a key protective factor against cerebrovascular events. Endothelium-derived NO was shown to be a critical factor for cerebrovascular health [9]. The importance of endothelial dysfunction in the pathogenesis of stroke is further highlighted by the fact that different treatment strategies that decrease the risk of cerebrovascular events also improve endothelial dysfunction [10,11]. The present review summarizes various treatment options for endothelial dysfunction and points out currently available evidence for their role in the prevention of cerebrovascular events.

Treatment

- There is no gold standard therapy for endothelial dysfunction. Moreover, there is currently no study that demonstrates an association between the improvement of endothelial function and reduction of stroke with any intervention. However, given that endothelial dysfunction is associated with all currently known cardiovascular risk factors, it is not surprising that treatment or elimination of these cardiovascular risk factors (eg, blood pressure-lowering therapy, smoking cessation) in general also translates into improvement of endothelial function. In addition to traditional cardiovascular risk factor modification, some specific dietary supplements, lifestyle modification strategies, and nonpharmacologic and pharmacologic therapies were shown to lead to an improvement in endothelial function. Notably, some of these treatment modalities have been shown to reduce the risk of cerebrovascular events as well. Nonetheless, it still remains to be determined whether an improvement of endothelial function also translates directly into clinical improvement in a patient's prognosis [3].
- Taken together, our current understanding is that it may be reasonable to conclude that modifications of lifestyle with the additional of statin and aspirin may be recommended as measures to improve endothelial function and stroke prevention.

Dietary supplements

Folic acid

- A small randomized crossover study of patients with coronary artery disease showed that folic acid supplementation resulted in improvement of peripheral endothelial dysfunction [12]. Endothelial function was also improved in patients with familial hypercholesterolemia [13]. Moreover, in a large prospective observational study of 43,732 healthy men, there was an inverse relationship between dietary folate intake and the risk of ischemic stroke [14]. Nonetheless, the results of the randomized controlled VISP (Vitamin Intervention for Stroke Prevention) trial that compared the effect of high doses of folic acid, pyridoxine (vitamin B₆), and cobalamin (vitamin B₁₂) to the effect of low doses of these vitamins failed to show a reduction in the risk of recurrent stroke over a 2-year period [15]. Moreover, the recently presented results

of the NORVIT (Norwegian Vitamin Trial) suggest that combined supplementation of high-dose pyridoxine and folic acid may even increase the risk of stroke in survivors of myocardial infarction [16].

Antioxidants

- With a few exceptions, acute administration of vitamin C was shown to reverse endothelial dysfunction of patients with cardiovascular risk factors and/or overt atherosclerosis [17]. In contrast, the effect of vitamin E on endothelial function remains controversial [18]. Nonetheless, human studies failed to show any benefit on stroke prevention with use of vitamin E or vitamin C [19].
- Other antioxidants that have been shown to improve endothelial function include glutathione [20] and *N*-acetylcysteine [21]. However, the benefits of glutathione [22] and *N*-acetylcysteine [23] on stroke have been limited to experimental models.

L-Arginine

- Conflicting data exist on the utility of L-arginine supplementation. In one study, long-term oral L-arginine supplementation for 6 months improved patients' symptoms of angina, coronary endothelial function, and plasma endothelin concentrations [24]. In another randomized, double-blind, placebo-controlled crossover trial of patients with established coronary artery disease and angina, flow-mediated brachial artery dilation, treadmill exercise time, and quality-of-life scores improved with a specially formulated nutrition bar rich in L-arginine [25]. However, a different study showed no benefit of L-arginine supplementation on endothelial function [26].
- No clinical trial has yet evaluated the effect of L-arginine supplementation on the risk of stroke. A recent systematic review of available data on experimental stroke showed that L-arginine was ineffective at reducing lesion volume in transient stroke, permanent stroke, or both [27]. In one study, L-arginine even significantly increased infarct volume [28].

Lifestyle modification

The type of diet

- The Mediterranean diet has been shown to have beneficial effect on peripheral endothelial function in diabetic [29] and hypercholesterolemic patients [30].
- A prospective cohort study looking at the effect of different dietary patterns found that a dietary style characterized by higher intakes of red and processed meats, refined grains, and sweets increased stroke rate, whereas a diet higher in fruits and vegetables, fish, and whole grains protected against stroke [31].

Weight loss

- Weight loss has been shown to improve endothelial dysfunction in obese women [32] and in children [33].
- To this day, no study has looked specifically at the relationship between weight loss and stroke risk reduction.

Physical exercise

- Exercise training was shown to prevent age-associated endothelial dysfunction and to improve endothelial dysfunction in previously sedentary middle-aged and older healthy men [34]. In patients with coronary artery disease, regular aerobic exercise was shown to improve coronary endothelial function [35].
- A meta-analysis of 23 studies looking at the overall association between physical activity and stroke incidence and mortality found that moderate or high levels of exercise are associated with a reduced risk of stroke [36].

Smoking cessation

- Cigarette smoking was demonstrated to be associated in a dose-related fashion with peripheral endothelial dysfunction. Smoking cessation was found to result in improvement of endothelial function in asymptomatic young adults [37].
- The risk of stroke in patients who stopped smoking was found to approach that of nonsmokers over the course of a few years [38].

Pharmacologic therapy

3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors (statins)

- There is strong evidence from both experimental and clinical studies that treatment with statins enhances endothelial NO bioavailability and improves endothelial function, an effect that seems to be at least partly mediated by lipid-independent effects of this class of drugs [39]. Notably, experimental studies in a normocholesterolemic murine model demonstrated that prophylactic treatment with statins increased cerebral blood flow, reduced cerebral infarct size, and improved neurologic function via an NO-mediated mechanism [40]. Numerous large-scale randomized clinical trials have shown that statins reduce the risk of stroke in patients with established coronary artery disease or in individuals who are considered at risk for cardiovascular events [41]. Based on the encouraging results of these trials, the Stroke Council of the American Heart Association/American Stroke Association issued an advisory statement indicating that the majority of patients with a history of ischemic stroke or transient ischemic attack of atherosclerotic origin could benefit from statin therapy. Moreover, initiation of statin therapy during hospitalization for first ischemic stroke is probably justified [42]. However, results of the ongoing SPARCL (Stroke Prevention by Aggressive Reduction in Cholesterol Levels) trial will have to provide definite information about the role of statins in patients with stroke but no history of other vascular disease. Nonetheless, given that hypercholesterolemia is not an established risk factor for stroke and given the favorable lipid-independent effects of statins on endothelial function, statin-induced improvement of endothelial function represents a likely mechanism contributing to the beneficial effect of statins on the risk of cerebrovascular events observed in large clinical trials.

Angiotensin-converting enzyme inhibitors

- Angiotensin-converting enzyme inhibitors have been shown to improve endothelial function in both patients with coronary artery disease and in patients at risk for atherosclerosis [43,44]. The proposed mechanism for improvement of endothelial function includes enhancing NO bioavailability by decreasing angiotensin II and increasing bradykinin levels.
- There is a paucity of trials looking specifically at the effect of angiotensin-converting enzyme inhibitors on prevention of stroke independent of the effect on blood pressure lowering [45]. In the HOPE (Heart Outcomes Prevention Evaluation) trial, by analyzing the end point of stroke separately from the primary composite end point of cardiovascular death, myocardial infarction, and stroke, ramipril was shown to have a significant protective effect on stroke risk [46]. The benefit of ramipril was thought to go beyond the small attained improvement in blood pressure [47].

Angiotensin II receptor blockers

- Several studies showed the benefit of angiotensin II receptor blockers on endothelial function in both patients with coronary artery disease and in patients at risk for atherosclerosis [48,49].
- In the LIFE (Losartan Intervention For Endpoint reduction for hypertension study) trial, losartan treatment resulted in fewer cardiovascular deaths, myocardial infarctions, and stroke when compared with atenolol. This effect was due to reduction in stroke risk driving the primary composite end point in favor of losartan. The conclusion was that the benefit of blocking the angiotensin II receptor goes beyond the blood pressure-lowering effect of losartan [50].

Estrogen replacement therapy

- The effect of estrogen replacement therapy on endothelial function in postmenopausal women is controversial [51,52]. Unfortunately, estrogen replacement therapy was found to be associated with an increased stroke risk in several studies [53••,54••]. Discrepancies between these randomized studies and earlier observational studies are difficult to reconcile but different hypotheses have been advanced [55].

Aspirin

- The role of aspirin in the reduction of the risk of cerebrovascular events is well established [56]. Preliminary studies demonstrated that acute administration of aspirin may improve peripheral endothelial function in humans [11].

References and Recommended Reading

Papers of particular interest, published recently, have been highlighted as:

- Of importance
 - Of major importance
1. Lerman A, Zeiher AM: **Endothelial function: cardiac events.** *Circulation* 2005, **111**:363–368.
 2. Anderson TJ, Uehata A, Gerhard MD, et al.: **Close relation of endothelial function in the human coronary and peripheral circulations.** *J Am Coll Cardiol* 1995, **26**:1235–1241.
 3. Bonetti PO, Lerman LO, Lerman A: **Endothelial dysfunction: a marker of atherosclerotic risk.** *Arterioscler Thromb Vasc Biol* 2003, **23**:168–175.
 4. Ganz P, Vita JA: **Testing endothelial vasomotor function: nitric oxide, a multipotent molecule.** *Circulation* 2003, **108**:2049–2053.

5. Perticone F, Ceravolo R, Pujia A, et al.: **Prognostic significance of endothelial dysfunction in hypertensive patients.** *Circulation* 2001, 104:191–196.
 6. Halcox JP, Schenke WH, Zalos G, et al.: **Prognostic value of coronary vascular endothelial dysfunction.** *Circulation* 2002, 106:653–658.
 7. Schachinger V, Britten MB, Zeiher AM: **Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease.** *Circulation* 2000, 101:1899–1906.
 8. Targonski PV, Bonetti PO, Pumper GM, et al.: **Coronary endothelial dysfunction is associated with an increased risk of cerebrovascular events.** *Circulation* 2003, 107:2805–2809.
- First study to show that coronary endothelial dysfunction is associated with increased cerebrovascular events.
9. McCarty MF: **Up-regulation of endothelial nitric oxide activity as a central strategy for prevention of ischemic stroke—just say NO to stroke!** *Med Hypotheses* 2000, 55:386–403.
 10. O'Driscoll G, Green D, Taylor RR: **Simvastatin, an HMG-coenzyme A reductase inhibitor, improves endothelial function within 1 month.** *Circulation* 1997, 95:1126–1131.
 11. Husain S, Andrews NP, Mulcahy D, et al.: **Aspirin improves endothelial dysfunction in atherosclerosis.** *Circulation* 1998, 97:716–720.
 12. Doshi SN, McDowell IF, Moat SJ, et al.: **Folate improves endothelial function in coronary artery disease: an effect mediated by reduction of intracellular superoxide?** *Arterioscler Thromb Vasc Biol* 2001, 21:1196–1202.
 13. Verhaar MC, Wever RM, Kastelein JJ, et al.: **Effects of oral folic acid supplementation on endothelial function in familial hypercholesterolemia. A randomized placebo-controlled trial.** *Circulation* 1999, 100:335–338.
 14. He K, et al.: **Folate, vitamin B6, and B12 intakes in relation to risk of stroke among men.** *Stroke* 2004, 35:169–174.
 15. Toole JF, Malinow MR, Chambless LE, et al.: **Lowering homocysteine in patients with ischemic stroke to prevent recurrent stroke, myocardial infarction, and death: the Vitamin Intervention for Stroke Prevention (VISP) randomized controlled trial.** *JAMA* 2004, 291:565–575.
 16. Bona KH, NORVIT: **Randomized trial of homocysteine-lowering with B-vitamins for secondary prevention of cardiovascular disease after acute myocardial infarction.** *Program and Abstracts from the European Society of Cardiology Congress 2005.* Stockholm, Sweden; September 3–7, 2005.
 17. Carr A, Frei B: **The role of natural antioxidants in preserving the biological activity of endothelium-derived nitric oxide.** *Free Radic Biol Med* 2000, 28:1806–1814.
 18. Visioli F: **Effects of vitamin E on the endothelium: equivocal? Alpha-tocopherol and endothelial dysfunction.** *Cardiovasc Res* 2001, 51:198–201.
 19. Lonn E, Bosch J, Yusuf S, et al.: **Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial.** *JAMA* 2005, 293:1338–1347.
 20. Prasad A, Andrews NP, Padder FA, et al.: **Glutathione reverses endothelial dysfunction and improves nitric oxide bioavailability.** *J Am Coll Cardiol* 1999, 34:507–514.
 21. Andrews NP, Prasad A, Quyyumi AA: **N-acetylcysteine improves coronary and peripheral vascular function.** *J Am Coll Cardiol* 2001, 37:117–123.
 22. De Haan JB, Crack PJ, Flentjar N, et al.: **An imbalance in antioxidant defense affects cellular function: the pathophysiological consequences of a reduction in antioxidant defense in the glutathione peroxidase-1 (Gpx1) knockout mouse.** *Redox Rep* 2003, 8:69–79.
 23. Khan M, Sekhon B, Jatana M, et al.: **Administration of N-acetylcysteine after focal cerebral ischemia protects brain and reduces inflammation in a rat model of experimental stroke.** *J Neurosci Res* 2004, 76:519–527.
 24. Lerman A, Burnett JC Jr, Higano ST, et al.: **Long-term L-arginine supplementation improves small-vessel coronary endothelial function in humans.** *Circulation* 1998, 97:2123–2128.
 25. Maxwell AJ, Zapien MP, Pearce GL, et al.: **Randomized trial of a medical food for the dietary management of chronic, stable angina.** *J Am Coll Cardiol* 2002, 39:37–45.
 26. Blum A, Hathaway L, Mincemoyer R, et al.: **Oral L-arginine in patients with coronary artery disease on medical management.** *Circulation* 2000, 101:2160–2164.
 27. Willmot M, Gray L, Gibson C, et al.: **A systematic review of nitric oxide donors and L-arginine in experimental stroke: effects on infarct size and cerebral blood flow.** *Nitric Oxide* 2005, 12:141–149.
 28. Zhang F, Casey RM, Ross ME, Iadecola C: **Aminoguanidine ameliorates and L-arginine worsens brain damage from intraluminal middle cerebral artery occlusion.** *Stroke* 1996, 27:317–323.
 29. Ryan M, McNerney D, Owens D, et al.: **Diabetes and the Mediterranean diet: a beneficial effect of oleic acid on insulin sensitivity, adipocyte glucose transport and endothelium-dependent vasoreactivity.** *QJM* 2000, 93:85–91.
 30. Fuentes F, Lopez-Miranda J, Sanchez E, et al.: **Mediterranean and low-fat diets improve endothelial function in hypercholesterolemic men.** *Ann Intern Med* 2001, 134:1115–1119.
 31. Fung TT, Stampfer MJ, Manson JE, et al.: **Prospective study of major dietary patterns and stroke risk in women.** *Stroke* 2004, 35:2014–2019.
 32. Ziccardi P, Nappo F, Giugliano G, et al.: **Reduction of inflammatory cytokine concentrations and improvement of endothelial functions in obese women after weight loss over one year.** *Circulation* 2002, 105:804–809.
 33. Woo KS, Chook P, Yu CW, et al.: **Effects of diet and exercise on obesity-related vascular dysfunction in children.** *Circulation* 2004, 109:1981–1986.
 34. DeSouza CA, Shapiro LE, Clevenger CM, et al.: **Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men.** *Circulation* 2000, 102:1351–1357.
 35. Hambrecht R, Wolf A, Gielen S, et al.: **Effect of exercise on coronary endothelial function in patients with coronary artery disease.** *N Engl J Med* 2000, 342:454–460.
 36. Lee CD, Folsom AR, Blair SN: **Physical activity and stroke risk: a meta-analysis.** *Stroke* 2003, 34:2475–2481.

37. Celermajer DS, Sorensen KE, Georgakopoulos D, et al.: Cigarette smoking is associated with dose-related and potentially reversible impairment of endothelium-dependent dilation in healthy young adults. *Circulation* 1993, 88(5 Pt 1):2149–2155.
38. Kawachi I, Colditz GA, Stampfer MJ, et al.: Smoking cessation and decreased risk of stroke in women. *JAMA* 1993, 269:232–236.
39. Bonetti PO, Lerman LO, Napoli C, Lerman A: Statin effects beyond lipid lowering—are they clinically relevant? *Eur Heart J* 2003, 24:225–248.
40. Endres M, Laufs U, Huang Z, et al.: Stroke protection by 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase inhibitors mediated by endothelial nitric oxide synthase. *Proc Natl Acad Sci U S A* 1998, 95:8880–8885.
41. Endres M: Statins and stroke. *J Cereb Blood Flow Metab* 2005, 25:1093–1110.
42. Adams H, Adams R, Del Zoppo G, Goldstein LB: Guidelines for the early management of patients with ischemic stroke: 2005 guidelines update a scientific statement from the Stroke Council of the American Heart Association/American Stroke Association. *Stroke* 2005, 36:916–923.
43. Anderson TJ, Elstein E, Haber H, Charbonneau F: Comparative study of ACE-inhibition, angiotensin II antagonism, and calcium channel blockade on flow-mediated vasodilation in patients with coronary disease (BANFF study). *J Am Coll Cardiol* 2000, 35:60–66.
44. Prasad A, Narayanan S, Husain S, et al.: Insertion-deletion polymorphism of the ACE gene modulates reversibility of endothelial dysfunction with ACE inhibition. *Circulation* 2000, 102:35–41.
45. Sokol SI, Portnay EL, Curtis JP, et al.: Modulation of the renin-angiotensin-aldosterone system for the secondary prevention of stroke. *Neurology* 2004, 63:208–213.
46. Yusuf S, Sleight P, Pogue J, et al.: Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000, 342:145–153. [Published erratum appears in *N Engl J Med* 2000, 42:1376.]
47. Sleight P, Yusuf S, Pogue J, et al.: Blood-pressure reduction and cardiovascular risk in HOPE study. *Lancet* 2001, 358:2130–2131.
48. Hornig B, Landmesser U, Kohler C, et al.: Comparative effect of ace inhibition and angiotensin II type 1 receptor antagonism on bioavailability of nitric oxide in patients with coronary artery disease: role of superoxide dismutase. *Circulation* 2001, 103:799–805.
49. Prasad A, Koh KK, Schenke WH, et al.: Role of angiotensin II type 1 receptor in the regulation of cellular adhesion molecules in atherosclerosis. *Am Heart J* 2001, 142:248–253.
50. Dahlöf B, Devereux RB, Kjeldsen SE, et al.: Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002, 359:995–1003.
51. Vita JA, Keaney JF Jr: Hormone replacement therapy and endothelial function: the exception that proves the rule? *Arterioscler Thromb Vasc Biol* 2001, 21:1867–1869.
52. Thompson LP, Pinkas G, Weiner CP: Chronic 17beta-estradiol replacement increases nitric oxide-mediated vasodilation of guinea pig coronary microcirculation. *Circulation* 2000, 102:445–451.
53. ●●Rossouw JE, Anderson GL, Prentice RL, et al.: Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002, 288:321–333.
Large randomized study that looked into the role of hormone therapy to prevent chronic disease in women with an intact uterus in the decades of life following menopause.
54. ●●Anderson GL, Limacher M, Assaf AR, et al.: Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* 2004, 291:1701–1712.
Large randomized, double-blind, placebo-controlled trial that tested the effects of the most commonly used postmenopausal hormone therapy preparation in the United States on chronic disease incidence in a healthy population of postmenopausal women.
55. Prentice RL, Langer R, Stefanick ML, et al.: Combined postmenopausal hormone therapy and cardiovascular disease: toward resolving the discrepancy between observational studies and the Women's Health Initiative clinical trial. *Am J Epidemiol* 2005, 162:404–414.
56. Patrono C, Garcia Rodriguez LA, Landolfi R, Baigent C: Low-dose aspirin for the prevention of atherothrombosis. *N Engl J Med* 2005, 353:2373–2383.