

In this article we, with certain restrictions, establish the weak equivalence of distribution functions of the eigenvalues and s-values of a compact operator. We give applications to differential operators, whose spectra cannot be condensed to a ray.

1. Let σ_∞ denote the class of totally continuous operators, defined in the separable Hilbert space H . Denote by $\lambda_n(A)$, $s_n(A)$ ($n = 1, 2, 3, \dots$), respectively, the eigenvalues and s-values of the operator $A \in \sigma_\infty$, numerated in the order of nonincrease of modules. We recall that $s_n^2(A) \leq \lambda_n(A^*A)$.

Denote by the number θ_T , for a linear operator T with region of definition D_T , the angle of the cone $W_T = \{Tf, f\}, f \in D_T\}$.

For the operator $A \in \sigma_\infty$, consider the functions

$$n(r, A) = \sum_{r \mid \lambda_n(A) \geq 1} 1, \quad \tilde{n}(r, A) = \sum_{rs_n(A) \geq 1} 1.$$

Moreover, for $p > 0$ set $\sigma_p(H) = \{A \in \sigma_\infty(H), \sum s_n^p(A) < +\infty\}$.

2. THEOREM 1. Let the dissipative operator $A \in \sigma_p$, where $p \leq \frac{\pi}{2} \theta_A^{-1}$, and let the following condition hold: $\tilde{n}(2r, A) = O(1) \tilde{n}(r, A)$. Then $\tilde{n}(r, A) \asymp n(r, A)$.

Remark. The example of the operator of fractional integration (see [1]) shows that the condition $A \in \sigma_p$, $p \leq \frac{\pi}{2} \theta_A^{-1}$, in Theorem 1 can never be replaced by the weaker statement $s_n(A) = O(n^{-1/2p})$, $p \leq \frac{\pi}{2} \theta_A^{-1}$.

3. Using the results of [2, §2], the proof of the theorem reduces to the case of integral p . Without loss of generality, we may assume that the values of the form $z = (Af, f)$ lie in the angle $0 \leq \arg z \leq \pi/2p$. Since $T = AP$ is a dissipative operator, it follows from Theorem 2 of [2] that $A = T^{1/p}$, in the sense of [2]. Applying once more the results of [2, §2], the proof reduces to the case $p = 1$, such that in this case the operator $B = A^{-1}$ exists, and $\operatorname{Re}(B + tE)^{-1} \geq 0$, $\operatorname{Im}(B + tE)^{-1} \geq 0$ for all $t > 0$. From the above inequalities we have

$$\|(B + tE)^{-1}\|_1 \leq 2 \int (t + \lambda)^{-1} d n(\lambda, A).$$

Hence, and since the operator $(B + tE)(\sqrt{B^*B} + tE)^{-1}$ is uniformly bounded in $t \in R_1^+$, we obtain the estimate

$$\int (\lambda + t)^{-1} d \tilde{n}(\lambda, A) \asymp \int (\lambda + t)^{-1} d n(\lambda, A),$$

since from the condition $\tilde{n}(2r, A) = O(1) \tilde{n}(r, A)$ it follows from the results of [3] that $n(r, A) = O(\tilde{n}(r, A))$. Moreover, the Tauber theorems apply.

4. Let T and L be invertible closed operators with discrete spectra; $T^* = T \geq 0$. Set $N(\lambda, T) = n(\lambda, T^{-1})$, $N(\lambda, L) = n(\lambda, L^{-1})$, $k(\lambda, T) = \tilde{n}(\lambda, T^{-1})$. Suppose that $D_T = D_L$ and for all $u \in D_T$ we have the inequalities

$$c_1 |Tu| \leq |Lu| \leq c_2 |Tu| \quad (1)$$

with some constants $c_1, c_2 > 0$. Since the function $k(\lambda, L)$ coincides with the maximal (in dimension of the linear manifolds M such that $|Lu| \leq \lambda|u|$ for all $u \in M$, from (1) we have the inequalities

$$N(c_2^{-1}\lambda, T) \leq k(\lambda, L) \leq N(c_1^{-1}\lambda, T).$$

Hence, if $N(2\lambda, T) = O(1)N(\lambda, T)$ then $N(\lambda, T) \asymp k(\lambda, L)$. If, moreover, the resolvent of the operator T belongs to the class σ_p , where $p \leq \frac{\pi}{2}\theta_L^{-1}$, and $\theta_L \leq \pi$, from Theorem 1 we obtain $N(\lambda, L) \asymp N(\lambda, T)$.

5. Let Ω be a bounded region in R_n . Denote by $\hat{H}_m(\Omega)$, where m is an integer, the closure of $C_0^\infty(\Omega)$ in the norm $|u|_m = \sum_{|\alpha| \leq m} |D^\alpha u|_{L_2(\Omega)}$, where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n)$, and $D^\alpha = \left(\frac{1}{i} \frac{\partial}{\partial x_1}\right)^{\alpha_1} \dots \left(\frac{1}{i} \frac{\partial}{\partial x_n}\right)^{\alpha_n}$. Let L denote the closure in $L_2(\Omega)$ of the operator

$$L_0 = \sum_{|\alpha|, |\beta| \leq m} D^\alpha (a_{\alpha\beta}(\cdot) D^\beta), \quad D_{L_0} = C^\infty(\Omega) \cap \hat{H}_m(\Omega),$$

where $a_{\alpha\beta}(\cdot) \in C^\infty(\Omega)$, for all $|\alpha|, |\beta| \leq m$. Suppose that the values of the function $z(x, s) = \sum_{|\alpha|=|\beta|=m} a_{\alpha\beta}(x) s^\alpha s^\beta$ ($x \in \Omega, s \in R_n$) form an angle of $\theta < \min\left(\frac{\pi m}{n}, \pi\right)$, where $z(x, s) \neq 0$ if $s \neq 0$.

THEOREM 2. For the conditions formulated above, the operator L has a discrete spectrum, and $N(\lambda, L) \asymp \lambda^{n/2m}$.

6. Consider the Sturm-Liouville operator $L = -\frac{d^2}{dt^2} + q(t)$ in $L_2(R_1^+)$, with boundary condition $y(0) = 0$. Suppose that $|\operatorname{Im} q(t)| \leq \operatorname{tg} \frac{\pi}{4p} \operatorname{Re} q(t)$ and $|q(t)|^{0.5-p} \in L_1(R_1^+)$, where $p > 1/2$. Let $\varphi(2\lambda) = O(\varphi(\lambda))$, where $\varphi(\lambda) = \operatorname{mes}\{t \mid |q(t)| \leq \lambda\}$ and $q'(t) = o(|q(t)|^{1.5})$.

THEOREM 3. For the conditions formulated above the operator L has discrete spectrum, and $N(\lambda, L) \asymp \sqrt{\lambda \varphi(\lambda)}$.

In connection with Theorem 3, we note [4-6], in which the asymptotics of the spectrum of a Sturm-Liouville operator with complex potential are also studied.

The author thanks A. G. Kostyuchenko and G. V. Radzievskii for their constant attention to this work, and also thanks the reviewer for his useful remarks.

LITERATURE CITED

1. I. C. Gohberg and M. G. Krein, Introduction to the Theory of Nonselfadjoint Operators, Amer. Math. Soc. (1969).
2. V. I. Matsaev and Yu. A. Palant, Ukr. Mat. Zh., 74, No. 3, 329-337 (1962).
3. A. S. Markus and V. I. Paraska, Izv. Mold. SSR, 7, 101-104 (1965).
4. M. V. Fedoryuk, I, Differents. Uravn., 8, No. 5, 811-816 (1972).
5. M. V. Fedoryuk and G. V. Zhdanova, II, Differents. Uravn., 10, No. 6, 1067-1073 (1974).
6. S. I. Chelkak, Questions in the Dynamic Theory of the Propagation of Seismic Waves [in Russian], Leningrad (1974), pp. 161-177.