

Treatment of midportion Achilles tendinopathy: an evidence-based overview

Ruben Zwiers · Johannes I. Wiegeler ·
C. Niek van Dijk

Received: 2 September 2014 / Accepted: 22 October 2014

© European Society of Sports Traumatology, Knee Surgery, Arthroscopy (ESSKA) 2014

Abstract In Achilles tendinopathy, differentiation should be made between paratendinopathy, insertional- and midportion Achilles tendinopathy. Midportion Achilles tendinopathy is clinically characterized by a combination of pain and swelling at the affected site, with impaired performance as an important consequence. The treatment of midportion Achilles tendinopathy contains both non-surgical and surgical options. Eccentric exercise has shown to be an effective treatment modality. Promising results are demonstrated for extracorporeal shockwave therapy. In terms of the surgical treatment of midportion Achilles tendinopathy, no definite recommendations can be made.

Level of evidence IV.

Keywords Achilles tendinopathy · Treatment · Conservative · Surgical

Introduction

In Achilles tendinopathy, differentiation should be made between paratendinopathy, insertional- and midportion Achilles tendinopathy [98]. Midportion Achilles tendinopathy is located 2–7 cm proximal of the Achilles insertion onto the calcaneus. Clinically, it is characterized by a combination of pain and swelling at the affected site, with impaired performance as an important consequence.

The treatment of midportion Achilles tendinopathy contains both non-surgical and surgical options. The initial treatment

consists of non-surgical management, but in approximately 25 % of the patients, a surgical intervention is still required [38, 44, 47]. Regarding non-surgical treatment, several rehabilitations protocols have been developed, with excellent results. Nevertheless, the treatment of midportion Achilles tendinopathy remains challenging. Over the years, many alternative treatment options have been developed. Although there is a lot of literature available, the strength of the evidence is often low. This review will provide an evidence-based overview of the current options in midportion Achilles tendinopathy. This overview may be useful for physicians to make evidence-based decisions in the treatment of Achilles tendinopathy.

Non-surgical non-invasive treatment

The initial treatment for midportion achilles tendinopathy is conservative. Activity adaptation or cessation, in the early phase sometimes with the use of NSAIDs, is often combined with specific training regimes and orthoses to distribute the peak forces on the Achilles tendon. Although commonly administered, the possible healing effect of NSAIDs is modest at best and is not supported by the literature [7] The positive (short term) effect may only lie in its analgesic effect. Regarding activity adaptation or cessation, a study showed allowing continued activity in terms of running and jumping but limiting the amount by the use of a pain-monitoring model is not detrimental for recovery [85]. Eccentric exercises are the mainstay of a prolonged rehabilitation programme [5, 20, 23, 49, 79–81, 104] (Table 1).

Eccentric and concentric training

A range of randomized trials and meta-analyses thereof have evaluated different eccentric training regimes showing

R. Zwiers (✉) · J. I. Wiegeler · C. N. van Dijk
Department of Orthopaedic Surgery, Academic Medical Center,
University of Amsterdam, P.O. Box 22660, 1100 DD Amsterdam,
The Netherlands
e-mail: r.zwiers@amc.uva.nl

beneficial effects. In general, a twelve-week programme is advocated, in which an eccentric exercises programme is developed under supervision of a physical therapist [5, 49, 50, 62, 65, 75, 87]. In addition, six-week programmes have also shown to be beneficial; however, this is a much less studied exercise programme [81]. The current literature on eccentric loading for midportion tendinopathy is graded as strong [82], with consistent statistically significant findings favouring eccentric loading over a night splint, deep friction massage, ultrasound, concentric loading or a wait and see policy [33, 80, 81]. Long-term results are good with some mild residual pain complaints [96]. Recently, the focus has again been shifted to a more pragmatic training regime: a combination of eccentric and concentric exercises [31, 105]. Previous work concluded that concentric was less effective; however, some studies showed good results [76]. This pragmatic regime is generally preferred as the exercises are performed more easily and can be advised in case of bilateral Achilles tendinopathy [58]. Studies evaluating this combination show promising results [31]; however, as some study concluded inferior results, this should be further analysed in the near future [82, 103].

Extracorporeal shock wave therapy

A randomized trial added shock wave therapy (ESWT) to an eccentric loading regime, showing superior results compared to a solitary eccentric loading regime [79]. Other studies have shown beneficial effects of ESWT predominantly in women, and improved outcome after previous non-surgical therapy had failed [28, 53].

The pathway of effect of both eccentric exercise and ESWT remains somewhat of a mystery [78]. Eccentric loading is known to induce a normalization of tendon structure with reduction of neovascularization, albeit it is unknown how this is achieved. For ESWT, the effect may lie in its disturbance of sensory nerve fibres and changes in dorsal root ganglia inducing a decrease of pain sensation. Healing responses may be further induced by changes in transforming growth factor-beta1 (TGF-B1), insulin-like growth factor-1 (IGF-1) and cavitation; however, this has only been studied in animal research.

Other non-surgical treatment options

Deep friction massage has been proposed to aid in the recovery of tendinopathy, but no evidence exists that evaluate the effectiveness of joint mobilization and/or soft tissue massage for midportion Achilles tendinopathy [11]. Stretching of the calf muscles is often recommended for patients with symptoms of the Achilles tendon. The purpose of stretching is to increase muscle-tendon length, reduce stiffness and increase the joint range of motion. Only one study has evaluated the effectiveness of stretching compared with eccentric exercise in patients with Achilles tendinopathy. Both treatment groups improved; however, no ankle range of motion measuring was performed [61]. Research is needed to evaluate whether and which patients benefit from stretching. Tape has been used for long time in patients with Achilles tendinopathy; however, no single study has reported on the effect. One study reported no beneficial effect of the use of kinesio tape (elastic therapeutic

Table 1 Overview of non-surgical non-invasive treatment options

Table 1	Overview of non-surgical non-invasive treatment options	
	Level of evidence	
	Strong evidence supporting use in treatment of midportion Achilles tendinopathy	
	Eccentric exercise	Multiple level I studies
	Moderate evidence	
	ESWT	Multiple level I studies
	Limited evidence	
	Iontophoresis	One level I studies
	Stretching	One level I studies
	GTN	>One level I studies
	Low-level laser	>One level I studies
	No evidence	
	Therapeutic US	One level I studies
	Night splints	>One level I studies
	NSAIDs	One level I studies
	Taping	Level IV studies
	Cryotherapy	Level IV studies
	Deep fraction massage	Level IV studies
	Orthosis	Level IV studies
	Heel lifts	Level IV studies
	ESWT extracorporeal shockwave therapy, GTN topical glycerine trinitrate, Therapeutic US therapeutic ultrasound	
		Beneficial in one study
		Beneficial in one study
		Conflicting results
		Only RTC no benefit
		No benefit in multiple studies
		Only RTC no benefit

ESWT extracorporeal shockwave therapy, GTN topical glycerine trinitrate, *Therapeutic US* therapeutic ultrasound

skin tape) [24]. Decreasing the amount of strain on the tendon by using heel lifts is commonly used; however, evidence is limited. Also, for the beneficial effects of orthotics is limited evidence [82]. One study showed physiotherapy and semirigid insoles to reduce pain after four weeks [57]. The use of night splints, whether or not in addition to eccentric exercise, for treatment of midportion Achilles tendinopathy, has not demonstrated to be of additional benefit [20, 21, 81]. Two systematic reviews support the effect of low-level laser therapy in patients with Achilles tendinopathy [89, 95]. Especially in combination with eccentric exercise, LLLT was showed to be beneficial [88]. It is suggested that therapeutic ultrasound may enhance tendon healing, although other studies did not show positive results of therapeutic ultrasound over placebo for tendinopathy [77, 97]. In the only clinical study on Achilles tendinopathy, a pilot study, both eccentric exercise and therapeutic ultrasound showed acceptable short-term results with no adverse events [14]. Conflicting evidence exists on the use of glyceryl trinitrate patches. A randomized trial and follow-up showed better results in patients treated with glyceryl trinitrate [69, 70]. However, in another trial, no superior results of topical glyceryl trinitrate were found [36]. Little evidence exists on the use of iontophoresis. However, in a randomized controlled trial, iontophoresis with dexamethasone has been shown to be of benefit for patients with Achilles tendinopathy [60].

Injections

Corticosteroid injections

The use of corticosteroid injections is controversial. Although corticosteroid injections are a commonly used treatment for chronic tendinopathies, the rationale for the use of anti-inflammatory injections in Achilles tendinopathy is debated [84]. The injection of corticosteroids around weight-loaded tendons is associated with an increased risk of tendon ruptures [6, 13, 51], although the evidence is not strong. Other studies did not report an increased risk [15, 26, 74]. A recent systematic review concluded that there is no significant evidence to draw any conclusions on the positive effect of local corticosteroid treatment in patients with midportion Achilles tendinopathy [50]. Together with the commonly assumed risk of tendon rupture, the use of corticosteroid injections is not recommended.

High-volume injections

The rationale of high-volume injections lay in its local mechanical effects, causing neo-vessels to stretch, break or occlude. The disruption of these vessels would also

damage the accompanying nerve supply, either by trauma or ischaemia, and therefore decrease symptoms [12]. Preliminary studies have shown that a high-volume injection with normal saline, local anaesthetic and corticosteroid can significantly reduce pain and improve short- and long-term function in patients with Achilles tendinopathy [12, 33, 34, 38–40, 46–49]. A recent study used aprotinin instead of hydrocortisone acetate resulting in significant improvement of the VISA-A score at one year follow-up, as well as a relatively high rate of return to sport. [45].

Sclerosing agents (Polidocanol)

Sclerosing therapy for chronic Achilles tendinopathy is based on the hypothesis that the process of neovascularization in the affected tendon is the source of the patient's pain. Some studies have implicated neovascularization as a possible (indirect) cause of symptoms in Achilles tendinopathy. These findings led to the hypothesis that the destruction of the vessels and accompanying nerves might lead to pain relief [3, 65].

A sclerosing agent is injected under ultrasound- and colour Doppler-guidance into the neovascularization in the anterior area outside the tendon. Retrospective case series with relatively short follow-up demonstrated promising clinical results [2, 42, 63, 64, 86]. However, the largest and most recent retrospective series included 53 tendon injections in patients with midportion Achilles tendinopathy. Only 44 % of the tendons were painless or minimally painful at 6 weeks following treatment, and 53 % of the treated tendons had undergone additional non-operative or surgical treatment [99]. Prospective comparative studies involved small numbers of patients: Polidocanol injections were superior to Lidocaine injections [3], and similar results were found when compared with patients undergoing mini-open surgery [4].

Protease inhibitors (Aprotinin)

Aprotinin is a broad-spectrum protease (including matrix metalloproteinase (MMP)) inhibitor. Initially, it was indicated for open heart and liver surgery to reduce blood loss. It is suggested that by inhibiting the enzymes that break down or degrade tendons, the healing response may be promoted. However, the clinical results of aprotinin injections have been conflicting. Aprotinin appears to have effect for patellar tendinopathy [10], whereas a prospective randomized controlled trial by Brown et al. [9] failed to show any significant benefit over placebo in patients with Achilles tendinopathy. The same group of authors subsequently reported successful treatment of non-insertional Achilles tendinopathy in a non-randomized study [66]. After studies associated aprotinin with an increased risk of complications

or death it was permanently withdrawn in May 2008, except for very restricted research use [52].

Hyperosmolar dextrose

Prolotherapy is based on the concept of creating irritation or injury to stimulate healing of a ligament. Hyperosmolar dextrose is said to initiate a local inflammatory response, causing fibroblast proliferation and subsequent collagen production, although the evidence for this is scant [27]. Dextrose acts by causing an osmotic shock to cells leading to the release of pro-inflammatory substances. Several studies have shown increased ligament mass and thickness after repeated prolotherapy injections [16, 50, 54, 56, 62, 64]. In an uncontrolled pilot study, Maxwell et al. [56] retrospectively analysed 33 patients with midportion and/or insertional Achilles tendinopathy and found that hyperosmotic injections yield good results in pain reduction in both rest and during activity.

A recent randomized clinical trial showed that combination of eccentric exercise with prolotherapy provides rapid improvements compared to eccentric exercise alone, but the long-term VISA-A scores were similar [104].

Low-dose heparin

Low-dose heparin has been used in the management of Achilles tendinopathies with the aim of limiting the formation of adhesions. However, there is no evidence for beneficial effect [38, 39, 64], and it has even been suggested that heparin, in itself, can *cause* degenerative tendinopathy in rats [91].

Deproteinized haemodialysate

Pförringer et al. [73] performed a double-blind, placebo-controlled trial to assess the effect of deproteinized haemodialysate, a filtered extract from calf blood. There was a significant reduction of tendon diameter and a reduction of pain as regards activity. No adverse effects were observed, though the follow-up was short.

Autologous blood

Autologous blood injections are increasingly popular treatment option for tendinopathy. A recent systematic review of these studies concluded that there were no advantages in using autologous blood products in patients with tendinopathies [18]. The proposed mechanism of action is that the cytokines and growth factors within the injected blood help to stimulate tissue healing and production of type I collagen [35].

Hitherto, only two randomized controlled trials are conducted in which peritendinous autologous blood injections

added to a standard eccentric training programme were compared with eccentric training alone. Both studies showed no additional benefit for autologous blood injections [8, 72].

Platelet-rich plasma

Platelet-rich plasma (PRP) is a bioactive component of autologous whole blood, which produced by the centrifugation of whole blood, yielding a concentration of platelets greater than baseline levels. The use of PRP has become widespread in different areas of orthopaedics and sports medicine. Theory behind the use of PRP is to enhance the process of tissue repair through the delivery of bioactive agents to provide chemotactic, proliferative and anabolic cellular responses and ultimately tissue function [25, 32].

Case series showing promising outcomes in patients with midportion Achilles tendinopathy has been published [29, 59, 67]. However, the only level one study regarding the use of PRP for chronic Achilles tendinopathy by de Vos et al. [17, 19] did not show any difference between exercises (standard care) combined with either a PRP injection (PRP group) or saline injection (placebo group). Also, at one year follow-up, the PRP group showed no clinical and ultrasonographic superiority compared with the control group. Caution is warranted when comparing different PRP studies. In literature, different types of PRP or PRP-derived products have been used, with a variety of platelet concentrations, inclusion of leucocytes, the use of anticoagulant and the use of activating agents (Table 2).

Surgical treatment

There is no high-level evidence on the surgical treatment. Literature on this topic consists of, mostly retrospective, case series often with small sample sizes. Surgical treatment for midportion Achilles tendinopathy is indicated after failure of appropriated non-surgical treatment, generally after a period of nonsurgical treatment of at least 6 months. Other factors associated with surgery are increased patient age, prolonged duration of symptoms and the presence of degenerative changes in the Achilles tendon [68]. Paavola et al. [68] reported that 29 % of the patients required surgical intervention. Contraindications to surgery include arterial insufficiency, active skin infection, poor soft tissue envelope and medical comorbidities that may increase the risk of surgical complications.

Over the years, different surgical procedures are described. Historically, surgical procedures consisted of debridement of degenerative nodules and fibrotic adhesions, with or without tenotomies to initiate neoangiogenesis in the tendon and so stimulate the healing response. Based on the theory, ingrowth of neovessels and accompanying nerves at

Table 2 Overview of injectable treatment options for midportion Achilles tendinopathy

Injection type	Substance	Level of evidence	+/-	Conclusion
Corticosteroid injections	Triamcinolone methylprednisolone	Two RCTs, several case series	– Conflicting outcomes – Possible risk on ruptures	Use should not be recommended
High-volume injections	Normal saline hydrocortisone aprotinin	No RCTs, several case series	+ Promising results – Limited evidence	No definitive conclusion possible
Sclerosing agents	Polidocanol	Three RCTs, several case series	+ Promising results in small RCTs – Small sample-size RCTs – Conflicting outcomes	No definitive conclusion possible
Protease inhibitors	Aprotinin	One RCT, some case series	– RCT has shown no benefit over placebo – Withdrawn after the suggestion of severe complications	No role in clinical practice
Prolotherapy	Hyperosmolar dextrose	One RCT, some case series	+ Promising short-term results – Limited evidence	No definitive conclusion possible
Anticoagulants	Low-dose heparin	One RCT	– Small sample size – No difference in comparison with placebo	No definitive conclusion possible
Haemodialysate	Deproteinized haemodialysate from calf blood (Actovegin®)	One RCTs	+ Promising short-term results – Limited evidence	No definitive conclusion possible
Autologous blood	Autologous whole blood	Two RCT	– No significant clinical effect	No definitive conclusion possible
	Platelet-rich plasma	One RCT, several case series	– No significant clinical effect	No definitive conclusion possible

Table 3 Overview of surgical techniques for midportion Achilles tendinopathy

Surgical technique	Rationale	Level of evidence
Open debridement		Level VI studies
	With stripping of paratendon	
	Without stripping	
Percutaneous tenotomies	Initiate neovascularization to stimulate healing response	Level VI studies
Minimally invasive tendon stripping	Denervation of the diseased tendon	Level VI studies
Mini-open “scraping”	Denervation of the diseased tendon	Level VI studies, one Level II (poor quality RCT)
Endoscopic debridement	Denervation of the diseased tendon	Level VI studies
Plantaris tendon release	Denervation of the diseased tendon	Level VI studies
Gastrocnemius recession	Unloading of diseased tendon	Level VI studies

the anterior aspect of the tendon are the presumable cause of pain, denervation of the diseased tendon. The technique used for operative management of tendinopathy depends on the stage of the disease and preference of the surgeon. Local degeneration and thickening are usually treated by excision and curettage. An insufficient Achilles tendon due to extensive degeneration can be reconstructed. Isolated paratendinopathy can be treated by excision or stripping of the affected paratendon (Table 3).

Open debridement

Open surgery consisted of a tenotomy with debridement of abnormal tissue degenerative nodules and fibrotic adhesions, with [41] or without stripping of the paratendon [40].

Usually, the tendon is even after thorough debridement strong enough. In case more than 50 % of the tendon is debrided, augmentation of the tendon using the FHL tendon is recommended [102].

The only systematic review on surgical treatment for midportion Achilles tendinopathy was published in 2001 [90]. The success rate varied widely; however, the mean success rate was 77 %. The outcome of this review should be interpreted with caution, since an inverse relation between study quality and outcome had been shown. Maffulli et al. [48], however, reported that only 36 % had good results after open debridement procedures.

Percutaneous tenotomies

Testa et al. [93] described a technique of percutaneous longitudinal tenotomy in which multiple, ultrasound-guided and percutaneous incisions are made through the diseased areas of the Achilles tendon. In their series, they described up to 75 % good to excellent results, even on the long term [46, 92].

Minimally invasive tendon stripping

Longo et al. [43] described a minimally invasive technique in which they used a large diameter suture to free the Achilles tendon from adhesions in the pre-Achilles fat pad. This technique is based on the precept that the process of neovascularization in the damaged tendon is the source of symptoms; however, no clinical results are published to date.

Mini-open “scraping”

Alfredson et al. [4] introduced a mini-open US and Doppler guided scraping technique based on the relationship between vessels and nerves on the ventral side of the Achilles tendon. In a region of degenerative changes and with a high blood flow, the tendon was completely released from the soft tissue. In addition, the plantaris tendon was frequently released. Good clinical outcomes were reported for this technique [83]. In a randomized controlled trial, the results for polidocinol injections and mini-open scraping technique were similar. In addition, in another randomized trial, no difference was found between mini-open scraping and a percutaneous approach using a needle [1].

Endoscopic debridement

To achieve faster recovery and less postoperative complications, endoscopic techniques have been developed. Steenstra and van Dijk were the first to describe an endoscopic technique to release the paratenon and debridement of the tendon. Good results have been described after endoscopic surgery although series are small and the follow-up is short [54, 55, 71, 94]. The main advantage of this technique is the quick recovery. Return to sport activities takes average 6–8 weeks compared to 5–6 months with tenotomy techniques.

Plantaris tendon

Van Sterkenburg and van Dijk [101] addressed the possible role of the plantaris tendon in midportion Achilles tendon. Small series demonstrated promising results after plantaris tendon release [71, 100].

Another new approach is gastrocnemius recession, first described by Gentchov et al. [30]. Lengthening of the gastrocnemius muscle was thought to decrease the load on the tendinopathy area and relieve symptoms [22, 37].

Conclusion

The most important finding of the present study was the large amount of available literature on treatment of midportion Achilles tendinopathy. The level of evidence of these studies, however, is mainly low. Only for eccentric and concentric exercise, there is strong evidence and is therefore recommended as treatment of first choice in patients with midportion Achilles tendinopathy. For the use extracorporeal shockwave therapy, promising results are reported. Regarding other treatment modalities, more high-level studies are required to make definite recommendations.

Limitation of this study is that it only gives a qualitative overview of the available literature. To actually compare outcomes of the different treatment modalities, a more quantitative evaluation is required.

Concerning the non-surgical management, eccentric-concentric exercise has proven to be an effective treatment modality. Promising results are demonstrated for extracorporeal shockwave therapy in several studies. For all other non-surgical treatment options, evidence is too limited to make reliable recommendations. In terms of the surgical treatment of midportion Achilles tendinopathy, no definite recommendations can be made.

References

1. Alfredson H (2011) Ultrasound and Doppler-guided mini-surgery to treat midportion Achilles tendinosis: results of a large material and a randomised study comparing two scraping techniques. *Br J Sports Med* 45:407–410
2. Alfredson H, Lorentzon R (2007) Sclerosing polidocanol injections of small vessels to treat the chronic painful tendon. *Cardiovasc Hematol Agents Med Chem* 5:97–100
3. Alfredson H, Ohberg L (2005) Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc* 13:338–344
4. Alfredson H, Ohberg L, Zeisig E, Lorentzon R (2007) Treatment of midportion Achilles tendinosis: similar clinical results with US and CD-guided surgery outside the tendon and sclerosing polidocanol injections. *Knee Surg Sports Traumatol Arthrosc* 15:1504–1509

5. Alfredson H, Pietila T, Jonsson P, Lorentzon R (1998) Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 26:360–366
6. Astrom M (1998) Partial rupture in chronic achilles tendinopathy. A retrospective analysis of 342 cases. *Acta Orthop Scand* 69:404
7. Astrom M, Westlin N (1992) No effect of piroxicam on achilles tendinopathy. A randomized study of 70 patients. *Acta Orthop Scand* 63:631–634
8. Bell KJ, Fulcher ML, Rowlands DS, Kerse N (2013) Impact of autologous blood injections in treatment of mid-portion Achilles tendinopathy: double blind randomised controlled trial. *BMJ* 346:f2310
9. Brown R, Orchard J, Kinchington M, Hooper A, Nalder G (2006) Aprotinin in the management of Achilles tendinopathy: a randomised controlled trial. *Br J Sports Med* 40:275–279
10. Capasso G, Testa V, Maffulli N, Bifulco G (1997) Aprotinin, corticosteroids and normosaline in the management of patellar tendinopathy in athletes: a prospective randomized study. *Sports Exerc and Inj* 3:111–115
11. Carcia CR, Martin RL, Houck J, Wukich DK (2010) Achilles pain, stiffness, and muscle power deficits: achilles tendinitis. *J Orthop Sports Phys Ther* 40:A1–A26
12. Chan O, O'Dowd D, Padhiar N, Morrissey D, King J, Jalan R, Maffulli N, Crisp T (2008) High volume image guided injections in chronic Achilles tendinopathy. *Disabil Rehabil* 30:1697–1708
13. Chechick A, Amit Y, Israeli A, Horoszowski H (1982) Recurrent rupture of the achilles tendon induced by corticosteroid injection. *Br J Sports Med* 16:89–90
14. Chester R, Costa ML, Shepstone L, Cooper A, Donell ST (2008) Eccentric calf muscle training compared with therapeutic ultrasound for chronic Achilles tendon pain—a pilot study. *Man Ther* 13:484–491
15. DaCruz DJ, Geeson M, Allen MJ, Phair I (1988) Achilles paratendonitis: an evaluation of steroid injection. *Br J Sports Med* 22:64–65
16. Dagenais S, Yelland MJ, Del MC, Schoene ML (2007) Prolotherapy injections for chronic low-back pain. *Cochrane Database Syst Rev*:CD004059
17. de Jonge S, de Vos RJ, Weir A, van Schie HT, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Tol JL (2011) One-year follow-up of platelet-rich plasma treatment in chronic Achilles tendinopathy: a double-blind randomized placebo-controlled trial. *Am J Sports Med* 39:1623–1629
18. de Vos RJ, van Veldhoven PL, Moen MH, Weir A, Tol JL, Maffulli N (2010) Autologous growth factor injections in chronic tendinopathy: a systematic review. *Br Med Bull* 95:63–77
19. de Vos RJ, Weir A, van Schie HT, Bierma-Zeinstra SM, Verhaar JA, Weinans H, Tol JL (2010) Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 303:144–149
20. de Vos RJ, Weir A, Visser RJ, de Winter WT, Tol JL (2007) The additional value of a night splint to eccentric exercises in chronic midportion Achilles tendinopathy: a randomised controlled trial. *Br J Sports Med* 41:e5
21. de Jonge S, de Vos RJ, van Schie HT, Verhaar JA, Weir A, Tol JL (2010) One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion Achilles tendinopathy. *Br J Sports Med* 44:673–677
22. Duthon VB, Lubbeke A, Duc SR, Stern R, Assal M (2011) Non-insertional Achilles tendinopathy treated with gastrocnemius lengthening. *Foot Ankle Int* 32:375–379
23. Fahlstrom M, Jonsson P, Lorentzon R, Alfredson H (2003) Chronic Achilles tendon pain treated with eccentric calf-muscle training. *Knee Surg Sports Traumatol Arthrosc* 11:327–333
24. Firth BL, Dingley P, Davies ER, Lewis JS, Alexander CM (2010) The effect of kinesiotape on function, pain, and motor-neuronal excitability in healthy people and people with Achilles tendinopathy. *Clin J Sport Med* 20:416–421
25. Foster TE, Puskas BL, Mandelbaum BR, Gerhardt MB, Rodeo SA (2009) Platelet-rich plasma: from basic science to clinical applications. *Am J Sports Med* 37:2259–2272
26. Fredberg U, Bolvig L, Pfeiffer-Jensen M, Clemmensen D, Jakobsen BW, Stengaard-Pedersen K (2004) Ultrasonography as a tool for diagnosis, guidance of local steroid injection and, together with pressure algometry, monitoring of the treatment of athletes with chronic jumper's knee and Achilles tendinitis: a randomized, double-blind, placebo-controlled study. *Scand J Rheumatol* 33:94–101
27. Freeman JW, Empson YM, Ekwueme EC, Paynter DM, Broolinson PG (2011) Effect of prolotherapy on cellular proliferation and collagen deposition in MC3T3-E1 and patellar tendon fibroblast populations. *Transl Res* 158:132–139
28. Furia JP (2008) High-energy extracorporeal shock wave therapy as a treatment for chronic noninsertional Achilles tendinopathy. *Am J Sports Med* 36:502–508
29. Gaweda K, Tarczynska M, Krzyzanowski W (2010) Treatment of Achilles tendinopathy with platelet-rich plasma. *Int J Sports Med* 31:577–583
30. Gentchos CE, Bohay DR, Anderson JG (2008) Gastrocnemius recession as treatment for refractory achilles tendinopathy: a case report. *Foot Ankle Int* 29:620–623
31. Gravare SK, Brorsson A, Karlsson J (2014) Rehabiliteringstraning ger barst effekt vid akillestendinopati. *Lakartidningen* 111
32. Hall MP, Band PA, Meislin RJ, Jazrawi LM, Cardone DA (2009) Platelet-rich plasma: current concepts and application in sports medicine. *J Am Acad Orthop Surg* 17:602–608
33. Herrington L, McCulloch R (2007) The role of eccentric training in the management of Achilles tendinopathy: a pilot study. *Phys Ther Sport* 8:191–196
34. Humphrey J, Chan O, Crisp T, Padhiar N, Morrissey D, Twycross-Lewis R, King J, Maffulli N (2010) The short-term effects of high volume image guided injections in resistant non-insertional Achilles tendinopathy. *J Sci Med Sport* 13:295–298
35. Kajikawa Y, Morihara T, Sakamoto H, Matsuda K, Oshima Y, Yoshida A, Nagae M, Arai Y, Kawata M, Kubo T (2008) Platelet-rich plasma enhances the initial mobilization of circulation-derived cells for tendon healing. *J Cell Physiol* 215:837–845
36. Kane TP, Ismail M, Calder JD (2008) Topical glyceryl trinitrate and noninsertional Achilles tendinopathy: a clinical and cellular investigation. *Am J Sports Med* 36:1160–1163
37. Kiewiet NJ, Holthuisen SM, Bohay DR, Anderson JG (2013) Gastrocnemius recession for chronic noninsertional Achilles tendinopathy. *Foot Ankle Int* 34:481–485
38. Kvist M (1991) Achilles tendon injuries in athletes. *Ann Chir Gynaecol* 80:188–201
39. Larsen AI, Egfjord M, Jelsdorff HM (1987) Low-dose heparin in the treatment of calcaneal peritendinitis. *Scand J Rheumatol* 16:47–51
40. Leach RE, Schepsis AA, Takai H (1992) Long-term results of surgical management of Achilles tendinitis in runners. *Clin Orthop Relat Res* (282):208–212
41. Leppilahti J, Karpakka J, Gorra A, Puranen J, Orava S (1994) Surgical treatment of overuse injuries to the Achilles tendon. *Clin J Sport Med* 4:100–107
42. Lind B, Ohberg L, Alfredson H (2006) Sclerosing polidocanol injections in mid-portion Achilles tendinosis: remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 14:1327–1332

43. Longo UG, Ramamurthy C, Denaro V, Maffulli N (2008) Minimally invasive stripping for chronic Achilles tendinopathy. *Disabil Rehabil* 30:1709–1713
44. Maffulli N (2005) Augmented repair of acute Achilles tendon ruptures using gastrocnemius-soleus fascia. *Int Orthop* 29:134
45. Maffulli N, Spiezia F, Longo UG, Denaro V, Maffulli GD (2013) High volume image guided injections for the management of chronic tendinopathy of the main body of the Achilles tendon. *Phys Ther Sport* 14(3):163–167
46. Maffulli N, Testa V, Capasso G, Bifulco G, Binfield PM (1997) Results of percutaneous longitudinal tenotomy for Achilles tendinopathy in middle- and long-distance runners. *Am J Sports Med* 25:835–840
47. Maffulli N, Walley G, Sayana MK, Longo UG, Denaro V (2008) Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 30:1677–1684
48. Maffulli N, Binfield PM, Moore D, King JB (1999) Surgical decompression of chronic central core lesions of the Achilles tendon. *Am J Sports Med* 27:747–752
49. Mafi N, Lorentzon R, Alfredson H (2001) Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc* 9:42–47
50. Magnusson RA, Dunn WR, Thomson AB (2009) Nonoperative treatment of midportion Achilles tendinopathy: a systematic review. *Clin J Sport Med* 19:54–64
51. Mahler F, Fritschy D (1992) Partial and complete ruptures of the Achilles tendon and local corticosteroid injections. *Br J Sports Med* 26:7–14
52. Mangano DT, Tudor IC, Dietzel C (2006) The risk associated with aprotinin in cardiac surgery. *N Engl J Med* 354:353–365
53. Mani-Babu S, Morrissey D, Waugh C, Screen H, Barton C (2014) The effectiveness of extracorporeal shock wave therapy in lower limb tendinopathy: a systematic review. *Am J Sports Med*. doi:10.1177/0363546514531911
54. Maquirriain J (2013) Surgical treatment of chronic achilles tendinopathy: long-term results of the endoscopic technique. *J Foot Ankle Surg* 52:451–455
55. Maquirriain J, Ayerza M, Costa-Paz M, Muscolo DL (2002) Endoscopic surgery in chronic achilles tendinopathies: a preliminary report. *Arthroscopy* 18:298–303
56. Maxwell NJ, Ryan MB, Taunton JE, Gillies JH, Wong AD (2007) Sonographically guided intratendinous injection of hyperosmolar dextrose to treat chronic tendinosis of the Achilles tendon: a pilot study. *AJR Am J Roentgenol* 189:W215–W220
57. Mayer F, Hirschmuller A, Muller S, Schubert M, Baur H (2007) Effects of short-term treatment strategies over 4 weeks in Achilles tendinopathy. *Br J Sports Med* 41:e6
58. Meyer A, Tumilty S, Baxter GD (2009) Eccentric exercise protocols for chronic non-insertional Achilles tendinopathy: how much is enough? *Scand J Med Sci Sports* 19:609–615
59. Monto RR (2012) Platelet rich plasma treatment for chronic Achilles tendinosis. *Foot Ankle Int* 33:379–385
60. Neeter C, Thomee R, Silbernagel KG, Thomee P, Karlsson J (2003) Iontophoresis with or without dexamethazone in the treatment of acute Achilles tendon pain. *Scand J Med Sci Sports* 13:376–382
61. Norregaard J, Larsen CC, Bieler T, Langberg H (2007) Eccentric exercise in treatment of Achilles tendinopathy. *Scand J Med Sci Sports* 17:133–138
62. Ohberg L, Alfredson H (2004) Effects on neovascularisation behind the good results with eccentric training in chronic mid-portion Achilles tendinosis? *Knee Surg Sports Traumatol Arthrosc* 12:465–470
63. Ohberg L, Alfredson H (2003) Sclerosing therapy in chronic Achilles tendon insertional pain-results of a pilot study. *Knee Surg Sports Traumatol Arthrosc* 11:339–343
64. Ohberg L, Alfredson H (2002) Ultrasound guided sclerosis of neovessels in painful chronic Achilles tendinosis: pilot study of a new treatment. *Br J Sports Med* 36:173–175
65. Ohberg L, Lorentzon R, Alfredson H (2004) Eccentric training in patients with chronic Achilles tendinosis: normalised tendon structure and decreased thickness at follow up. *Br J Sports Med* 38:8–11
66. Orchard J, Massey A, Brown R, Cardon-Dunbar AI, Hofmann J (2008) Successful management of tendinopathy with injections of the MMP-inhibitor Aprotinin. *Clin Orthop Relat Res* 466:1625
67. Owens RF Jr, Ginnetti J, Conti SF, Latona C (2011) Clinical and magnetic resonance imaging outcomes following platelet rich plasma injection for chronic midsubstance Achilles tendinopathy. *Foot Ankle Int* 32:1032–1039
68. Paavola M, Kannus P, Orava S, Pasanen M, Jarvinen M (2002) Surgical treatment for chronic Achilles tendinopathy: a prospective seven month follow up study. *Br J Sports Med* 36:178–182
69. Paoloni JA, Appleyard RC, Nelson J, Murrell GA (2004) Topical glyceryl trinitrate treatment of chronic noninsertional achilles tendinopathy. A randomized, double-blind, placebo-controlled trial. *J Bone Joint Surg Am* 86-A:916–922
70. Paoloni JA, Murrell GA (2007) Three-year followup study of topical glyceryl trinitrate treatment of chronic noninsertional Achilles tendinopathy. *Foot Ankle Int* 28:1064–1068
71. Pearce CJ, Carmichael J, Calder JD (2012) Achilles tendinopathy and plantaris tendon release and division in the treatment of non-insertional Achilles tendinopathy. *Foot Ankle Surg* 18:124–127
72. Pearson J, Rowlands D, Highet R (2011) Autologous blood injection for treatment of Achilles tendinopathy? A randomised controlled trial. *J Sport Rehabil*
73. Pförringer W, Pfister A, Kuntz GÄ (1994) The treatment of Achilles paratendinitis: results of a double-blind, placebo-controlled study with a deproteinized hemodialysate. *Clin J Sport Med* 4:92–99
74. Read MT (1999) Safe relief of rest pain that eases with activity in achillodynia by intrabursal or peritendinous steroid injection: the rupture rate was not increased by these steroid injections. *Br J Sports Med* 33:134
75. Rees JD, Lichtwark GA, Wolman RL, Wilson AM (2008) The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology (Oxford)* 47:1493–1497
76. Rees JD, Wolman RL, Wilson A (2009) Eccentric exercises; why do they work, what are the problems and how can we improve them? *Br J Sports Med* 43:242–246
77. Robertson VJ, Baker KG (2001) A review of therapeutic ultrasound: effectiveness studies. *Phys Ther* 81:1339–1350
78. Rompe JD, Furia J, Maffulli N (2008) Eccentric loading compared with shock wave treatment for chronic insertional achilles tendinopathy. A randomized, controlled trial. *J Bone Joint Surg Am* 90:52–61
79. Rompe JD, Furia J, Maffulli N (2009) Eccentric loading versus eccentric loading plus shock-wave treatment for midportion achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 37:463–470
80. Rompe JD, Nafe B, Furia JP, Maffulli N (2007) Eccentric loading, shock-wave treatment, or a wait-and-see policy for tendinopathy of the main body of tendo Achillis: a randomized controlled trial. *Am J Sports Med* 35:374–383
81. Roos EM, Engstrom M, Lagerquist A, Soderberg B (2004) Clinical improvement after 6 weeks of eccentric exercise in

- patients with mid-portion Achilles tendinopathy—a randomized trial with 1-year follow-up. *Scand J Med Sci Sports* 14:286–295
82. Rowe V, Hemmings S, Barton C, Malliaras P, Maffulli N, Morrissey D (2012) Conservative management of midportion Achilles tendinopathy: a mixed methods study, integrating systematic review and clinical reasoning. *Sports Med* 42:941–967
 83. Ruergard A, Alfredson H (2014) Major physical but also psychological effects after pain relief from surgical scraping in patients with Achilles tendinopathy: a 1-year follow-up study. *Pain Stud Treat* 2:21
 84. Shrier I, Matheson GO, Kohl HW III (1996) Achilles tendonitis: are corticosteroid injections useful or harmful? *Clin J Sport Med* 6:245–250
 85. Silbernagel KG, Thomee R, Eriksson BI, Karlsson J (2007) Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med* 35:897–906
 86. Speed CA (2007) Injection therapies for soft-tissue lesions. *Best Prac Res Clin Rheumatol* 21:333–347
 87. Stanish WD, Rubinovich RM, Curwin S (1986) Eccentric exercise in chronic tendinitis. *Clin Orthop Relat Res* 208:65–68
 88. Stergioulas A, Stergioula M, Aarskog R, Lopes-Martins RA, Bjordal JM (2008) Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *Am J Sports Med* 36:881–887
 89. Sussmilch-Leitch SP, Collins NJ, Bialocerkowski AE, Warden SJ, Crossley KM (2012) Physical therapies for Achilles tendinopathy: systematic review and meta-analysis. *J Foot Ankle Res* 5:15
 90. Tallon C, Coleman BD, Khan KM, Maffulli N (2001) Outcome of surgery for chronic Achilles tendinopathy. A critical review. *Am J Sports Med* 29:315–320
 91. Kosay C, Baran O, Ozcan E, Ozer E, Ulukus C (2001) Effect of heparin on tendon degeneration: an experimental study on rats. *Knee Surg Sports Traumatol Arthrosc* 9:247–253
 92. Testa V, Capasso G, Benazzo F, Maffulli N (2002) Management of Achilles tendinopathy by ultrasound-guided percutaneous tenotomy. *Med Sci Sports Exerc* 34:573–580
 93. Testa V, Maffulli N, Capasso G, Bifulco G (1996) Percutaneous longitudinal tenotomy in chronic Achilles tendonitis. *Bull Hosp Jt Dis* 54:241–244
 94. Thermann H, Benetos IS, Panelli C, Gavrilidis I, Feil S (2009) Endoscopic treatment of chronic mid-portion Achilles tendinopathy: novel technique with short-term results. *Knee Surg Sports Traumatol Arthrosc* 17:1264–1269
 95. Tumilty S, Munn J, McDonough S, Hurley DA, Basford JR, Baxter GD (2010) Low level laser treatment of tendinopathy: a systematic review with meta-analysis. *Photomed Laser Surg* 28:3–16
 96. van der Plas A, de Jonge S, de Vos RJ, van der Heide HJ, Verhaar JA, Weir A, Tol JL (2012) A 5-year follow-up study of Alfredson's heel-drop exercise programme in chronic midportion Achilles tendinopathy. *Br J Sports Med* 46:214–218
 97. van der Windt D, van der Heijden G, van den Berg S, ter Riet G, de Winter A, Bouter L (1999) Ultrasound therapy for musculoskeletal disorders: a systematic review. *Pain* 81:257–271
 98. van Dijk CN, van Sterkenburg MN, Wieringer JI, Karlsson J, Maffulli N (2011) Terminology for Achilles tendon related disorders. *Knee Surg Sports Traumatol Arthrosc* 19(5):835–841
 99. van Sterkenburg MN, de Jonge MC, Siersevelt IN, van Dijk CN (2010) Less promising results with sclerosing etoxysclerol injections for midportion Achilles tendinopathy: a retrospective study. *Am J Sports Med* 38(11):2226–2232
 100. van Sterkenburg MN, Kerkhoffs GM, van Dijk CN (2011) Good outcome after stripping the plantaris tendon in patients with chronic mid-portion Achilles tendinopathy. *Knee Surg Sports Traumatol Arthrosc* 19:1362–1366
 101. van Sterkenburg MN, van Dijk CN (2011) Mid-portion Achilles tendinopathy: why painful? An evidence-based philosophy. *Knee Surg Sports Traumatol Arthrosc* 19:1367–1375
 102. Wapner KL, Taras JS, Lin SS, Chao W (2006) Staged reconstruction for chronic rupture of both peroneal tendons using Hunter rod and flexor hallucis longus tendon transfer: a long-term followup study. *Foot Ankle Int* 27:591–597
 103. Woodley BL, Newsham-West RJ, Baxter GD (2007) Chronic tendinopathy: effectiveness of eccentric exercise. *Br J Sports Med* 41:188–198
 104. Yelland MJ, Sweeting KR, Lyftogt JA, Ng SK, Scuffham PA, Evans KA (2011) Prolotherapy injections and eccentric loading exercises for painful Achilles tendinosis: a randomised trial. *Br J Sports Med* 45:421–428
 105. Yu J, Park D, Lee G (2013) Effect of eccentric strengthening on pain, muscle strength, endurance, and functional fitness factors in male patients with achilles tendinopathy. *Am J Phys Med Rehabil* 92:68–76