

# Protection of chickens against experimental fowl typhoid using a *nuoG* mutant of *Salmonella* serotype Gallinarum

Li Zhang-Barber\*†§, Arthur K. Turner\*‡, Gordon Dougan† and Paul A. Barrow\*

*A nuoG mutation in NADH dehydrogenase I was introduced into a virulent strain of Salmonella serotype Gallinarum, the causative agent of fowl typhoid, using gene replacement with a nuoG open reading frame inactivated by insertion of DNA encoding a kanamycin resistance determinant. The S. Gallinarum nuoG mutant, named SG9NGK, was highly attenuated in chickens. SG9NGK colonized the caeca of chickens less efficiently than the S. Gallinarum parental strain, was less invasive and showed no evidence of multiplication in the liver or spleen. Using a single oral immunization with live bacteria SG9NGK reduced mortality in 2-week-old chickens following challenge with virulent S. Gallinarum from 75% to less than 8%. © 1998 Elsevier Science Ltd. All rights reserved*

**Keywords:** *Salmonella* Gallinarum; *nuoG* mutant; vaccine

*Salmonella* serotype Gallinarum is a cause of fowl typhoid in chickens, a systemic disease primarily involving the reticuloendothelial system<sup>1</sup>. The course of infection is either acute or chronic. Traditionally, control of *S. Gallinarum* and the biotype *S. Pullorum*, in flocks has involved elimination of carriers identified by serological detection, followed by improvements in hygiene and management to prevent re-infection. However, in countries where the ambient temperature necessitates the use of open-sided housing, the possibilities for control are relatively limited. Thus, although *Salmonella* serotype Gallinarum has been largely eradicated from some countries<sup>2</sup>, it is still of considerable, and in some cases increasing, economic importance in many areas, including large parts of Latin America, the Middle East, the Pacific rim, Africa, and some parts of Southern Europe<sup>3–5</sup>. It is likely, therefore, that an effective live vaccine would be of considerable practical value.

A considerable amount of work has already been carried out into the feasibility of controlling fowl

typhoid by vaccination. Early work on the development of vaccines against *S. Gallinarum*, showed killed vaccines were of little value<sup>6,7</sup>. The live *S. Gallinarum* 9R vaccine was found to produce strong protection against experimental oral infection with a virulent *S. Gallinarum* strain<sup>7</sup>, but it possessed residual virulence in some breeds of chicken<sup>8–10</sup>. In addition, the attenuating lesion of the *S. Gallinarum* 9R vaccine has not been identified. A strain of *S. Gallinarum* cured of the virulence-associated plasmid showed some attenuation, but was less protective than the 9R strain<sup>11</sup>. The development of live *Salmonella* vaccines for use in humans and other animals has resulted in the identification of a number of attenuating mutations<sup>12–17</sup>, and some of these have been evaluated in *S. Gallinarum*. An *aroA* mutant of *S. Gallinarum* was shown to be attenuated for two-week-old chickens inoculated intramuscularly or orally. Although no immunity developed following oral administration of the vaccine, some protection followed intra-muscular inoculation, but this was less than with the 9R strain<sup>2</sup>.

Previous studies with the 9R vaccine strain indicated that this vaccine could induce some protection also against experimental *S. Enteritidis* infection in laying hens<sup>18,19</sup>. Given the extent of *S. Enteritidis* infection in the poultry industry world-wide, there could be considerable advantage in using a vaccine strain which was effective against two serotypes of major importance.

During investigations in our laboratory into aspects of intestinal colonization in chickens, a *TnphoA* mutant

\*Institute for Animal Health, Compton, Newbury Berkshire RG20 7NN, UK. †Department of Biochemistry, Imperial College of Science, Technology and Medicine, London SW7 2AZ, UK. ‡Present address: Peptide Therapeutics, 321 Cambridge Science Park, Cambridge CB4 4WG, UK. §Author to whom all correspondence should be addressed. Tel.: 44 171 594 5254; email: l.z.barber@i-c.ac.uk. (Received 31 July 1997; accepted 18 November 1997)

of the *S. Typhimurium* strain F98 was obtained which showed reduced virulence in one-day-old chickens compared with the wild-type (Turner *et al.*, in preparation). The *TnphoA* insertion site was found to be located within *nuoG*, a gene in the *nuo* locus encoding NADH dehydrogenase I<sup>20,21</sup>. It was considered worthwhile to assess the virulence of a *nuoG* mutant of *S. Gallinarum* as well as the protective properties against fowl typhoid. This paper describes the construction of a defined non-polar *nuoG* mutant of *S. Gallinarum* which combines the desired phenotype of attenuation and immunogenicity, and reports on the potential of the strain as a live vaccine against fowl typhoid.

## MATERIALS AND METHODS

### Bacterial strains, plasmids and culture media

A nalidixic acid resistant (Nal<sup>r</sup>) derivative of *S. Gallinarum* strain 9 was used to construct the *nuoG* mutant, and as the challenge strain in protection studies. This strain is highly virulent for chickens<sup>1,22,23</sup>. *E. coli* MC1016 harbouring plasmid pSFDK1 containing a non-polar kanamycin cassette was obtained from Dr Sansonetti of the Institute Pasteur, France. *E. coli* K12 strains SY327*ipir* and SM10*ipir*<sup>24</sup> were used as the host for suicide vector pGP704-derivatives<sup>24</sup> and for conjugative transfer of the relevant construct to *S. Gallinarum* 9. *E. coli* strains were grown on Luria-Bertani (LB) agar or in LB broth, and *S. Typhimurium* and *S. Gallinarum* strains were grown on MacConkey agar (CM7; Oxoid). Unless otherwise indicated, broth cultures contained  $1-3 \times 10^9$  c.f.u. ml<sup>-1</sup>. Antibiotics were used at the following concentrations: ampicillin, 100 µg ml<sup>-1</sup>; kanamycin, 50 µg ml<sup>-1</sup>; spectinomycin, 30 µg ml<sup>-1</sup>; nalidixic acid, 20 µg ml<sup>-1</sup> and novomycin, 1 µg ml<sup>-1</sup>.

### DNA manipulations and reagents

Plasmid DNA was prepared using the Wizard DNA miniprep kit (Promega, UK) in accordance with the manufacturer's instructions. Restriction endonucleases and DNA-modifying enzymes were obtained from Boehringer (Mannheim, Germany) and used according to the manufacturer's instructions. Polymerase chain reactions (PCR) were performed according to standard procedures<sup>25</sup>.

### Construction of a *nuoG* mutant of *S. Gallinarum*

The defined *nuoG* mutation was constructed by PCR amplifying a 1.259 kbp DNA fragment from the *nuoG* gene of *S. Typhimurium* F98<sup>22,23</sup>. The nucleotide sequence of the 5' primer was 5'-GCGAATTCTCCG-GTAACCT, and 3' primer was 5'-GCTCTA-GAAAGCTTGCCGCGGAAAGCAC. The amplified DNA fragment was digested with restriction endonucleases *EcoRI* and *XbaI* and ligated into pGP704 digested with the same enzymes. A 850 bp non-polar cassette containing a kanamycin resistance determinant, *aphA-3*, was amplified by PCR from plasmid pSFB1<sup>26</sup> using 5' primer (5'TCCCCGGGT-GACTAAGTAGGAGG) and downstream 3' primer (5'-TCCCCCGGGGTCATTATCCCT). The PCR products were digested with *SmaI*, and cloned into the *SmaI* site of the pGP704-*nuoG* fragment derivative

720 bp from the start of the *nuoG* open reading frame. This plasmid was transferred to *S. Gallinarum* strain 9 by conjugation, and kanamycin resistant, ampicillin sensitive transconjugants were selected. One such transconjugant, named SG9NGK, was shown by Southern hybridization<sup>25</sup> and PCR analysis to have incorporated the *nuoG* mutation. SG9NGK was used in all subsequent vaccination experiments described in this report.

### Experimental animals

Three-week-old specific pathogen-free Light Sussex chickens bred at the Institute for Animal Health were used throughout. These are outbred, of moderate sensitivity to salmonellosis<sup>27</sup>, and were housed and reared as described previously<sup>28</sup>.

### Determination of virulence of the SG9NGK

Groups of 24 chickens were housed separately and inoculated orally with 0.3 ml of a 24 h LB broth culture (ca.  $6 \times 10^8$  c.f.u.). Chickens were observed for typical signs of fowl typhoid (anorexia, ruffled feathers, cyanosis and diarrhoea) for a 3-week period, and those severely affected were killed humanely. For the estimation of LD<sub>50</sub> values, groups of five chickens were inoculated intramuscularly with decimal dilutions of broth cultures. Chickens were observed for 1 week and the LD<sub>50</sub> values were calculated<sup>29</sup>.

### Estimation of intestinal invasiveness

To assess the intestinal invasiveness of SG9NGK and the wild-type *S. Gallinarum* strains, two groups of 12 chickens were each inoculated orally with 0.3 ml of broth culture, as above, of SG9NGK or wild type. At intervals, three chickens from each group were killed and viable counts of *Salmonellae* in the liver, spleen, and caeca determined<sup>23</sup>.

### Persistence in the reticuloendothelial system

Groups of 20 or 30 chickens were inoculated intravenously with  $10^4$  c.f.u. of the wild-type strain or  $10^6$  c.f.u. of SG9NGK. Immediately and at intervals thereafter, groups of three chickens were killed and the viable counts of bacteria in the blood, livers and spleens were determined<sup>23</sup>.

### Assessment of protection

Fifteen chickens were inoculated orally with 0.3 ml of an overnight culture of SG9NGK ( $10^9$  c.f.u.), and another group of 12 birds were untreated. After three weeks three birds from the group inoculated with SG9NGK were killed to count the numbers of *Salmonella* in the tissues. All birds were then inoculated orally with 0.3 ml of a 24 h old culture of wild-type *S. Gallinarum* 9, and monitored for symptoms of disease, as above.

## RESULTS

### Comparison of virulence of SG9NGK and *S. Gallinarum* wild-type

*S. Typhimurium* strains harbouring *nuoG* mutations grow more poorly on acetate as the sole carbon

source<sup>30</sup>, and a 24 h old LB broth culture will not suppress the growth of small numbers of wild-type cells inoculated into that culture<sup>31</sup>. SG9NGK was found to display the same phenotypes as the *S. Typhimurium nuoG* mutants (results not shown). When 24 chickens were inoculated with wild-type *S. Gallinarum*, 18 died or were killed humanely within the following 3-week period as a result of fowl typhoid. In contrast, all 24 chickens in the group inoculated with SG9NGK remained healthy. At post-mortem examination, chickens inoculated with the wild-type *S. Gallinarum* strain showed grossly enlarged livers, and pure, heavy growth of the inoculated strain was obtained from this organ. By contrast, examination of the chickens inoculated with SG9NGK at the end of the experiment revealed healthy organs and no growth of the inoculated organisms from liver. By intra-muscular inoculation the log<sub>10</sub> LD<sub>50</sub> for the *S. Gallinarum nuoG* mutant was found to be >7.08 compared to <0.38 for the wild-type, indicating that SG9NGK was significantly attenuated by the *nuoG* mutation.

#### Comparison of invasiveness of SG9NGK and the wild-type *S. Gallinarum*

Wild-type *S. Gallinarum* was detected in the caeca from 6 h to 5 days after oral inoculation while SG9NGK was detected in the caeca up to 12 h post-inoculation, but not thereafter (Table 1). Wild-type *S. Gallinarum* were detectable in the livers 3 days after inoculation, but the numbers of SG9NGK were significantly less than the wild-type. Wild-type *S. Gallinarum* bacteria were detected in the spleens in significant numbers from three days post-inoculation, whereas SG9NGK was not detected at all in this organ (Table 1).

#### Persistence of SG9NGK and wild-type *S. Gallinarum* in the reticuloendothelial system

For this experiment chickens were inoculated with higher numbers of SG9NGK than wild-type *S. Gallinarum* to allow persistence over a longer period to be assessed. Immediately following inoculation, both strains were detected in the livers, spleens and blood of chickens (Table 2). The wild-type *S. Gallinarum* strain multiplied in the spleen and liver and eventually reappeared in the blood, producing a bacteraemia consistent with the later stages of disease. This was accompanied by the appearance of necrotic lesions in the liver and spleen. In contrast, SG9NGK gradually declined in numbers in the liver and spleen, and was not isolated from the blood after 2 days post-inoculation. Despite the gradual decline in numbers of the mutant in these organs, necrotic lesions developed within a few days of inoculation. SG9NGK persisted in the liver and spleen for 42 days after inoculation, when the experiment was terminated (Table 2).

#### Protection afforded by vaccination with SG9NGK

A group of chickens were immunized orally with SG9NGK and a similar control group were left unimmunized. Immediately prior to challenge, no SG9NGK bacteria were detected in the livers or spleens of three birds killed from the vaccinated group. By 3 weeks after challenge with wild-type *S. Gallinarum*, nine birds in the non-vaccinated group had died or been killed because of severe fowl typhoid. All nine birds, and two of the three survivors, had enlarged livers, and pure, heavy growth of wild-type *S. Gallinarum* was obtained from this organ. In the group vaccinated with SG9NGK, none of the birds showed any signs of disease during the course of the experi-

**Table 1** Comparison of intestinal invasiveness of the *S. Gallinarum nuoG* mutant and wild type

Time after infection	Log <sub>10</sub> viable count per gram of organs <sup>a</sup>					
	Liver		Spleen		Caeca	
	<i>nuoG</i> mutant	Wild type	<i>nuoG</i> mutant	Wild type	<i>nuoG</i> mutant	Wild type
6 h	<2	<2	<2	<2	4.6	5.1
12 h	<2	<2	<2	<2	2.9	2.5
3 days	2.6	4.4	<2	4.5	<2	3.3
5 days	2	5.1	<2	4.7	<2	2.8

<sup>a</sup>Mean value from three birds

**Table 2** Comparison of presence in the tissues of the *S. Gallinarum nuoG* mutant and wild-type following intravenous inoculation

Days after infection	Log <sub>10</sub> viable count per gram of organs <sup>a</sup>					
	Liver		Spleen		Blood	
	<i>nuoG</i> mutant	Wild type	<i>nuoG</i> mutant	Wild type	<i>nuoG</i> mutant	Wild type
0	4.5	2.5	6.3	4.3	3.6	0.7
2	5.1	4.6	5.7	5.0	0.8	<0.5
4	4.2 <sup>b</sup>	4.9 <sup>b</sup>	5.6	5.4 <sup>b</sup>	<0.5	1.0
7	3.4 <sup>b</sup>	5.5 <sup>b</sup>	5.3 <sup>b</sup>	5.0 <sup>b</sup>	<0.5	1.7
10	2.7 <sup>b</sup>	5.7 <sup>b</sup>	4.6 <sup>b</sup>	5.6 <sup>b</sup>	<0.5	2.4
14 <sup>c</sup>	3.3 <sup>b</sup>		4.5 <sup>b</sup>		<0.5	
21	3.3 <sup>b</sup>		5.0 <sup>b</sup>		<0.5	
35	2.7		4.5		<0.5	
42	1.7		2.6		<0.5	

<sup>a</sup>Mean value from three birds<sup>b</sup>Necrotic lesions visible<sup>c</sup>At 14 days birds infected with the wild-type were killed

ment. On completion of the experiment, the vaccinated birds were healthy and no bacterial growth was obtained from the livers.

## DISCUSSION

We have introduced a mutation into the *nuoG* gene of *S. Gallinarum* which codes for a subunit of the NADH dehydrogenase I complex. The *nuoG* mutation attenuates this serotype such that inoculation with  $10^7$  viable organisms produced no outward effects. The mechanism for the attenuation is unclear. However, it is known that *nuoG* mutants grow very poorly on acetate as the sole carbon source<sup>30</sup>. Since the NADH dehydrogenase I complex is the first component of the electron transport chain, it is likely that *nuo* mutants are less efficient than the wild-type at extracting energy from non-fermentable carbon sources, and this may be the reason for the attenuation.

Three days after oral inoculation SG9NGK had not colonized the caeca in detectable numbers, unlike the wild-type *S. Gallinarum*. In addition, SG9NGK was less invasive, as indicated by reduced numbers of salmonellae detected in the liver and none detected in the spleen (Table. 1). Unpublished data using the mouse virulent *S. Typhimurium* C5 strain also indicates that the *nuoG* mutation may confer partially reduced invasiveness in tissue culture cells (Barber, unpublished results). Following inoculation of chickens intravenously, SG9NGK exhibited no evidence for multiplication in the liver or spleen, nevertheless, necrotic lesions were observed on these organs several days post-inoculation after the bacterial numbers had begun to decline.

SG9NGK has a number of potential advantages over existing live vaccine candidates<sup>2,11</sup>. The degree of attenuation was similar to that observed with an *S. Gallinarum aroA* mutant<sup>2</sup> ( $LD_{50}$  values in excess of  $10^7$  for both). However, the *S. Gallinarum aroA* strain did not induce a strong protective immunity, reducing mortality after challenge with the wild-type strain from 63% to 30%, and then only after intra-muscular administration. No protection was obtained by oral vaccination with the *S. Gallinarum aroA* strain. The decline in numbers of salmonellae in the liver and spleen was slower for SG9NGK than the *S. Gallinarum aroA* mutant<sup>2</sup>, and this may be the reason for the stronger protection obtained with the former. The 9R *S. Gallinarum* vaccine strain reduced mortality to 11.5% by oral vaccination<sup>2</sup>. Although fewer birds were used in the present experiments none of them showed any signs of illness, and this represents a mortality of less than 8%. Also, here we used 3-week-old rather than 2-week-old birds, but previous experience indicates that this makes little difference to the virulence of this serotype<sup>2,11</sup> (Barrow, unpublished observations). Thus, SG9NGK conferred a degree of protection at least similar to, or possibly better than, the 9R vaccine strain. However, the attenuating lesion in the 9R strain has not been identified, and in this respect SG9NGK is probably more acceptable as a live vaccine.

## ACKNOWLEDGEMENTS

The authors wish to thank Miss K. Page and Mr A. Grey for excellent technical assistance with animal experiments. This study was supported by Ministry of Agriculture, Fisheries and Food and a Wellcome trust grant to G. Dougan.

## REFERENCES

- 1 Barrow, P.A., Huggins, M.B. and Lovell, M.A. Host specificity of *Salmonella* infection in chickens and mice is expressed in vivo primarily at the level of the reticuloendothelial system. *Infect Immun.* 1994, **62**, 4602–4610.
- 2 Griffin, H.G. and Barrow, P.A. Construction of an *aroA* mutant of *Salmonella* serotype Gallinarum: its effectiveness in immunization against experimental fowl typhoid. *Vaccine* 1993, **11**, 457–462.
- 3 Pomeroy, B.S. and Nagaraja, K.V. Fowl typhoid. In: *Poultry Diseases* 9th edn (Eds Calnek, B.W., Barnes, H.J., Beard, C.W., Reid, W.M. and Yoder, H.W.). Iowa State University Press, Ames, 1991, pp. 87–99.
- 4 Silva, E.N. *The Salmonella gallinarum Problem in Central and South America*. American Association Avian Pathologists, Philadelphia, 1985, pp. 150–156.
- 5 Sato, Y., Sato, G. and Tuchili, L. *et al.* Status of *Salmonella gallinarum-pullorum* infections in poultry in Zambia. *Avian Dis.* 1997, **41**, 490–495.
- 6 Hall, W.J., MacDonald, A.D. and Legenhausen, D.H. Studies of fowl typhoid II. Control of the disease. *Poultry Sci.* 1949, **28**, 789–901.
- 7 Smith, H.W. The use of live vaccines in experimental *Salmonella gallinarum* infection in chickens with observations on their interference effect. *J. Hyg.* 1956, **54**, 419–432.
- 8 Gordon, R.F., Garside, J.S. and Tucker, J.F. The use of living attenuated vaccines in the control of fowl typhoid. *Vet. Rec.* 1959, **71**, 300–305.
- 9 Gordon, W.A.M. and Luck, D. A note on the use of the 9R fowl typhoid vaccine in poultry breeding flocks. *Vet. Rec.* 1959, **71**, 926–927.
- 10 Silva, E.N., Snoeyenbos, G.H., Weinack, O.M. and Smyser, C.F. Studies on the use of 9R vaccine of *Salmonella gallinarum* as a vaccine in chickens. *Avian Dis.* 1981, **25**, 38–52.
- 11 Barrow, P.A. Immunity to experimental fowl typhoid in chickens induced by a virulence plasmid-cured derivative of *Salmonella gallinarum*. *Infect. Immun.* 1990, **58**, 2283–2288.
- 12 Collins, F.M. Effect of specific immune mouse serum on the growth of *Salmonella enteritidis* in mice preimmunized with living or ethyl alcohol-killed vaccines. *J. Bacteriol.* 1969, **97**, 676–683.
- 13 Robertsson, J.A., Lindberg, A.A., Hoiseth, S. and Stocker, B.A. *Salmonella typhimurium* infection in calves: protection and survival of virulent challenge bacteria after immunization with live or inactivated vaccines. *Infect. Immun.* 1983, **41**, 742–750.
- 14 Jones, P.W., Dougan, G., Hayward, C., Mackensie, N., Collins, P. and Chatfield, S.N. Oral vaccination of calves against experimental salmonellosis using a double *aro* mutant of *Salmonella typhimurium*. *Vaccine* 1991, **9**, 29–34.
- 15 Curtiss, R. *Attenuated Salmonella Strains as Live Vectors for the Expression of Foreign Antigens*. Marcel Dekker, New York, 1990.
- 16 Dougan, G., Smith, L. and Heffron, F. *Live Bacterial Vaccines and their Application as Carriers for Foreign Antigens*. Academic Press, London, 1989, pp. 271–300.
- 17 Griffin, H.G. Attenuated *Salmonella* as live vaccines: prospects for multivalent poultry vaccines. *W. Poult. Sci. J.* 1991, **47**, 129–140.
- 18 Barrow, P.A., Lovell, M.A. and Berchieri, A. Immunisation of laying hens against *Salmonella enteritidis* with live attenuated vaccines. *Vet. Rec.* 1990, **126**, 241–242.
- 19 Witvliet, M., Vorstermans, T., Scharr, H., Van Empel, P. and Van den Bosch, H. The *Salmonella gallinarum* 9R vaccine: homologous protection and cross protection against *Salmon-*

- ella enteritidis*. In: *Proceedings of the Second International Symposium on Salmonella and Salmonellosis*. ZOOPÔLE développement, Ploufragan, France, 1997, pp. 503–505.
- 20 Anraku, Y. and Gennis, R.B. The aerobic respiratory chain of *Escherichia coli*. *Trends Biochem. Sci.* 1987, **12**, 262–266.
- 21 Anraku, Y. Bacterial electron transport chains. *Ann. Rev. Biochem.* 1988, **57**, 101–132.
- 22 Barrow, P.A., Huggins, M.B., Lovell, M.A. and Simpson, J.M. Observations on the pathogenesis of experimental *Salmonella typhimurium* infection in chickens. *Res. Vet. Sci.* 1987, **42**, 194–199.
- 23 Barrow, P.A., Simpson, J.M., Lovell, M.A. and Binns, M.M. Contribution of *Salmonella gallinarum* large plasmid toward virulence in fowl typhoid. *Infect. Immun.* 1987, **55**, 388–392.
- 24 Simon, R., Priefer, U. and Puhler, A. A broad host range mobilisation system for in vivo genetic engineering: transposon mutagenesis in Gram negative bacteria. *Biotechnology* 1983, **1**, 784–789.
- 25 Sambrook, J., Fritsch, E.F. and Maniatis, T. *Molecular Cloning*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.
- 26 Menard, R., Sansonetti, P.J. and Parsot, C. Nonpolar mutagenesis of the *ipa* genes defines IpaB, IpaC, and IpaD as effectors of *Shigella flexneri* entry into epithelial cells. *J. Bacteriol* 1993, **175**, 5899–5906.
- 27 Bumstead, N., Millard, B.M., Barrow, P. and Cook, J.K.A. *Genetic Basis of Disease Resistance in Chickens*. CAB International, UK, 1991, pp. 10–23.
- 28 Smith, H.W. and Tucker, J.F. The effect of antibiotic therapy on the faecal excretion of *Salmonella typhimurium* by experimentally infected chickens. *J. Hyg.* 1975, **75**, 275–292.
- 29 Reed, L.J. and Muench, H. A simple method of estimating fifty percent endpoints. *Am. J. Hyg.* 1938, **27**, 493–497.
- 30 Archer, C.D., Jin, J. and Elliott, T. Stabilization of a HemA-LacZ hybrid protein against proteolysis during carbon starvation in *atp* mutants of *Salmonella typhimurium*. *J. Bacteriol.* 1996, **178**, 2462–2464.
- 31 Zhang-Barber, L., Turner, A.K., Martin, G., Frankel, G., Dougan, G. and Barrow, P. Influence of genes encoding proton-translocating enzymes on suppression of *Salmonella typhimurium* growth and colonization. *J. Bacteriol.* 1997, **179**, 7186–7190.