

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281226617>

Expeditious Synthesis of 2-Phenylquinazolin-4-amines via a Fe/Cu Relay-Catalyzed Domino Strategy

ARTICLE *in* ORGANIC LETTERS · AUGUST 2015

Impact Factor: 6.36 · DOI: 10.1021/acs.orglett.5b02020 · Source: PubMed

CITATION

1

READS

12

6 AUTHORS, INCLUDING:



Xu Cheng

Central China Normal University

5 PUBLICATIONS 12 CITATIONS

SEE PROFILE


Expeditious Synthesis of 2-Phenylquinazolin-4-amines via a Fe/Cu Relay-Catalyzed Domino Strategy

Feng-Cheng Jia, Zhi-Wen Zhou, Cheng Xu, Qun Cai, Deng-Kui Li, and An-Xin Wu*

Key Laboratory of Pesticide & Chemical Biology, Ministry of Education, College of Chemistry, Central China Normal University, Hubei, Wuhan 430079, P. R. China

Supporting Information

ABSTRACT: A highly efficient Fe/Cu relay-catalyzed domino protocol has been developed for the synthesis of 2-phenylquinazolin-4-amines from commercially available *ortho*-halogenated benzonitriles, aldehydes, and sodium azide. This elegant domino process involved consecutive iron-mediated [3 + 2] cycloaddition, copper-catalyzed S_NAr , reduction, cyclization, oxidation, and copper-catalyzed denitrogenation sequences. The formed structure is the privileged core in drugs and bioactive molecules.



Quinazoline represents an important and abundant class of nitrogen-containing heterocycles.¹ In particular, as one of the diverse quinazoline derivatives, the 4-aminoquinazoline nucleus is exemplified as a privileged structure that exists in many pharmaceutical molecules and biologically active compounds,^{2,3} such as erlotinib (I),^{2a} gefitinib (II),^{2b} prazosin (III),^{2c} and human adenosine A_3 receptor antagonist (IV)^{2d} (Figure 1). In addition, 4-aminoquinazoline derivatives are often used as synthetic intermediates for the direct synthesis of biologically active molecules.^{2d,3}





Figure 1. Selected drugs or biologically active compounds with a 4-aminoquinazoline moiety.

Because of their great value, the synthesis of 4-aminoquinazolines has gained much attention. The current synthetic methods of this skeleton are mainly summarized as the following three types: (i) the nucleophilic addition/cyclization reaction of anthranilonitrile with benzonitriles;^{2d,4} (ii) the coupling/cyclization reaction of 2-bromobenzonitriles with amidines;⁵ and (iii) the S_NAr /cyclization reaction of 2-fluorobenzonitriles with amidines;⁶ Alternatively, 2-substituted 4-aminoquinazolines can be prepared by the decoration of the existing quinazoline nucleus^{3b,7} (Scheme 1). Although these reactions provide efficient access to 4-aminoquinazolines, their applications are

Scheme 1. Synthetic Routes to 2-Substituted 4-Aminoquinazolines



limited by a lack of suitable substrates, poor substitution diversity, and the requirement for harsh reaction conditions. Therefore, the development of effective new methods for the facile construction of 4-aminoquinazolines is highly desirable.

Sodium azide (NaN_3), which was used as a convenient nitrogen source, has been widely applied in organic synthesis.^{8–15} The common functions of NaN_3 mainly includes two types: (i) a 1,3-dipole to react with electron-deficient olefins,⁸ alkynes,⁹ or nitriles¹⁰ and (ii) a coupling partner participating in copper-catalyzed S_NAr reactions.¹¹ Substantial progress has been made in developing domino reactions based on these two fundamental reactions involving NaN_3 .^{12–14} As part of our ongoing efforts toward developing novel copper-catalyzed domino reaction related to sodium azide,¹⁵ herein we present a

Received: July 14, 2015

novel Fe/Cu relay-catalyzed domino strategy for the direct synthesis of pharmaceutically significant 2-phenylquinazolin-4-amine derives from commercially available *ortho*-halogenated benzonitriles, aldehydes, and sodium azide (Scheme 1).

To explore the feasibility of this domino protocol, our study commenced with *o*-bromobenzonitrile (**1a**), benzaldehyde (**2a**), and sodium azide as model substrates to optimize the reaction conditions. Initially, various Lewis acids were screened in view of their potential catalytic activity toward initial [3 + 2] cycloaddition of nitriles with NaN₃ according to the existing literature,¹⁰ and FeCl₃ showed the highest efficiency in the presence of CuI/L-proline in DMF at 110 °C in a sealed vessel under air (Table 1, entries 1–9). Then several solvents were

Table 1. Optimization of the Reaction Conditions^a

entry	catalyst	solvent	temp (°C)	yield ^b (%)
1	CAN	DMF	110	62
2	FeCl ₃	DMF	110	71
3	ZnCl ₂	DMF	110	42
4	AlCl ₃	DMF	110	45
5	InBr ₃	DMF	110	68
6	ZnBr ₂	DMF	110	11
7	Cu(OAc) ₂	DMF	110	trace
8	Pd(OAc) ₂	DMF	110	trace
9	AgNO ₃	DMF	110	trace
10	FeCl ₃	DMSO	110	6
11	FeCl ₃	1,4-dioxane	110	trace
12	FeCl ₃	toluene	110	trace
13	FeCl ₃	DMF	80	53
14	FeCl ₃	DMF	100	64
15	FeCl ₃	DMF	120	70
16		DMF	100	trace
17 ^c	FeCl ₃	DMF	110	trace
18 ^d	FeCl ₃	DMF	110	trace
19 ^e	FeCl ₃	DMF	110	80


^aReactions conditions: **1a** (0.5 mmol), **2a** (0.5 mmol), NaN₃ (2.0 mmol), CuI (10%), L-proline (20%), and catalyst (10%) were heated in 3 mL of solvent in a sealed vessel under air for 12 h. ^bIsolated yield.

^cAbsence of CuI. ^dAbsence of L-proline. ^e30 mol % of FeCl₃ was used.

tested (Table 1, entries 10–12), and DMF proved to be the most effective solvent (Table 1, compare entries 2 and 10–12). Increasing or decreasing the temperature of the reaction did not lead to any further improvements in the yield (Table 1, entries 13–15). A control experiment confirmed that FeCl₃, CuI, and L-proline are indispensable elements in our catalytic system (Table 1, entries 16–18). Slightly improved efficiency was observed when the loading of FeCl₃ was increased from 10 to 30 mol % (Table 1, entry 19). Overall, the optimized reaction conditions were identified as **1a** (0.5 mmol), 1.0 equiv of **2a**, 4.0 equiv of sodium azide, 30 mol % of FeCl₃, 10 mol % of CuI, and 20 mol % of L-proline in 3 mL of DMF at 110 °C in a sealed vessel under air.

With the optimal reaction conditions in hand, we next investigated the scope of the domino process. A variety of aromatic aldehydes bearing different substituents were tested, and the results are summarized in Scheme 2. It was found that the transformation was very general; electron-neutral (4-H, 2-Me, 4-Me), electron-donating (4-OMe, 4-OEt, 3,4-(OMe)₂), and

Scheme 2. Scope of Aryl Aldehydes^{a,b}




^aReaction conditions: **1a** (0.5 mmol), **2** (0.5 mmol), NaN₃ (2.0 mmol), FeCl₃ (0.15 mmol), CuI (0.05 mmol), and L-proline (0.1 mmol) in DMF (3 mL) at 110 °C in a sealed vessel under air for 12 h.

^bIsolated yields.

electron-deficient (3-NO₂) groups were well tolerated, giving the corresponding products in moderate to good yields (42%–82%, **3aa–ag**). To our delight, the optimized conditions were mild enough to allow halo-substituted substrates (67%–73%, **3ah–aj**), which provided the possibility for further functionalization. Furthermore, sterically hindered substrates such as 1-naphthaldehyde and 2-naphthaldehyde were also found to be suitable for this transformation (**3ak–al**, 57% and 74%). Meanwhile, the optimized conditions could be applied to heteroaryl aldehydes including furan-2-carbaldehyde, thiophene-2-carbaldehyde, and thiophene-3-carbaldehyde (**3am–ao**, 65%–78%). Furthermore, the structure of **3aa** was unambiguously determined by X-ray crystallographic analysis (see the Supporting Information).


To further expand the scope of the substrates, a variety of *ortho*-halogenated benzonitriles and aryl aldehydes were then examined. Gratifyingly, electron-neutral (4-Me, 5-Me) groups on the phenyl rings of 2-bromobenzonitriles were compatible and provided the corresponding products in moderate to good yields (Scheme 3, 53–84%, **3ba–cg**). Halogen-substituted 2-bromobenzonitriles (5-F, 5-Cl) also afforded the desired products in moderate yields (Scheme 3, 45% and 67%, **3da** and **3ea**). In addition, other *ortho*-halogenated benzonitriles such as 2-fluorobenzonitrile, 2-chlorobenzonitrile, and 2-iodobenzonitrile all also exhibit good reactivity under the optimized conditions (Scheme 3, 74–82%, **3aa–aa**).

Notably, this method could also be successfully applied in the convenient synthesis of 1-(2-methoxyphenyl)-3-(2-(pyridin-3-yl)quinazolin-4-yl)urea (**IV**), which is a potent and selective human adenosine A₃ receptor antagonist demonstrated by van Muijlwijk-Koezen.^{2d} As shown in Scheme 4, the reaction of *o*-bromobenzonitrile (**1a**) with sodium azide and nicotinaldehyde occurred smoothly under the standard conditions to afford the corresponding products **3ap** in 77% yield. The product **3ap** was

Scheme 3. Scope of *o*-Halogenated Benzonitriles and Aryl Aldehydes^{a,b}

^aReaction conditions: **1** (0.5 mmol), **2a** (0.5 mmol), NaN₃ (2.0 mmol), FeCl₃ (0.15 mmol), CuI (0.05 mmol), and L-proline (0.1 mmol) in DMF (3 mL) at 110 °C in a sealed vessel under air for 12 h.
^bIsolated yields.


Scheme 4. Synthetic Application



subsequently transformed to pharmaceutically active molecular **IV** according to the reported procedure.^{2d}


Having established the scope of our new domino reaction, we turned our attention to evaluate the reaction mechanism. We initially investigated the reaction of *o*-bromobenzonitrile (**1a**) with sodium azide (2 equiv) in DMF in the presence of FeCl₃ at 110 °C for 12 h, which gave 5-(2-bromophenyl)-1*H*-tetrazole (**4**) in 71% yield (Scheme 5a). When 5-(2-bromophenyl)-1*H*-tetrazole (**4**) was treated with benzaldehyde (**2a**) and NaN₃ (2 equiv) in the presence of CuI in DMF at 110 °C in a sealed vessel under air for 12 h, the target product 2-phenylquinazolin-4-amine (**3aa**) was isolated in 84% yield (Scheme 5b). Furthermore, the reactions of 2-(1*H*-tetrazol-5-yl)aniline (**5**) and benzaldehyde (**2a**) were conducted under standard conditions, and the desired product **3aa** was obtained in 83% yield (Scheme 5c). When 5-(2-bromophenyl)-1*H*-tetrazole (**4**) was treated with benzaldehyde (**2a**) and NaN₃ (2 equiv) in the presence of CuI and L-proline in DMF at 80 °C for 6 h, 5-phenyltetrazolo[1,5-*c*]quinazoline (**6**) and 2-phenylquinazolin-4-amine (**3aa**) were obtained in 51% and 27% yields, respectively (Scheme 5d). Next, when 5-phenyltetrazolo[1,5-*c*]quinazoline (**6**) was heated at 110 °C for 12 h in DMF in the presence of CuI and L-proline, the substrate could be converted to the desired product **3aa** in almost quantitative yield (Scheme 5e). Taken together, these control experiments clearly demonstrated that 5-(2-bromophenyl)-1*H*-tetrazole (**4**), 2-(1*H*-tetrazol-5-yl)aniline (**5**), and 5-phenyltetrazolo[1,5-*c*]quinazoline (**6**) may be key intermediates in this reaction.

Scheme 5. Control Experiments



On the basis of the above observations and literature precedent,^{10–19} a possible reaction mechanism of this transformation was represented in Scheme 6. Initially, the sodium 5-

Scheme 6. Possible Mechanism



(2-bromophenyl)tetrazol-1-ide (**A**) was generated through an iron-mediated [3 + 2] cycloaddition of *o*-bromobenzonitrile (**1a**) with NaN₃.¹⁰ Subsequently, intermediate **A** would undergo a copper-catalyzed S_NAr with NaN₃ to afford intermediate **B** in the light of the *ortho*-substituent effect.^{10,16} Coordination of azide to copper, followed by an electrocyclicization with the concomitant release of N₂, would give the Cu(III) complex **D**,¹⁷ which would undergo a reduction with the aid of trace H₂O in DMF to give intermediate 2-(1*H*-tetrazol-5-yl)aniline (**5**).^{11c–e} Next, 2-(1*H*-1,2,3-triazol-5-yl)aniline (**5**) could easily condense with benzaldehyde (**2a**) to give imine intermediate **E**. Then intramolecular nucleophilic attack of nitrogen to imine in **E** followed by oxidative dehydrogenation led to **F**. Eventually, the target product **3aa** was obtained after final copper-catalyzed denitrogenation process.¹⁸ It is also possible that 5-phenyltetrazolo[1,5-*c*]quinazoline (**6**) and 2-(1*H*-tetrazol-5-yl)aniline (**5**) could be formed via a synergistic oxidation–

reduction reaction between intermediates **B** and **F**.¹⁹ Further mechanistic studies of the detailed process of reduction and oxidation in this reaction system are in progress.

In conclusion, we have developed a highly efficient Fe/Cu relay-catalyzed domino reaction for the facile synthesis of pharmaceutically significant 2-phenylquinazolin-4-amines from commercially available *ortho*-halogenated benzonitriles, aldehydes, and sodium azide. This elegant domino process involved consecutive iron-mediated [3 + 2] cycloaddition, copper-catalyzed S_NAr, reduction, cyclization, oxidation, and copper-catalyzed denitrogenation sequences. Notably, sodium azide acted as dual nitrogen source in the construction of these fused N-heterocycles. Moreover, the free NH₂ generated from this reaction can be utilized for further manipulation. Application of this self-sequence strategy utilizing NaN₃ as a simple nitrogen donor for the synthesis of other fascinating N-heterocycles are underway in our laboratory.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b02020.

Crystallographic data of **3aa** (CIF)

Experimental procedures, product characterizations, and copies of the ¹H and ¹³C NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: chwuax@mail.ccnu.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to the National Natural Science Foundation of China (Grant Nos. 21272085 and 21472056) for financial support.

■ REFERENCES

- (1) For selected papers, see: (a) Parhi, A. K.; Zhang, Y.; Saionz, K. W.; Pradhan, P.; Kaul, M.; Trivedi, K.; Pilch, D. S.; LaVoie, E. J. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 4968–4974. (b) Ugale, V. G.; Bari, S. B. *Eur. J. Med. Chem.* **2014**, *80*, 447–501. (c) Juvele, K.; Gallus, J.; Wiese, M. *Bioorg. Med. Chem.* **2013**, *21*, 7858–7873.
- (2) (a) Gundla, R.; Kazemi, R.; Sanam, R.; Muttineni, R.; Sarma, J. A. R. P.; Dayam, R.; Neamati, N. J. *Med. Chem.* **2008**, *51*, 3367–3377. (b) Omar, M. A.; Conrad, J.; Beifuss, U. *Tetrahedron* **2014**, *70*, 3061–3072. (c) Wilson, L. J. *Org. Lett.* **2001**, *3*, 585–588. (d) van Muijlwijk-Koezen, J. E.; Timmerman, H.; van der Goot, H.; Menge, W. M. P. B.; Frijtag von Drabbe Künzel, J.; de Groote, M.; IJzerman, A. P. *J. Med. Chem.* **2000**, *43*, 2227–2238.
- (3) (a) El-Azab, A. S.; Al-Omar, M. A.; Abdel-Aziz, A. A. M.; Abdel-Aziz, N. I.; El-Sayed, M. A. A.; Aleisa, A. M.; Sayed-Ahmed, M. M.; Abdel-Hamide, S. G. *Eur. J. Med. Chem.* **2010**, *45*, 4188–4198. (b) Marvania, B.; Lee, P. C.; Chaniyara, R.; Dong, H.; Suman, S.; Kakadiya, R.; Chou, T. C.; Lee, T. C.; Shah, A.; Su, T. L. *Bioorg. Med. Chem.* **2011**, *19*, 1987–1998.
- (4) Seijas, J. A.; Vázquez-Tato, M. P.; Montserrat Martínez, M. *Tetrahedron Lett.* **2000**, *41*, 2215–2217.
- (5) Yang, X.; Liu, H.; Fu, H.; Qiao, R.; Jiang, Y.; Zhao, Y. *Synlett* **2010**, *2010*, 101–106.
- (6) (a) Cargill, M. R.; Linton, K. E.; Sandford, G.; Yufit, D. S.; Howard, J. A. K. *Tetrahedron* **2010**, *66*, 2356–2362. (b) Yan, S.; Dong, Y.; Peng, Q.; Fan, Y.; Zhang, J.; Lin, J. *RSC Adv.* **2013**, *3*, 5563–5569.
- (7) (a) Ahmad, O. K.; Hill, M. D.; Movassaghi, M. *J. Org. Chem.* **2009**, *74*, 8460–8463. (b) Yadav, M. R.; Grande, F.; Chouhan, B. S.; Naik, P. P.; Giridhar, R.; Garofalo, A.; Neamati, N. *Eur. J. Med. Chem.* **2012**, *48*, 231–243. (c) Itoh, T.; Mase, T. *Tetrahedron Lett.* **2005**, *46*, 3573–3577.
- (8) (a) Quiclet-Sire, B.; Zard, S. Z. *Synthesis* **2005**, *19*, 3319–3326. (b) Zhang, Y.; Li, X.; Li, J.; Chen, J.; Meng, X.; Zhao, M.; Chen, B. *Org. Lett.* **2012**, *14*, 26–29. (c) Quan, X. J.; Ren, Z. H.; Wang, Y. Y.; Guan, Z. H. *Org. Lett.* **2014**, *16*, 5728–5731.
- (9) Li, J.; Wang, D.; Zhang, Y.; Li, J.; Chen, B. *Org. Lett.* **2009**, *11*, 3024–3027.
- (10) For selected papers, see: (a) Gawande, S. D.; Raihan, M. J.; Zanzwar, M. R.; Kavala, V.; Janreddy, D.; Kuo, C. W.; Chen, M. L.; Kuo, T. S.; Yao, C. F. *Tetrahedron* **2013**, *69*, 1841–1848. (b) Kumar, S.; Dubey, S.; Saxena, N.; Awasthi, S. K. *Tetrahedron Lett.* **2014**, *55*, 6034–6038. (c) Mani, P.; Singh, A. K.; Awasthi, S. K. *Tetrahedron Lett.* **2014**, *55*, 1879–1882.
- (11) (a) Zhu, W.; Ma, D. *Chem. Commun.* **2004**, 888–889. (b) Andersen, J.; Madsen, U.; Björklund, F.; Liang, X. *Synlett* **2005**, 2209–2213. (c) Markiewicz, J. T.; Wiest, O.; Helquist, P. *J. Org. Chem.* **2010**, *75*, 4887–4890. (d) Zhao, H.; Fu, H.; Qiao, R. *J. Org. Chem.* **2010**, *75*, 3311–3316. (e) Goriya, Y.; Ramana, C. V. *Tetrahedron* **2010**, *66*, 7642–7650.
- (12) For representative papers of domino reactions initiated by [3 + 2] cycloaddition involving NaN₃, see: (a) Arigela, R. K.; Samala, S.; Mahar, R.; Shukla, S. K.; Kundu, B. *J. Org. Chem.* **2013**, *78*, 10476–10484. (b) Chen, Z. Y.; Wu, M. J. *Org. Lett.* **2005**, *7*, 475–477. (c) Gulevskaya, A. V.; Tyaglyi, A. S.; Pozharskii, A. F.; Nelina-Nemtseva, J. I.; Steglenko, D. V. *Org. Lett.* **2014**, *16*, 1582–1585. (d) Yan, J.; Zhou, F.; Qin, D.; Cai, T.; Ding, K.; Cai, Q. *Org. Lett.* **2012**, *14*, 1262–1265.
- (13) For representative papers related to the formation of organic azide and sequential relay of click reaction with alkynes, see: (a) Pericherla, K.; Jha, A.; Khungar, B.; Kumar, A. *Org. Lett.* **2013**, *15*, 4304–4307. (b) Ackermann, L.; Potukuchi, H. K.; Landsberg, D.; Vicente, R. *Org. Lett.* **2008**, *10*, 3081–3084. (c) Liu, Z.; Zhu, D.; Luo, B.; Zhang, N.; Liu, Q.; Hu, Y.; Pi, R.; Huang, P.; Wen, S. *Org. Lett.* **2014**, *16*, 5600–5603. (d) Qian, W.; Winterheimer, D.; Allen, J. *Org. Lett.* **2011**, *13*, 1682–1685.
- (14) For representative papers related to the formation of organic azide and sequential relay of intramolecular N-atom transfer reaction, see: (a) Shang, X.; Zhao, S.; Chen, W.; Chen, C.; Qiu, H. *Chem. - Eur. J.* **2014**, *20*, 1825–1828. (b) Kumar, M. R.; Park, A.; Park, N.; Lee, S. *Org. Lett.* **2011**, *13*, 3542–3545. (c) Kim, Y.; Kumar, M. R.; Park, N.; Heo, Y.; Lee, S. *J. Org. Chem.* **2011**, *76*, 9577–9583. (d) Goriya, Y.; Ramana, C. V. *Chem. Commun.* **2013**, *49*, 6376–6378. (e) Li, K.; Chen, J.; Li, J.; Chen, Y.; Qu, J.; Guo, X.; Chen, C.; Chen, B. *Eur. J. Org. Chem.* **2013**, *2013*, 6246–6248. (f) Ou, Y.; Jiao, N. *Chem. Commun.* **2013**, *49*, 3473–3475. (g) Goriya, Y.; Ramana, C. V. *Chem. Commun.* **2014**, *50*, 7790–7792.
- (15) Jia, F. C.; Xu, C.; Zhou, Z. W.; Cai, Q.; Li, D. K.; Wu, A. X. *Org. Lett.* **2015**, *17*, 2820–2823.
- (16) For papers involving the *ortho*-substituent effect, see: (a) Yang, D.; Wang, Y.; Yang, H.; Liu, T.; Fu, H. *Adv. Synth. Catal.* **2012**, *354*, 477–482. (b) Jia, F. C.; Xu, C.; Cai, Q.; Wu, A. X. *Chem. Commun.* **2014**, *50*, 9914–9916.
- (17) For papers related to the Cu(III) complex, see: (a) Zhang, H.; Zhao, L.; Wang, D. X.; Wang, M. X. *Org. Lett.* **2013**, *15*, 3836–3839. (b) Huang, H.; Ji, X.; Tang, X.; Zhang, M.; Li, X.; Jiang, H. *Org. Lett.* **2013**, *15*, 6254–6257. (c) Yu, D. G.; Suri, M.; Glorius, F. *J. Am. Chem. Soc.* **2013**, *135*, 8802–8805. (d) Dhar, D.; Tolman, W. B. *J. Am. Chem. Soc.* **2015**, *137*, 1322–1329.
- (18) Shi, L. L.; Yang, H. J.; Jiang, Y. Y.; Fu, H. *Adv. Synth. Catal.* **2013**, *355*, 1177–1184.
- (19) (a) Cenini, S.; Gallo, E.; Penoni, A.; Ragaini, F.; Tollari, S. *Chem. Commun.* **2000**, 2265–2266. (b) Ragaini, F.; Penoni, A.; Gallo, E.; Tollari, S.; Gotti, C. L.; Lapadula, M.; Mangioni, E.; Cenini, S. *Chem. - Eur. J.* **2003**, *9*, 249–259. (c) Nguyen, Q.; Nguyen, T.; Driver, T. G. *J. Am. Chem. Soc.* **2013**, *135*, 620–623.