

## ORIGINAL ARTICLE

# Overview of the Special Projects of National Significance Program's 10 Models of Adolescent HIV Care

ELIZABETH R. WOODS, M.D., M.P.H.

Ten models of adolescent human immunodeficiency virus (HIV) care were funded in 1993 by the Special Projects of National Significance (SPNS) Program, HIV/Acquired Immunodeficiency Virus (AIDS) Bureau, Health Resources and Services Administration, through the Ryan White CARE Act. These models were supported to advance knowledge about the engagement of HIV-positive and at-risk adolescents and young adults in care. This article provides an overview of the SPNS Program's adolescent initiative, which developed and evaluated innovative models of HIV care, and provides background information on and summarizes the 10 models of care. The models are organized into four groups emphasizing different concepts: (a) youth involvement; (b) outreach to bring youth into services; (c) case management and linkage to services; and (d) a comprehensive continuum of care for youth. © Society for Adolescent Medicine, 1998

## KEY WORDS:

HIV care  
Models of care  
Adolescent  
HIV/AIDS  
Special Projects of National Significance Program

In 1993, 10 models of adolescent human immunodeficiency virus (HIV) care were funded by the Special Projects of National Significance (SPNS) Program of

the Health Resources and Services Administration (see Preface to this issue). Because many HIV-positive adults were infected during adolescence and because young people are particularly hard to engage in care, these 10 innovative models were funded to advance the understanding of the issues around connecting adolescents and young adults (ages 12-24 years) to HIV care. This article summarizes the background information leading to the development of the adolescent models, and introduces the articles contained in this issue which describe in detail the 10 models of care.

## *National HIV/Acquired Immunodeficiency Virus (AIDS) Statistics*

National HIV/AIDS statistics indicate that adolescents and young adults represent important risk groups. The Centers for Disease Control and Prevention (CDC) report that as of June 30, 1997, there were 612,078 reported cases of AIDS in the United States, with 2953 in the 13-19-year-old age group, and 22,070 cases in the 20-24-year-old age group (1). Owing to the long delay in developing symptoms associated with HIV infection, many of the young adults were probably infected during adolescence. The male/female ratio for AIDS cases is 1.6:1 for adolescents and 2.8:1 for young adults (20-24 years old), reflecting the increased number of young women with HIV in these age groups compared to adults (1). Estimates indicate that there are as many as two to three HIV-positive youth for every known case of AIDS (2). A 1996 report released by the White House Office of National AIDS Policy stated that

*From the Division of Adolescent/Young Adult Medicine, Children's Hospital, Harvard Medical School, Boston, Massachusetts.*

*Address correspondence to: Elizabeth R. Woods, M.D., M.P.H., Division of Adolescent/Young Adult Medicine, Mailstop JB-335, Children's Hospital, 300 Longwood Ave., Boston, MA 02115.*

*Manuscript accepted February 1, 1998.*

25% of all new cases of HIV infection occur in young people between 13 and 20 years of age (3). Sexual exposures continue to account for the majority of adolescent AIDS cases (1). The majority of sexual risk for young men is through same-sex contact, and for young women, is heterosexual contact (1). HIV disease occurs disproportionately in black and Hispanic adolescents and young adults (1).

### *Special Issues for Adolescents*

Adolescent subpopulations are particularly difficult to engage in the care needed for the diagnosis and treatment of HIV (4). Adolescents also are one of the most medically underserved populations in the United States (5). Even when physician-adolescent encounters occur, communication about HIV risk behaviors is frequently limited (6,7). In one study, fewer than half of the adolescents with histories of "survival sex," injection drug use (IDU), same-gender sexual behavior, or a sexually transmitted disease (STD) associated with genital ulcers or sores had sought or received help for these HIV-related issues (8). STDs are synergistic with HIV in terms of transmission and offer a marker of increased health risk behaviors (9,10). Taking risks and testing limits are necessary for normal adolescent psychosocial development. For most adolescents, however, these developmental tasks are accomplished without risk to health. When risk taking and limit testing include sexual risk behaviors, the chances of STDs and HIV increase. Developmentally appropriate and culturally competent HIV counseling and testing is particularly important for youth, since a significant number of young people state that they would commit suicide if they test HIV positive (11). Gay and bisexual adolescents have high rates of depression and a threefold increased risk in suicide intent and sevenfold increase in suicide attempts (12). Use of alcohol and other recreational drugs contributes substantially to potential HIV exposure (13,14), since both increase the likelihood of teens' participating in sexual risk behaviors and reduce the likelihood of condom use (15,16).

### **Homeless and Street Youth**

Currently, there are between half a million and two million homeless and runaway teenagers in the United States (4). Many homeless youth have run away from family violence and sexual abuse or are "throw-away" youth because of issues such as sexual

orientation and pregnancy (4,17). Many of these adolescents exchange sex for food, shelter, or drugs to survive ("survival sex") (17-19). Homeless youth have particularly high seroprevalence rates (20) and require sensitive youth-oriented programs to connect them to care (21-23). Specialized training is required for health care providers to reach these populations (6,7,21,24-28).

### **Previous Efforts**

The knowledge and experience of earlier programs for HIV-positive and homeless youth were helpful in the development of the SPNS Program's adolescent models of care. The Larkin Street Youth Center in San Francisco established a central drop-in center and medical clinic that demonstrated the importance of basic needs (such as food, clothing, and shelter) as a stepping stone to health care (28). The Adolescent AIDS Program at Montefiore Medical Center in the Bronx showed the importance of outreach to referring agencies to connect youth to care (29,30). The Los Angeles system of care for homeless youth discovered the importance of collaborating with agencies in a defined geographic area solidified by frequent network functions to care for homeless and street youth (31-34). CDC-funded programs found that HIV interventions in specific communities needed to be "based on local needs" using an empirically tested theoretical framework to develop prevention activities (35). Guidelines from the Title I programs for evaluation of local HIV service delivery programs emphasize the need for both qualitative and quantitative methods to evaluate service provision, consumer opinion, and HIV-related outcome measures (36). These prior efforts demonstrated important components and evaluation mechanisms for the SPNS Program's Adolescent Initiative.

### **Theoretical Background**

Adolescents engage in clusters of HIV risk behaviors, often in association with other health-risk and problem behaviors, such as drug and alcohol use (13,14,31,32,37,38). Many of the effective health education programs aimed at preventing or changing adolescents' health-impairing behaviors have been based on social cognitive theory (39). According to social cognitive theory, interventions aimed at motivating self-directed behavior changes, thereby reducing the risk that adolescents will engage in HIV risk behaviors, should include four components: (a) in-

formation designed to increase awareness and knowledge of the consequences of behavior; (b) social and self-regulative skills development aimed at translating knowledge into preventive action; (c) opportunities for guided practice and corrective feedback in applying the skills in high-risk situations, resulting in skills enhancement and resilient self-efficacy; and (d) changes in social norms and social supports for desired behavior change (39). Many programs emphasize all four of these components through youth involvement in program design, management, outreach, and skills building risk-reduction efforts (40–42). Peer educators can help shift normative behaviors and integrate skills into the individual's values (33,43–45). Interactive skills-based education has greater impact on behavior than knowledge-based education (46,47).

### **The Need for Outreach**

Homeless adolescents are particularly difficult to treat because of their behaviors and vulnerabilities (4), the inability of providers to work with adolescents, and the legal status of adolescents in some states (48). Outreach has been shown to be effective for connecting homeless youth to services (20,22,31,49). Outreach programs are more effective when they are linked to integrated services including food, clothing, shelter, risk-reduction education, and health services (50). The CDC and others have reported that peers within the community should be recruited to deliver intervention messages and to reach less accessible populations (35,51–53), and help link hard-to-reach youth to services (54).

### *Issues of HIV Care*

Once adolescents and young adults are connected to services, a balanced approach to HIV management is essential to maintain youth in care (29,55–57). The emphasis should be on continuing to engage youth in care rather than on obtaining particular laboratory results. Working with the adolescents to evaluate how tight a control of their infection they desire and enlisting their involvement in maintaining this control are essential (55,56). As in other chronic diseases, youth adherence to complex care plans is difficult and often not a high priority to the young person (55,56). Since resistance to antiretroviral medications develops so rapidly unless regimens are carefully followed, one should be careful not to eliminate medications which might later be useful (55,58,59). Providers should work with the young person to

balance personal issues and develop a care plan. These SPNS Program models of care provide a variety of culturally sensitive projects for at-risk populations which provide youth-oriented services, linked services networks, case management, and/or comprehensive HIV health care. Integration of culturally sensitive mental health and substance use care into all health care visits is important to serve vulnerable populations (60).

### *Models of Adolescent Care*

Ten models of adolescent HIV care have been supported by the SPNS Program to demonstrate innovative concepts in youth-oriented HIV care. Those 10 models are described in the articles that follow. Several programs have developed linked services networks which cross institutional barriers. Many have demonstrated the importance of intensive case management to coordinate and maintain youth in care. Two programs are managed by young people and offer support to historically disenfranchised youth. Most of the programs contain youth advisors, providers, and educators. Outreach, HIV counseling and testing, and interagency referrals are important gateways to case identification and early intervention. Many programs link adolescents and young adults to comprehensive services, which include HIV care. Comprehensive care is more appealing to young people than specialized clinics or locations (54). Integrating ideas from youth themselves into program development improves the accessibility and appeal to young people who frequently would not obtain care.

Based on the literature reviewed above and the findings of the 10 grantees described below, it is very important to emphasize the word “adolescent” in “adolescent models of care.” These perspectives converge on the idea that programs for youth must have significant youth input in their design and continuing operations to be attractive to the clients they wish to serve. Because of their ages, young people may not have had technical training in many areas which are needed to build and sustain a program. Hence, it is essential that trusting partnerships be formed between youth and traditional professional service providers.

### *Description of 10 Models: The Consensus Model*

In the next article (pages 11–27), Huba and Melchior described a consensus model for the SPNS Program's

adolescent projects that summarize some major themes explored later in this issue. Tables are included which show the ways in which the five major consensus elements were expressed in the different projects and ways that future projects might implement these concepts. While there are several consistent themes among the programs, each model addresses the issues somewhat differently and offers unique solutions to the question of how to develop youth-friendly services that will attract large numbers of clients, keep them engaged in care, and produce useful medical and other outcomes. This synopsis is provided so that the reader can choose to read about models that are of particular interest or can compare the models as they are reviewed. In terms of their overall functional similarities, the projects can be clustered into four major groups. Those groupings are as follows:

#### I. Models stressing youth involvement

- Bay Area Young Positives (pp. 28–36)
- Health Initiatives for Youth (pp. 71–82)

#### II. Models stressing outreach to bring youth into services

- Bridgeport's Team Outreach and Primary Services (TOPS) Project (pp. 49–58)
- The University of Alabama Teenage Access Project (pp. 107–114)

#### III. Models stressing case management and linkage to services

- The Indiana Youth Access Project (pp. 83–95)
- Seattle YouthCare's Prevention, Intervention, and Education Program (pp. 96–106)
- The University of Minnesota Youth and AIDS Projects' Adolescent Early Intervention Program (pp. 115–121)

#### IV. Models stressing a comprehensive continuum of care for youth

- Boston HAPPENS Program (pp. 37–48)
- Childrens Hospital Los Angeles (pp. 59–70)
- Walden House (pp. 122–131)

### *National Evaluation*

A special feature of the grant initiative was HRSA's decision to fund a national cross-cutting evaluation center. The Measurement Group has staffed this center throughout the history of the grants and helped develop, collect, and analyze standardized

information across sites. Some of these data are presented in the articles throughout this issue for the individual sites; the cross-cutting national results are available elsewhere (61–63). Fax-in forms and questionnaires can be obtained directly from the Measurement Group (5811A Uplander Way, Culver City, CA 90230) or on its Web site (<http://www.tmg-web.com>) (64–69). The evaluation modules are also published in the SPNS Program's reports (61–63). In addition, the three SPNS reports summarize the national data from their study of the infrastructure for adolescent services in their communities and cross-cutting analyses (61–63) and can be obtained from the Measurement Group.

Special thanks to the Special Projects of National Significance Program of the Health Resources and Services Administration for the support of these projects and the national evaluation center, the editorial board members for their careful reviews and assistance to the projects, and Maria Louni and Rebecca Lamm of Boston's Children's Hospital for coordination of the supplement. Also, special thanks to Robert H. DuRant, Ph.D., Barney Singer, JD, and George Huba, Ph.D., for their comments, and to Robin Guilfooy and Michelle McCarte of Children's Hospital, Boston, for preparation of the manuscript. This publication was supported in part by the Special Projects of National Significance Program's Project BRH 970155-04-0 and Project MCJ-MA259195 from the Maternal and Child Health Bureau, Health Resources and Services Administration, Department of Health and Human Services. This publication's contents are solely the responsibility of the author and do not necessarily represent the official view of the funding agencies.

### *References*

1. Centers for Disease Control and Prevention. HIV/AIDS Surveillance Report, Mid-Year, 1997;9(1).
2. Karon JM, Rosenberg PS, McQuillan G, et al. Prevalence of HIV infection in the United States, 1984 to 1992. *JAMA* 1996;276:126–31.
3. Office of National AIDS Policy, Youth and HIV/AIDS: An American Agenda. Washington, DC: The White House, March 1996.
4. Ryan C, Futterman D. Lesbian and gay youth: Care and counseling. In: State of the Art Reviews: Adolescent Medicine, Vol. 8. Philadelphia, PA: Hanley and Belfus, 1997:259–61.
5. Blum R. Contemporary threats to adolescent health in the United States. *JAMA* 1987;257:3390–5.
6. Friedman LS, Johnson B, Brett AS. Evaluation of substance abusing adolescents by primary care physicians. *J Adolesc Health Care* 1990;11:227–30.
7. Rawitscher LA, Saitz R, Friedman LS. Adolescent's preferences regarding human immunodeficiency virus (HIV)-related physician counseling and HIV testing. *Pediatrics* 1995; 96:52–8.
8. Stiffman AR, Earls F. Behavioral risks for human immunodeficiency virus infection in adolescent medical patients. *Pediatrics* 1990;85:303–10.
9. The Contraceptive Report. In: STDs as Cofactors for HIV Infection; Taking a Sexual History. David A. Grimes, ed. Warren, NJ: Emron; 1996:VII(2).



10. Wasserheit JN. Epidemiologic synergy, interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis* 1992;19:61-7.
11. Goodman E, Cohall AT. AIDS and adolescents: knowledge, attitudes, beliefs, and behaviors in a New York City adolescent minority population. *Pediatrics* 1989;84:36-42.
12. Remafedi G, French S, Story M, et al. The relationship between suicide risk and sexual orientation: Results of a population-based study. *Am J Public Health* 1998;88:57-60.
13. Elifson KW, Boles J, Sweat M. Risk factors associated with HIV infection among male prostitutes. *Am J Public Health* 1993;83:79-83.
14. Edlin BR, Irwin KL, Farvoue S, et al. Intersecting epidemics—crack cocaine use and HIV infection among inner city young adults. *N Engl J Med* 1994;331:1422-7.
15. Friedman LS, Strunin L, Hingson R. A survey of attitudes, beliefs, behaviors and knowledge about AIDS and HIV testing by adolescents and young adults enrolled in alcohol and drug treatment. *J Adolesc Health* 1993;14:442-5.
16. Hingson RW, Strunin L, Berlin B, et al. Beliefs about AIDS, use of alcohol and drugs and unprotected sex among Massachusetts adolescents. *Am J Public Health* 1990;80:1-5.
17. Sondheim DL. HIV infection and disease among homeless adolescents. In: RJ DiClemente, ed. *Adolescents and AIDS: A Generation in Jeopardy*. Newbury Park, CA: Sage, 1992:71-85.
18. Friedman LS, Goodman E. Adolescents at risk for HIV infection: AIDS and HIV infection in office practice. *Primary Care* 1992;19:171-90.
19. Morey MA, Friedman LS. Health care needs of homeless adolescents. *Curr Opin Pediatrics* 1993;5:395-9.
20. Pfeifer RW, Oliver J. A study of HIV seroprevalence in a group of homeless youth in Hollywood, California. *J Adolesc Health* 1997;20:339-42.
21. Melchiono M, Price V. Outreach to sexual minority street population, adolescents and youth. In: *Community Street Outreach Guide*, Massachusetts Department of Public Health pub. no. 16,547-63-1000-4-91-C.R., Boston, MA: June 1991.
22. Kipke MD, Simon TR, Montgomery SB, et al. Homeless youth and their exposure to and involvement in violence while living on the streets. *J Adolesc Health* 1997;20:360-7.
23. Goulart M, Madover S. An AIDS prevention program for homeless youth. *J Adolesc Health* 1991;12:573-95.
24. Melchiono M, Lauerman JF, Palfrey J. *Reaching Out to at Risk Youth*. Boston: Harvard University Press, 1990.
25. Melchiono M, Pine V. Hustling—in substance abuse treatment: Considerations for lesbians and gay men. Boston, MA: Mobile AIDS Resource Team, 1992.
26. Sugarman ST, Hergenroeder AC, Chacko MR, Parcel GS. Acquired immunodeficiency syndrome and adolescents: Knowledge, attitudes, and behaviors of runaway and homeless youths. *Am J Dis Child* 1991;145:431-6.
27. Rotheran-Borus MJ, Koopman C. Sexual risk behaviors, AIDS knowledge, and beliefs about AIDS among runaways. *Am J Public Health* 1991;81:206-8.
28. Schor EL, Goodman E. Out of home living arrangements. In: Friedman S, Schonberg K, Fisher M, Alderman E, eds. *Comprehensive Adolescent Health Care*, 2nd ed. St. Louis, MO: Mosby-Yearbook, 1998:718-22.
29. Futterman D, Hein K, Reuben N, et al. Human immunodeficiency virus-infected adolescents: The first 50 patients in a New York City program. *Pediatrics* 1993;91:730-5.
30. DiLorenzo TA, Abramo DM, Hein K, et al. The evaluation of targeted outreach in an adolescent HIV/AIDS program. *J Adolesc Health* 1993;13:301-6.
31. Yates GL, MacKenzie RG, Pennbridge J, Swofford A. A risk profile comparison of homeless youth involved in prostitution. *J Adolesc Health* 1991;12:555-60.
32. Kipke MD, Montgomery S, MacKenzie R. Substance use among youth seen at a community based health clinic. *J Adolesc Health* 1993;14:289-94.
33. Kipke MD, Boyer C, Hein K. An evaluation of an AIDS risk reduction education and skills training (ARREST) program. *J Adolesc Health* 1993;14:533-9.
34. Kipke MD, O'Connor S, Palmer R, MacKenzie RG. Street youth in Los Angeles, profile of a group at high risk for human immunodeficiency virus infection. *Arch Pediatr Adolesc Med* 1995;149:513-9.
35. Centers for Disease Control and Prevention. Community-level prevention of human immunodeficiency virus infection among high-risk populations: the AIDS community demonstration project. *MMWR* 1996;45(Suppl):1-24.
36. Ryan White Title I Meeting Summary. Evaluation of local HIV service delivery: Issues, approaches and strategies under Title I, the Ryan White Care Act. Health Resources and Services Administration, May 1996.
37. Jessor R. Risk behavior in adolescence: A psychosocial framework for understanding and action. *J Adolesc Health* 1991;12:597-605.
38. Cohen E, MacKenzie RG, Yates GL. HEADSS, a psychosocial risk assessment instrument: implications for designing effective intervention programs for runaway youth. *J Adolesc Health* 1991;12:539-44.
39. Bandura A. A social cognitive approach to the exercise of control over AIDS infection. In: RJ DiClemente, ed. *Adolescents and AIDS: A Generation in Jeopardy*. Newbury Park, CA: Sage; 1992:89-116.
40. Jay MS, DuRant RH, Shoffitt T, et al. Effect of peer counselors on adolescent compliance in use of oral contraceptives. *Pediatrics* 1984;73:125-31.
41. Lewis CE, Lewis MA. Peer pressure and risk taking behaviors in children. *Am J Public Health* 1984;74:580-4.
42. Cohen BR, Fuccillo R. *Peer Leadership: Preventing AIDS*. Boston: The Medical Foundation, 1991.
43. Dryfoos JG. Adolescents at risk: A summation of works in the field—programs and policies. *J Adolesc Health* 1991;12:619-29.
44. Levitt MZ, Selman RL, Richmond JB. The psychosocial foundations of early adolescents' high risk behavior: Implications for research and practice. *J Res Adolesc* 1991;1:349-78.
45. Zimmerman MA, Janz NK, Wren PA. AIDS prevention in the community: lessons from the first decade. *Am Public Health Assoc* 1995 (June) report and reprinted in Atlantic Information Services, Inc., Sept. 1995;1414:1-12.
46. Anglin T. Skills-based health education for adolescents. *Adolescent Health Update* 1993;5.
47. DuRant RH, Treiber F, Getts A, et al. Comparison of two violence prevention curricula for middle school adolescents. *J Adolesc Health* 1996;19:111-7.
48. English A. Expanding access to HIV services for adolescents: Legal and ethical issues. In: RJ DiClemente, ed. *Adolescents and AIDS: A Generation in Jeopardy*. Newbury Park, CA: Sage; 1992:249-61.
49. McCormick B, Newald J. Outreach works in treating homeless youth. *Hospitals* 1986;60:162.
50. Farrow JA, Deisher RW, Brown R, et al. Position paper on the health and health needs of homeless and runaway youth. *J Adolesc Health* 1992;13:717-26.
51. Podschun GD. Teen peer outreach-street work project: HIV prevention education for runaway and homeless youth. *Public Health Rep* 1993;108:150-5.

52. Luna GC, Rotheram-Borus MJ. Street youth and the AIDS pandemic. *AIDS Educ Prev* 1992;Suppl:1-13.
53. Springer E, Mandeville K, Newell PJ, Santiago P. Peer education for street youth in Times Square, New York City. *Int Conf AIDS* 1992;8:D395.
54. Rosenfeld SL, Fox DJ, Keenan PM, et al. Findings from a focus group pilot project: Primary care experiences and preferences of urban youth. *J Pediatr Health Care* 1996;10:151-60.
55. Samples CL, Goodman E, Woods ER. Epidemiology and medical management of adolescents. In Pizzo P, Wilfert C, eds. *Pediatric AIDS: the Challenge of HIV Infection in Infants, Children and Adolescents*, 3rd ed. Baltimore, Maryland: Williams & Wilkins, 1998 (in press).
56. Samples CL. Human immunodeficiency virus in young women. In: Emans SJ, Goldstein DP, Laufer M, eds. *Pediatric and Adolescent Gynecology*, 4th ed. Philadelphia, PA: Lippincott-Raven, 1998:531-552.
57. Rogers AS, Futterman D, Levin L, D'Angelo L. A profile of human immunodeficiency virus-infected adolescents receiving health care services at selected sites in the United States. *J Adolesc Health* 1996;19:401-8.
58. Carpenter CC, Fischl MA, Hammer SM, et al. Antiretroviral therapy for HIV infection in 1997: Updated recommendations of the International AIDS Society—USA Panel. *JAMA* 1997;277:1962-9.
59. National Institutes of Health. Panel on Principles of Therapy of HIV Infection. Report of the NIH Panel to Define Principles of Therapy of HIV Infection. Rockville, MD: Department of Health and Human Services, 1997.
60. Feingold A, Slammon WR. A model integrating mental health and primary care services for a family with HIV. *Gen Hosp Psychiatry* 1993;15:290-300.
61. Huba GJ, Melchior LA, Panter AT, et al. Outreach activities in 10 adolescent-targeted projects for HIV/AIDS services: A cross-cutting evaluation of the Health Resources and Services Administration Special Projects of National Significance. Technical Monograph Series from The Measurement Group Evaluation and Dissemination Center for the HRSA SPNS Models of Adolescent HIV/AIDS Care. Culver City, CA: The Measurement Group in collaboration with the 10 adolescent SPNS Program Projects, 1996.
62. Huba GJ, Melchior LA, Panter AT, et al. Enrolled clients in 10 adolescent-targeted projects for HIV/AIDS services: A cross-cutting evaluation of the Health Resources and Services Administration Special Projects of National Significance. Technical Monograph Series from The Measurement Group Evaluation and Dissemination Center for the HRSA SPNS Models of Adolescent HIV/AIDS Care. Culver City, CA: The Measurement Group in collaboration with the 10 adolescent SPNS Program Projects, 1996.
63. Huba GJ, Melchior LA, Panter AT, et al. Interventions provided in 10 adolescent-targeted projects for HIV/AIDS services: A cross-cutting evaluation of the Health Resources and Services Administration Special Projects of National Significance. Technical Monograph Series from The Measurement Group Evaluation and Dissemination Center for the HRSA SPNS Models of Adolescent HIV/AIDS Care. Culver City, CA: The Measurement Group in collaboration with the 10 adolescent SPNS Program Projects, 1996.
64. Huba G, Melchior L. Brief Natural History Interview: Youth Version, 1994. (On-line). Available: <http://www.tmg-web.com>.
65. Huba G, Melchior L. Abbreviated Natural History Interview: Youth Version, 1994. (On-line). Available: <http://www.tmg-web.com>.
66. Huba G, Melchior L. Adolescent SPNS Contact Form, 1994. (On-line). Available: <http://www.tmg-web.com>.
67. Huba G, Melchior L. Adolescent SPNS Intervention Form, 1994. (On-line). Available: <http://www.tmg-web.com>.
68. Huba G, Melchior L. Adolescent SPNS Residential Form, 1994. (On-line). Available: <http://www.tmg-web.com>.
69. Huba G, Melchior L. Adolescent SPNS Presentation Form, 1994. (On-line). Available: <http://www.tmg-web.com>.