

## A PROGRAM FOR THE EXTRACTION OF RADIATIVE LIFETIMES FROM EXPERIMENTAL BEAM-FOIL INTENSITY DECAY DATA

D.J.G. IRWIN <sup>☆</sup> and A.E. LIVINGSTON

*Department of Physics, The University of Alberta, Edmonton, Alberta, Canada*

Received 22 November 1972

### PROGRAM SUMMARY

*Title of program:* HOMER

*Catalogue number:* AAED

*Computer:* IBM 360/67; *Installation:* University of Alberta

*Operating system:* Michigan Terminal System (MTS)

*Programming language used:* FORTRAN IV

*High speed storage required:* 16335 words

*No. of bits in a word:* 32

*Overlay structure:* None

*No. of magnetic tapes required:* None

*Other peripherals used:* Card reader, line printer, card punch

*No. of cards in combined program and test deck:* 2021

*Keywords:* Atomic, ion, life-time, excited state, beam-foil, decay curve, chi-square fit.

#### *Nature of physical problem*

The problem is to analyse experimental beam-foil intensity decay data into a sum of exponential terms, each with an amplitude and lifetime parameter. The most apt number of parameters and the optimum value of each is to be determined, together with estimates of standard deviation.

<sup>☆</sup> Present address: Department of Physics, St. Francis Xavier University-Sydney Campus, Sydney, Nova Scotia, Canada.

#### *Method of solution*

The minimum of chi-square is sought in an iterative procedure. Normal equations are obtained from a first order Taylor expansion of the multi-exponential fitting function, together with an auxiliary binomial expansion to first order. Non-convergence is impeded by two devices. A modified Marquardt method is built in as an option [1]. Initial estimates may be computed from the data. Rejection of extra-statistical data points may be allowed [2].

#### *Restrictions on the complexity of the problem*

The complexity of the fitting function is arbitrarily limited to, at most, a six-parameter fit in the form of a sum of three exponential terms. Provision is made for adjustment of the number of parameters during the computation so that a 6-, 5-, 4-, 3- or 2- parameter best fit may be obtained.

#### *Typical running time*

Typical running time is about 2 seconds on the IBM 360/67 with the G compiler for 25 data points and a six-parameter fitting function. This estimate does not include compilation of the program. It is recommended that a compiled version of the program be used in the interests of economy.

#### *References*

- [1] N.R. Draper and H. Smith, *Applied regression analysis* (Wiley, New York, 1967) p. 272.
- [2] E.M. Pugh and G.H. Winslow, *The analysis of physical measurements* (Addison-Wesley, Reading, Massachusetts, 1966) p. 108.