



Review

Contagious agalactia due to *Mycoplasma* spp. in small dairy ruminants: Epidemiology and prospects for diagnosis and control

Ángel Gómez-Martín, Joaquín Amores, Ana Paterna, Christian De la Fe*

Department of Animal Health, Faculty of Veterinary Sciences, Regional Campus of International Excellence 'Campus Mare Nostrum', Universidad de Murcia, Campus de Espinardo s/n, 30100 Murcia, Spain

ARTICLE INFO

Article history:

Accepted 21 April 2013

Keywords:

Contagious agalactia
Mycoplasma agalactiae
Small dairy ruminants
Epidemiology
Control

ABSTRACT

Contagious agalactia (CA) is a serious disease of small dairy ruminants that has a substantial economic impact on the goat and sheep milk industries. The main aetiological agent of the disease is *Mycoplasma agalactiae*, although other species, such as *Mycoplasma mycoides* subsp. *capri*, *Mycoplasma capricolum* subsp. *capricolum* and *Mycoplasma putrefaciens*, are pathogenic in goats. There are two clinical–epidemiological states of CA in sheep and goats; herds and flocks may exhibit outbreaks of CA or may be chronically infected, the latter with a high incidence of subclinical mastitis and only occasional clinical cases. The complex epidemiology of CA is related to the genetic characteristics and mechanisms of molecular variation of the *Mycoplasma* spp. involved, along with presence of CA-mycoplasmas in wild ruminant species. In goats, the situation is particularly complex and asymptomatic carriers have been detected in chronically infected herds. The coexistence of other non-pathogenic mycoplasmas in the herd further complicates the diagnosis of CA and the design of efficient strategies to control the disease. Routes of infection, such as the venereal route, may be involved in the establishment of chronic infection in herds. Current challenges include the need for improved diagnostic methods for detection of chronic and sub-clinical infections and for the design of more efficient vaccines.

© 2013 Elsevier Ltd. All rights reserved.

Introduction

Contagious agalactia (CA) is a multi-aetiological syndrome that has a serious economic impact in countries with goat and/or sheep dairy industries, mainly in the Mediterranean. Clinical disease typically involves the mammary glands, joints and eyes, although respiratory signs and reproductive disorders, including abortions, have been also reported. Clinical CA outbreaks in herds and flocks are often attributable to the entry of infected animals or to a decrease in herd immunity (Bergonier et al., 1997). In areas where CA is endemic, the most common clinical–epidemiological situation is one of chronically infected herds, with no apparent signs of disease. This situation is characterised by the presence of animals with subclinical mastitis caused by *Mycoplasma* spp., with progression to clinical mastitis in some animals. In infected goat herds, large numbers of asymptomatic carriers exist and these are often serologically negative for CA-causing *Mycoplasma* spp.

Developments in the molecular diagnosis of CA and the genetic typing of isolates are starting to clarify epidemiological aspects of the disease. These advances have improved our prospects for the

design of effective strategies to prevent, diagnose and control CA. In this paper, we review the epidemiology of CA in dairy goats and sheep, and discuss future prospects for its control.

Taxonomy of *Mycoplasma* spp. causing contagious agalactia

Contagious agalactia is caused by four different species of *Mycoplasma*: *Mycoplasma agalactiae* (Ma), the classic agent of the disease in sheep and goats, *Mycoplasma mycoides* subsp. *capri* (Mmc), *Mycoplasma capricolum* subsp. *capricolum* (Mcc) and *Mycoplasma putrefaciens* (Mp). In addition to Ma, Mmc has also been identified in sheep and, in decreasing order, Mmc, Mcc and Mp have been implicated in disease in goats (Bergonier et al., 1997; Chazel et al., 2010).

Among the four *Mycoplasma* spp. incriminated in CA, Mmc and Mcc phylogenetically belong to the *M. mycoides* cluster within the group *Spiroplasma*. The causal agent of CA, previously defined as *M. mycoides* subsp. *mycoides* large colony (LC), has been reclassified as Mmc on the basis of phylogenetic data (Fig. 1) (Manso-Silván et al., 2007 and 2009; Shahram et al., 2010). Mp also occurs in the *Spiroplasma* group and is phylogenetically close to the *M. mycoides* cluster. In contrast, Ma is phylogenetically distant from the remaining *Mycoplasma* spp. causing CA and has been ascribed to the group *Hominis* (Fig. 2). Horizontal gene transfer takes place

* Corresponding author. Tel.: +34 868887259.

E-mail address: cdelafe@um.es (C. De la Fe).




Fig. 1. Phylogenetic tree of *Mycoides* cluster, *Mycoplasma putrefaciens*, *Mycoplasma yeatsii* and *Mycoplasma cotewii* derived from distance analysis of five concatenated protein coding sequences, *fusA*, *glpQ*, *gyrB*, *lepA* and *rpoB* (Manso-Silván et al., 2007).

among the four *Mycoplasma* spp. and it is estimated that Ma shares ~18% of its genome with the *M. mycoides* cluster (Sirand-Pugnet et al., 2007).

Economic impact of contagious agalactia

In herds and flocks of domestic small ruminants, economic losses due to CA are mainly due to loss of milk production in affected animals, mortality, abortions, reduced growth rates, early culling and the costs generated by control measures. The impact of mycoplasmas on milk quality is probably underestimated (Contreras et al., 2008). The risk of exceeding the legal limits for somatic cell counts (SCC) in bulk tank milk (BTM) samples is high during an outbreak of CA and affected dairy goat and sheep herds may suffer serious mastitis problems. However, SCC cannot be used for the indirect detection of chronically infected herds of goats (Corrales et al., 2004) or flocks of sheep (Gonzalo et al., 2005), particularly in areas where the disease is endemic.

Mycoplasma spp. may also affect fertility, since Ma, Mmc and Mp exhibit tropism for the reproductive tract in sheep and goats (Gil et al., 2003; Szeredi et al., 2003; Kidanemariam et al., 2005a). In goats, there is a negative correlation between mycoplasma infection, gestation rates and the number of kids weaned (Gil et al., 2003), and it is likely that the effects of mycoplasmas on reproductive performance have been underestimated.

Antimicrobial therapy is one of the main tools used to control CA (Bergonier et al., 1997), but requires a withholding period, resulting in milk production losses. In addition, the presence of antimicrobial residues in milk poses a risk to the consumer (Allison, 1985; Dewdney et al., 1991) and could impair bacterial fermentation required for cheese production (Mourot and Loussouarn, 1981). None of four screening tests for detection of residues of 20 antimicrobial agents (including members of the aminoglycoside, macrolide, tetracycline, sulphonamide and quinolone groups) in goat's milk were able to detect European Union maximum permissible levels of tetracyclines or quinolones currently used against CA (Sierra et al., 2009), indicating a need to improve




Fig. 2. *Mycoplasma* phylogenetic tree of the 16S rRNA (Thompson et al., 2011).

the sensitivity of the methods routinely used in antimicrobial residue programmes.

Molecular epidemiology of *Mycoplasma* spp.

Multilocus sequence (MLST), variable number tandem repeats (VNTR) and pulsed field gel electrophoresis (PFGE) have been used to identify several strains of Ma circulating in Europe (McAuliffe et al., 2008, 2011). In goats, molecular studies have revealed a high genetic diversity of Ma isolates in an area endemic for Spanish CA (De la Fe et al., 2012), in contrast with a relatively low genetic diversity in Ma from sheep in France (Nouvel et al., 2012), Italy (Tola et al., 1999) and Spain (Ariza-Miguel et al., 2013). This genetic diversity of Ma in goats needs to be considered in the design of vaccines.

A single pulsotype of Mmc was detected during a CA outbreak in goats, while different pulsotypes were observed in asymptomatic auricular carriers and these in turn differed from isolates obtained from clinical cases, suggesting the existence of non-pathogenic strains of Mmc (Tardy et al., 2007). However, clinical signs developed in goats after experimental inoculation using from the clinical outbreak, as well as isolates from asymptomatic animals, indicating a lack of correlation between genomic diversity and pathogenicity (Tardy et al., 2011).

Genetic systems that allow mycoplasmas to undergo high frequency surface antigenic variations include *Vpma* in Ma and *Vmc* in Mcc (Glew et al., 2000; Wise et al., 2006). Molecular mechanisms generating high antigenic variability has been recently reviewed (Citti et al., 2010). In Ma, *Vpma* phase variation is not necessary for establishing infection, but may influence the survival and persistence of the organism (Chopra-Dewasthaly et al., 2012). In a comparison of two Ma genomes, most differences were attributable to mobile genetic elements, including integrating conjugative elements (ICE) and insertion sequences (IS), and elements related to the variable expression of surface proteins (Nouvel et al., 2010).

Genomic sequencing has revealed a Ma prophage resembling a prophage identified in *M. conjunctivae*, which has been linked to outbreaks of keratoconjunctivitis in the Alpine ibex (*Capra ibex*) (Tardy et al., 2012). These findings point to the highly dynamic nature of Ma. Several coding sequences identified in Ma are conserved in other ruminant *Mycoplasma* spp., and some of these genome fragments have been implicated in a substantial horizontal gene transfer event between Ma and other ruminant mycoplasmas, such as *M. bovis* and members of the *Mycoides* cluster (Skapski et al., 2011).

Analytical epidemiology of contagious agalactia

CA is distributed worldwide and has been reported in Africa, Asia and America (OIE, 2012). However, the Mediterranean region traditionally has been the focus of attention because of its large dairy goat industry and the endemic status of CA in the region (Bergonier et al., 1997).

Although Ma is the main agent responsible for CA in sheep, Mmc has also been identified sporadically in this species and is able to produce lesions (Trichard et al., 1993; Kidanemariam et al., 2005a; Chazel et al., 2010). In goat herds, the aetiology of CA seems to be more complex, since up to four species of *Mycoplasma* spp. have been defined as causative agents (Ma, Mcc, Mmc and Mp; OIE, 2012) and mixed infections are common at individual or herd level. This, along with the presence of other non-pathogenic mycoplasmas, makes the diagnosis and control of CA particularly difficult (Gil et al., 2003; Gómez-Martín et al., 2012a). Contact between different *Mycoplasma* spp. may also promote the transfer of genetic material (Nouvel et al., 2010).

Outbreaks of disease caused by Ma and Mmc have been detected in several wild ruminants, including the Spanish ibex (*Capra*

pyrenaica) (Verbisck et al., 2008, 2010), Alpine ibex (Chazel et al., 2010; Giangaspero et al., 2010) and chamois (Tardy et al., 2012). Pneumonia associated with Mcc has been reported in the markhor (*Capra falconeri*) (Ostrowski et al., 2011). The occurrence of these outbreaks in wild ruminants has been linked to interactions with domestic species sharing grazing lands and water troughs, movement of livestock, high population density of wild ruminants, unfavourable climate conditions and the availability of forage sources (Verbisck et al., 2008; Ostrowski et al., 2011). Further work is needed to clarify the role of wild ruminants in the epidemiology of CA in domestic sheep and goats.

The epidemiological role of asymptomatic carriers, especially auricular carriers, has been investigated extensively. All four *Mycoplasma* spp. involved in CA can be isolated from the external ear canal of goats, both in animals with clinical disease and asymptomatic animals (Bergonier et al., 1997). Carriers of Ma and Mmc have been also detected in wild goat populations (Chazel et al., 2010). Several hypotheses have been evoked to explain the localisation of *Mycoplasma* spp. in the ear: (1) spread by vectors, such as blood-feeding mites (Cottew and Yeats, 1982; DaMassa and Brooks, 1991); the presence of Mp is correlated with the presence of mites in the external ear canal (Otero et al., 2009); (2) damage to the tympanic membrane; simultaneous infection of the middle ear and external auditory canal has been reported in goats (Cottew and Yeats, 1982) and this mode of localisation appears to occur in calves infected with *M. bovis* (Walz et al., 1997); (3) via the auditory tube, which favours spread of Mmc from the respiratory system to the middle ear (Gómez-Martín et al., 2012a); Mmc is able to colonise the ear via the oronasal route following experimental infection (DaMassa and Brooks, 1991); and (4) via the bloodstream; after experimental infection in the absence of mites, the external ear canal is colonised by Ma following mycoplasmaemia (De la Fe et al., 2011). In a study of bucks that were auricular carriers of Mmc, the organism was detected in chronic lesions in the respiratory, genitourinary and nervous systems (Gómez-Martín et al., 2012a).

The host humoral immune response does not seem to be able to prevent colonisation of the ears by *Mycoplasma* spp. (Castro-Alonso et al., 2009; De la Fe et al., 2011) and a high proportion of goats with auricular colonisation by Ma and Mmc are seronegative (De la Fe et al., 2010; Gómez-Martín et al., 2012b). In endemic areas, in which CA in most herds is chronic, the presence of many asymptomatic carriers indicates that the infection is constantly perpetuated, compromising disease control and eradication measures (Thiaucourt and Bölske, 1996; Mercier et al., 2007). In these chronically infected herds, the presence of mycoplasmas may lead to outbreaks under conditions of stress and/or a diminished immune response (Thiaucourt and Bölske, 1996; De la Fe et al., 2007a). Mmc isolates associated with septicaemia cannot be distinguished on the basis of genetic or pathogenic characteristics from isolates carried in the external ear canal (Tardy et al., 2011). Further work is needed to clarify the pathogenesis and epidemiological role of CA carriers.

The possibility of venereal transmission of CA in goats has been suggested by: (1) detection of Ma and Mmc in the semen of naturally infected bucks (De la Fe et al., 2009a; Gómez-Martín et al., 2012b); (2) isolation of Mp and Mmc from the female reproductive tract (Gil et al., 2003; Szeredi et al., 2003); and (3) the detection of Ma, Mmc and Mcc in bucks in artificial insemination centres (Amores et al., 2011a). Viable Ma, Mmc and Mp have also been detected at several sites in the male reproductive tract, including the testes, bulbourethral glands and prepuce of naturally infected bucks (Gil et al., 2003; De la Fe et al., 2010; Gómez-Martín et al., 2012a, 2012b).

The potential risk of transmitting mycoplasmal infection from males to females has prompted the design of specific control measures linked to genetic improvement programmes for dairy goat breeds: (1) preventing the entry of asymptomatic auricular carriers



Fig. 3. Contagious agalactia is characterised by the classical triad of (a) joint, (b) mammary and (c) eye disease. (d) Disease is sometimes severe and can lead to emaciation and death. (e) Respiratory signs range from coughing to dyspnoea, leading to death in severe cases, especially in young animals. (f) Abortion can occur sporadically or as outbreaks.

into genetic selection centres based on the results of ear swab cultures and PCR; and (2) detecting infected animals at artificial breeding centres by routine testing of semen ejaculates by culture and PCR (Gómez-Martín et al., 2012b). Ma is shed intermittently in goat buck semen (De la Fe et al., 2009), indicating a need to continuously monitor the semen produced at these centres (Amores et al., 2011a; Gómez-Martín et al., 2012a).

Clinical disease associated with contagious agalactia

In addition to the classic triad of diseases associated with CA (mastitis, arthritis and conjunctivitis) (Fig. 3), respiratory and reproductive disorders have been observed in goats infected with CA-associated mycoplasmas (Bergonier et al., 1997). In chronically infected dairy goat herds, subclinical mastitis is mostly frequently observed, among which episodes of clinical mastitis occur sporadically. However, in these flocks, mycoplasmal infection does not have substantial effects on milk quality (Corrales et al., 2004; De la Fe et al., 2009b).

Mmc and Mp have been identified in the brain and meninges of goats with neurological disease, as well as in animals without clinical signs or lesions (DaMassa et al., 1987; Gómez-Martín et al., 2012a). Neurological signs associated with CA include opisthotonus, circling and coma (DaMassa et al., 1987; Kinde et al., 1994; Bajmocy et al., 2000).

The presence of mycoplasmas in the reproductive tract of females may be associated with reduced fertility (Gil et al., 2003; Di Provvido et al., 2009). Ulcerative balanoposthitis and vulvitis in sheep have also been associated with reduced fertility (Trichard et al., 1993; Kidanemariam et al., 2005a). However, Mmc has been detected in the genital tract of apparently healthy sheep (Kidanemariam et al., 2005a). Therefore, it remains to be determined whether mycoplasmas in the genital tract of small ruminants represent a reservoir of infection within a herd or flock, as suggested for mycoplasmas in horses (Spergser et al., 2002).

Diagnosis of contagious agalactia

The diagnosis of CA, especially in endemic zones, is complex and depends on laboratory testing (Bergonier et al., 1997). Recent studies indicate that milk samples for isolating mycoplasmas should not be frozen, since freezing reduces the viability of the organisms (Amores et al., 2010a). Therefore, it is preferable to use fresh samples or to treat samples with a bacteriostatic preservative, such as azidiol (Amores et al., 2010a). The advent of diagnostic molecular tests, especially real-time PCR has provided rapid, sensitive and robust tests for identifying CA-causing mycoplasmas (Fitzmaurice et al., 2008; Oravcová et al., 2009; Becker et al., 2012). These tests have proved valid for the systematic anal-

Table 1

Characteristics of *Mycoplasma* spp. infecting small ruminants.

Species	Growth in vitro	Typical colony size	Biochemical features ^a						Host	Disease	Samples
			S	A	U	T	F	C			
<i>M. agalactiae</i>	Good	Small	–	–	–	+	+	+	Sheep/goat	Contagious agalactia	Milk, joint fluid, ocular swabs, nasal swabs, ear swabs, semen, udder, lymph nodes, lung lesions, brain
<i>Mycoplasma mycoides</i> subsp. <i>capri</i>	Good	Large	+	–	–	+	–	–	Goat/sheep	Contagious agalactia	Milk, joint fluid, ocular swabs, nasal swabs, ear swabs, semen, udder, lymph nodes, lung lesions, brain
<i>Mycoplasma capricolum</i> subsp. <i>capricolum</i>	Good	Large	+	–	–	+	+	–	Goat	Contagious agalactia	Milk, joint fluid, ocular swabs, nasal swabs, ear swabs, udder, lymph nodes, lung lesions, brain
<i>M. putrefaciens</i>	Good	Small	+	–	–	+	+	+	Goat	Contagious agalactia	Milk, joint fluid, udder, lymph nodes, lung lesions
<i>Mycoplasma capricolum</i> subsp. <i>capripneumoniae</i>	Fastidious	Large	+	–	–	+	+	–	Goat	Caprine pleuropneumonia	Nasal, ear swabs, lung lesions, pleural fluid, mediastinal lymph nodes
<i>M. ovipneumoniae</i>	Fastidious	Centreless	+	–	–	+	–	–	Sheep/goat	Non-progressive pneumonia	Nasal swabs, bronchoalveolar lavage, lung lesions, pleural fluid, bronchopulmonary lymph nodes
<i>M. conjunctivae</i>	Fastidious	Small	+	–	–	+	–	–	Sheep	Infectious keratoconjunctivitis	Eye swabs
<i>M. arginini</i>	Good	Small	–	+	–	–	–	–	Sheep/goat	Opportunistic pathogen	Nasal swabs, bronchoalveolar lavage, lung lesions, pleural fluid, bronchopulmonary lymph nodes
<i>M. cotewii</i>	Good	Variable	+	–	–	+	–	–	Goat/sheep	Non-pathogenic	Ear swabs
<i>M. yeatsii</i>	Good	Variable	+	–	–	+	–	–	Goat/sheep	Non-pathogenic	Ear swabs, milk
<i>M. auris</i>	Good	Variable	–	+	–	+	+	–	Goat	Non-pathogenic	Ear swabs

^a S, fermentation of sugars; A, hydrolysis of arginine; U, hydrolysis of urea; T, reduction of tetrazolium; F, phosphatase activity; C, formation of films and spots in solid medium.

ysis of BTM (Chazel et al., 2010), even in the presence of preservatives, such as bronopol or sodium azide (Amores et al., 2011b).

In endemic areas, BTM testing is recommended to determine the health status of herds (Contreras et al., 2008), noting that detection of mycoplasmas in milk is affected by the number of shedding animals and the final tank volume (Amores et al., 2012). Intermittent Ma shedding in milk has also been reported in sheep BTM samples in CA endemic areas; 26.3% of farms with a positive test result for the disease returned a negative result at the next sampling (Ariza-Miguel et al., 2013). It has been proposed that samples from clinical cases of mastitis should be tested at the same time as BTM to increase diagnostic sensitivity at the herd level, since herds have been identified in which five consecutive BTM samples were negative for Ma and Mmc, whereas CA infection was confirmed on testing of samples from cases of clinical mastitis (Amores et al., 2012).

Culture combined with PCR is the best method of detecting CA in individual cases (Amores et al., 2010b; Gómez-Martín et al., 2012a) and in BTM for diagnosis at the herd level (Ariza-Miguel et al., 2012). However, non-pathogenic *Mycoplasma* spp., such as *Mycoplasma yeatsii*, *Mycoplasma cotewii* and *Mycoplasma auris*, have also been isolated from small ruminants, especially goats (Chazel et al., 2010), indicating the need to correctly identify the species of *Mycoplasma* involved (Table 1).

Serological tests are available for detection of antibodies against CA (Bergonier et al., 1997; Nicholas et al., 2008). However, some commercial ELISA kits have a poor ability to detect antibodies against some Ma strains and test performance varies according to host species and geographical origin of samples (Poumarat et al., 2012). The low sensitivity of some serological tests may explain the lack of a detectable antibody response in some infected animals (Amores et al., 2011a). Conversely, cross-reactions among *Mycoplasma* spp. will yield false positive results, sometimes corresponding to the presence of non-pathogenic mycoplasmas (Di Provvido et al., 2009; Poumarat et al., 2012). These shortcomings have limited the use of serology to assess the health status of herds or flocks (Bergonier et al., 1997; Schubert et al., 2011; Gómez-Martín et al., 2012b; Agnone et al., 2013a). The sensitivity of serological

testing may be improved by combining ELISA with Western blot analysis (Schubert et al., 2011; Agnone et al., 2013a).

Control of contagious agalactia

Current measures to control CA in affected areas have mainly been through the use of antibiotics and vaccines, with few changes over the last 15 years (Bergonier et al., 1997). These treatments are an alternative to culling infected animals of high genetic value, as well as rare domestic breeds and wild species in danger of extinction. Several studies have determined antibiotic resistance profiles of CA-causing *Mycoplasma* spp. (Loria et al., 2003; Al-Momani et al., 2006; Antunes et al., 2007a, 2007b, 2008); resistance may lead to failure of some antibiotic treatments (Gómez-Martín et al., 2013).

The antibiotics of choice against Mmc and Ma are fluoroquinolones, tetracyclines and macrolides. While erythromycin is effective against Mmc, Mcc and Mp, it is inefficient against the strains of Ma tested so far (Loria et al., 2003; Al-Momani et al., 2006; Antunes et al., 2007a, 2007b, 2008). Enrofloxacin, florfenicol, oxytetracycline and spiramycin have been used to treat ulcerative balanitis and vulvitis in sheep caused by Mmc (Kidanemariam et al., 2005b); Mmc exhibits resistance to nalidixic acid, gentamicin, streptomycin and spectinomycin. Similarly, Ma is resistant to streptomycin and nalidixic acid (Antunes et al., 2007a, 2008). The *Hominis* group shows an intrinsic resistance to erythromycin and streptomycin (Furneri et al., 2001; Königsson et al., 2002). A low sensitivity to tetracyclines has also been observed in some strains of Mp (Antunes et al., 2007b).

In a recent study, systemic marbofloxacin treatment was assessed in goat bucks that were asymptomatic auricular carriers of CA-causing mycoplasmas at an artificial breeding centre (Gómez-Martín et al., 2013). Despite the known high susceptibility of these microorganisms to fluoroquinolones and their good tissue penetration, these agents were unable to eliminate Ma and Mmc from the external ear canal of affected animals and also had detrimental effects on sperm motility. Thus, current antimicrobial strat-

egies have limited effects on carriers in chronically infected herds, with implications for controlling CA in artificial breeding centres.

The strategies employed for vaccination against CA have also varied little over the years and conventional monovalent or polyvalent killed vaccines are still used (Nicholas et al., 2008). Under field conditions, vaccines may prevent the appearance of new clinical signs and reduce mycoplasmal excretion, but are unable to prevent transmission of infection (De la Fe et al., 2007b; Agnone et al., 2013a, 2013b). The low efficacy of currently available vaccines may be due to the multi-aetiological nature of CA, the molecular characteristics of the causative species and the ability of CA-causing mycoplasmas to avoid the host immune response.

The best clinical results with inactivated vaccines have been observed when saponin has been used as an adjuvant or inactivating agent (De la Fe et al., 2007a, 2007b; Agnone et al., 2013a, 2013b). The beneficial effect of saponin in vaccines is linked to the timing of T lymphocyte memory cell activation (Agnone et al., 2013a). In an experimental study in sheep, a live attenuated vaccine against Ma also induced good clinical protection (Agnone et al., 2013b). It is likely that a new generation of recombinant vaccines in the next 10 years will provide better protection against mycoplasmal infection, especially against Ma.

Another tool proposed to eliminate a source of CA infection for offspring is pasteurisation of colostrum. Treatment of experimentally contaminated colostrum samples for 60 min at 60 °C successfully eliminates viable Mmc colonies (Paterna et al., 2012). Although treatment of colostrum for 30 min at 56 °C significantly reduces Ma counts, this *Mycoplasma* spp. is able to survive after treatment of colostrum for 120 min at 60 °C (Paterna et al., 2012).

Conclusions

A better understanding of the genetic characteristics of the mycoplasma species involved in CA and the molecular mechanisms driving their variability and pathogenicity is essential to explain the complex epidemiology of CA, especially in goat herds. There is a need to clarify the consequences of the coexistence of several mycoplasmas in the same animal or herd, the epidemiological role of asymptomatic carriers and the potential risk of transmitting mycoplasmal infection from wild animals to domestic small ruminants. More sensitive diagnostic methods should be targeted at detecting asymptomatic carriers in chronically infected herds. Finally, to achieve real control of CA, new efficient control tools are needed to prevent the infection entering a herd.

Conflict of interest statement

None of the authors of this paper has a financial or personal relationship with other people or organisations that could inappropriately influence or bias the content of the paper.

Acknowledgements

The work of our Research group is supported by the Fundación SENECA (Agencia Regional de Ciencia y Tecnología de la Región de Murcia, Project 11785/PI/09) and by Project AGL2009-09128 of the Spanish Ministry of Science and Innovation, Government of Spain. The authors thank Dr A. Contreras, Dr J.C. Corrales and Dr A. Sánchez (Research group 'Ruminant Health' Departamento de Sanidad Animal, Universidad de Murcia, Spain) for their help and for critical reading of the manuscript.

References

Agnone, A., La Manna, M.P., Loria, G.R., Puleio, R., Villari, S., Nicholas, R.A.J., Guggino, G., Sireci, G., 2013a. Timing of activation of CD4⁺ memory cells as a possible

- marker to establish the efficacy of vaccines against contagious agalactia in sheep. *Veterinary Immunology and Immunopathology* 152, 252–259.
- Agnone, A., La Manna, M., Sireci, G., Puleio, R., Usticino, A., Ozdemir, U., Nicholas, R., Chiaracane, V., Dieli, F., Di Marco, V., Loria, G., 2013b. A comparison of the efficacy of commercial and experimental vaccines for contagious agalactia in sheep. *Small Ruminant Research* 112, 230–234.
- Allison, J.R., 1985. Antibiotic residues in milk. *British Veterinary Journal* 141, 9–16.
- Al-Momani, W., Nicholas, R.A., Janakat, S., Abu-Basha, E., Ayling, R.D., 2006. The in vitro effect of six antimicrobials against *Mycoplasma putrefaciens*, *Mycoplasma mycoides* subsp. *mycoides* LC and *Mycoplasma capricolum* subsp. *capricolum* isolated from sheep and goats in Jordan. *Tropical Animal Health and Production* 38, 1–7.
- Amores, J., Sánchez, A., Gómez-Martín, A., Corrales, J.C., Contreras, A., De la Fe, C., 2010a. Viability of *Mycoplasma agalactiae* and *Mycoplasma mycoides* subsp. *capri* in goat milk samples stored under different conditions. *Veterinary Microbiology* 145, 347–350.
- Amores, J., Corrales, J.C., Gómez-Martín, A., Sánchez, A., Contreras, A., De la Fe, C., 2010b. Comparison of culture and PCR to detect *Mycoplasma agalactiae* and *Mycoplasma mycoides* subspecies *capri* in ear swabs taken from goats. *Veterinary Microbiology* 140, 105–108.
- Amores, J., Gómez-Martín, A., Corrales, J.C., Sánchez, A., Contreras, A., De la Fe, C., 2011a. Presence of contagious agalactia causing mycoplasmas in Spanish goat artificial insemination centres. *Theriogenology* 75, 1265–1270.
- Amores, J., De la Fe, C., Gómez-Martín, A., Corrales, J.C., Contreras, A., Sánchez, A., 2011b. Preserved goat milk as a valid sample for the PCR detection of *Mycoplasma agalactiae*. *Small Ruminant Research* 99, 61–64.
- Amores, J., Sánchez, A., Gómez-Martín, A., Corrales, J.C., Contreras, A., De la Fe, C., 2012. Surveillance of *Mycoplasma agalactiae* and *Mycoplasma mycoides* subsp. *capri* in dairy goat herds. *Small Ruminant Research* 102, 89–93.
- Antunes, N.T., Tavio, M.M., Assunção, P., Rosales, R.S., Aquili, V., De la Fe, C., Poveda, J.B., 2007a. In vitro susceptibilities of field isolates of *Mycoplasma mycoides* subsp. *mycoides* large colony type to 15 antimicrobials. *Veterinary Microbiology* 119, 72–75.
- Antunes, N.T., Tavio, M.M., Mercier, P., Ayling, R.D., Al-Momani, W., Assunção, P., Rosales, R.S., Poveda, J.B., 2007b. In vitro susceptibilities of *Mycoplasma putrefaciens* field isolates. *Antimicrobial Agents and Chemotherapy* 5, 3452–3454.
- Antunes, N.T., Tavio, M.M., Assunção, P., Rosales, R.S., Poveda, C., De la Fe, C., Gil, M.C., Poveda, J.B., 2008. In vitro susceptibilities of field isolates of *Mycoplasma agalactiae*. *The Veterinary Journal* 177, 436–438.
- Ariza-Miguel, J., Rodríguez-Lázaro, D., Hernández, M., 2012. A survey of *Mycoplasma agalactiae* in dairy sheep farms in Spain. *BMC Veterinary Research* 8, 171.
- Ariza-Miguel, J., Rodríguez-Lázaro, D., Hernández, M., 2013. Molecular characterization of *Mycoplasma agalactiae* reveals the presence of an endemic clone in Spain. *Journal of Clinical Microbiology* 51, 656–660.
- Bajmocy, E., Turcsanyi, I., Bolske, G., Bacsadi, A., Kiss, I., 2000. Disease caused by *Mycoplasma mycoides* subspecies *mycoides* LC in Hungarian goat herds. *Acta Veterinaria Hungarica* 48, 277–283.
- Becker, C.A., Ramos, F., Sellal, E., Moine, S., Poumarat, F., Tardy, F., 2012. Development of a multiplex real-time PCR for contagious agalactia diagnosis in small ruminants. *Journal of Microbiological Methods* 90, 73–79.
- Bergonier, D., Berthelot, X., Poumarat, F., 1997. Contagious agalactia of small ruminants: Current knowledge concerning epidemiology, diagnosis and control. *Revue Scientifique et Technique et Technique (Office International des Épizooties)* 16, 848–873.
- Castro-Alonso, A., Rodríguez, F., De la Fe, C., Espinosa de los Monteros, A., Poveda, J.B., Andrada, M., 2009. Correlating the immune response with the clinical-pathological course of persistent mastitis experimentally induced by *Mycoplasma agalactiae* in dairy goats. *Research in Veterinary Science* 86, 274–280.
- Chazel, M., Tardy, F., Le Grand, D., Calavas, D., Poumarat, F., 2010. Mycoplasmoses of ruminants in France: Recent data from the national surveillance network. *BMC Veterinary Research* 6, 32.
- Chopra-Dewasthaly, R., Baumgartner, M., Gamper, E., Innerebner, C., Zimmermann, M., Schilcher, F., Tichy, A., Winter, P., Jechlinger, W., Rosengarten, R., Spersger, J., 2012. Role of *Vpma* phase variation in *Mycoplasma agalactiae* pathogenesis. *FEMS Immunology and Medical Microbiology* 63, 307–322.
- Citti, C., Nouvel, L.X., Baranowski, E., 2010. Phase and antigenic variation in mycoplasmas. *Future Microbiology* 5, 1073–1085.
- Contreras, A., Miranda, R.E., Sánchez, A., De la Fe, C., Sierra, D., Luengo, C., Corrales, J.C., 2008. Presence of *Mycoplasma* species and somatic cell counts in bulk-tank goat milk. *Small Ruminant Research* 75, 247–251.
- Corrales, J.C., Sánchez, A., Luengo, C., Poveda, J.B., Contreras, A., 2004. Effect of clinical contagious agalactia on the bulk tank milk somatic cell count in Murciano Granadina goat herds. *Journal of Dairy Science* 87, 3165–3171.
- Cottew, G.S., Yeats, F.R., 1982. Mycoplasmas and mites in the ears of clinically normal goats. *Australian Veterinary Journal* 59, 77–81.
- DaMassa, A.J., Brooks, D.L., 1991. The external ear canal of the goats and other animals as a mycoplasma habitat. *Small Ruminant Research* 4, 85–93.
- DaMassa, A.J., Brooks, D.L., Holmberg, C.A., Moe, A.I., 1987. Caprine mycoplasmosis: An outbreak of mastitis and arthritis requiring the destruction of 700 goats. *Veterinary Record* 120, 409–413.
- De la Fe, C., Gutiérrez, A., Poveda, J.B., Assunção, P., Ramírez, A.S., Fabelo, F., 2007a. First isolation of *Mycoplasma capricolum* subsp. *capricolum*, one of the causal agents of caprine contagious agalactia, on the island of Lanzarote (Spain). *The Veterinary Journal* 173, 440–442.

- De la Fe, C., Assunção, P., Saavedra, P., Tola, S., Poveda, C., Poveda, J.B., 2007b. Field trial of two dual vaccines against *Mycoplasma agalactiae* and *Mycoplasma mycoides* subsp. *mycoides* (large colony type) in goats. *Vaccine* 8, 2340–2345.
- De la Fe, C., Amores, J., Gómez-Martín, A., Sánchez, A., Contreras, A., Corrales, J.C., 2009a. *Mycoplasma agalactiae* detected in the semen of goat bucks. *Theriogenology* 72, 1278–1281.
- De la Fe, C., Sánchez, A., Gutiérrez, A., Contreras, A., Corrales, J.C., Assunção, P., Poveda, C., Poveda, J.B., 2009b. Effects on goat milk quality of the presence of *Mycoplasma* spp. in herds without symptoms of contagious agalactia. *Journal of Dairy Research* 76, 20–23.
- De la Fe, C., Gómez-Martín, A., Amores, J., Corrales, J.C., Sánchez, A., Poveda, J.B., Contreras, A., 2010. Latent infection of male goats with *Mycoplasma agalactiae* and *Mycoplasma mycoides* subspecies *capri* at an artificial insemination centre. *The Veterinary Journal* 186, 113–115.
- De la Fe, C., Castro-Alonso, A., Herráez, P., Poveda, J.B., 2011. Recovery of *Mycoplasma agalactiae* from the ears of goats experimentally infected by the intramammary route. *The Veterinary Journal* 190, 94–97.
- De la Fe, C., Amores, J., Tardy, F., Sagne, E., Nouvel, L.X., Citti, C., 2012. Unexpected genetic diversity of *Mycoplasma agalactiae* caprine isolates from an endemic geographically restricted area of Spain. *BMC Veterinary Research* 27, 146.
- Dewdney, J.M., Maes, L., Raynaud, J.P., Blanc, F., Scheid, J.P., Jackson, T., Lens, S., Verschueren, C., 1991. Risk assessment of antibiotic residues of beta-lactams and macrolides in food products with regard to their immuno-allergic potential. *Food and Chemical Toxicology* 29, 477–483.
- Di Provvido, A., Scacchia, M., Varasano, V., Churchward, C., De Caro, C., Al-Momani, W., Di Francesco, G., Ayling, R.D., Lelli, R., Nicholas, R.A., 2009. Experimental infection of goats with an unusual strain of *Mycoplasma mycoides* subsp. *capri* isolated in Jordan. *Journal of Comparative Pathology* 141, 121–126.
- Fitzmaurice, J., Sewell, M., Manso-Silván, L., Thiaucourt, F., McDonald, W.L., O'Keefe, J.S., 2008. Real-time polymerase chain reaction assays for the detection of members of the *Mycoplasma mycoides* cluster. *New Zealand Veterinary Journal* 56, 40–47.
- Furneri, P.M., Rappazzo, G., Musumarra, M.P., Di Pietro, P., Catania, L.S., Roccasalva, L.S., 2001. Two new point mutations at A2062 associated with resistance to 16-membered macrolide antibiotics in mutant strains of *Mycoplasma hominis*. *Antimicrobial Agents and Chemotherapy* 45, 2958–2960.
- Giangaspero, M., Orusa, R., Nicholas, R.A.J., Harasawa, R., Ayling, R.D., Churchward, C.P., Whatmore, A., Bradley, D., Robetto, S., Sacchi, L., Domenis, L., 2010. Characterization of *Mycoplasma* isolated from an ibex (*Capra ibex*) suffering from keratoconjunctivitis in northern Italy. *Journal of Wildlife Diseases* 46, 1070–1078.
- Gil, M.C., Pena, F.J., Hermoso, D.M., Gómez, L., 2003. Genital lesions in an outbreak of caprine contagious agalactia caused by *Mycoplasma agalactiae* and *Mycoplasma putrefaciens*. *Journal of Veterinary Medicine B. Infectious Diseases and Veterinary Public Health* 50, 484–487.
- Glew, M.D., Papazisi, L., Poumarat, F., Bergonier, D., Rosengarten, R., Citti, C., 2000. Characterization of a multigene family undergoing high-frequency DNA rearrangements and coding for abundant variable surface proteins in *Mycoplasma agalactiae*. *Infection and Immunity* 68, 4539–4548.
- Gómez-Martín, A., De la Fe, C., Amores, J., Sánchez, A., Contreras, A., Paterna, A., Buendía, A.J., Corrales, J.C., 2012a. Anatomic location of *Mycoplasma mycoides* subsp. *capri* and *Mycoplasma agalactiae* in naturally infected goat male auricular carriers. *Veterinary Microbiology* 157, 355–362.
- Gómez-Martín, A., Corrales, J.C., Amores, J., Sánchez, A., Contreras, A., Paterna, A., De la Fe, C., 2012b. Controlling contagious agalactia in artificial insemination centers for goats and detection of *Mycoplasma mycoides* subspecies *capri* in semen. *Theriogenology* 77, 1252–1256.
- Gómez-Martín, A., Sánchez, A., Amores, J., Corrales, J.C., Contreras, A., De la Fe, C., 2013. Effect of marbofloxacin on mycoplasma carrier state and sperm quality in goat bucks. *Small Ruminant Research* 112, 186–190.
- Gonzalo, C., Carriedo, J.A., Blanco, M.A., Beneitez, E., Juarez, M.T., De La Fuente, L.F., Primitivo, F.S., 2005. Factors of variation influencing bulk-tank somatic cell count in dairy sheep. *Journal of Dairy Science* 88, 969–974.
- Kidanemariam, A., Gouws, J., van Vuuren, M., Gummow, B., 2005a. Ulcerative balanitis and vulvitis of Dorper sheep in South Africa: A study on its aetiology and clinical features. *Journal of the South African Veterinary Association* 76, 197–203.
- Kidanemariam, A., Gouws, J., Van Vuuren, M., Gummow, B., 2005b. In vitro antimicrobial susceptibility of *Mycoplasma mycoides mycoides* large colony and *Arcanobacterium pyogenes* isolated from clinical cases of ulcerative balanitis and vulvitis in Dorper sheep in South Africa. *Journal of the South African Veterinary Association* 76, 204–208.
- Kinde, H., DaMassa, A.J., Wakenell, P.S., Petty, R., 1994. *Mycoplasma* infection in a commercial goat dairy caused by *Mycoplasma agalactiae* and *Mycoplasma mycoides* subsp. *mycoides* (caprine biotype). *Journal of Veterinary Diagnostic Investigation* 6, 423–427.
- Königsson, M.H., Bölske, G., Johansson, K.E., 2002. Intraspecific variation in the 16S rRNA gene sequences of *Mycoplasma agalactiae* and *Mycoplasma bovis* strains. *Veterinary Microbiology* 85, 209–220.
- Loria, G.R., Sammartino, C., Nicholas, R.A., Ayling, R.D., 2003. In vitro susceptibilities of field isolates of *Mycoplasma agalactiae* to oxytetracycline, tylosin, enrofloxacin, spiramycin and lincomycin-spectinomycin. *Research in Veterinary Sciences* 75, 3–7.
- Manso-Silván, L., Perrier, X., Thiaucourt, F., 2007. Phylogeny of the *Mycoplasma mycoides* cluster based on analysis of five conserved protein-coding sequences and possible implications for the taxonomy of the group. *International Journal of Systematic and Evolutionary Microbiology* 57, 2247–2258.
- Manso-Silván, L., Vilei, E.M., Sachse, K., Djordjevic, S.P., Thiaucourt, F., Frey, J., 2009. *Mycoplasma leachii* sp. nov. as a new species designation for *Mycoplasma* sp. bovine group 7 of Leach, and reclassification of *Mycoplasma mycoides* subsp. *mycoides* LC as a serovar of *Mycoplasma mycoides* subsp. *capri*. *International Journal of Systematic and Evolutionary Microbiology* 59, 1353–1358.
- McAuliffe, L., Churchward, C., Lawes, J.R., Loria, G., Ayling, R.D., Nicholas, R., 2008. VNTR analysis reveals unexpected genetic diversity within *M. agalactiae*, the main causative agent of contagious agalactia. *BMC Microbiology* 8, 193.
- McAuliffe, L., Gosney, F., Hlusek, M., de Garnica, M.L., Sperser, J., Kargl, M., Rosengarten, R., Ayling, R.D., Nicholas, R.A., Ellis, R.J., 2011. Multilocus sequence typing of *Mycoplasma agalactiae*. *Journal of Medical Microbiology* 6, 803–811.
- Mercier, P., Pellet, M., Morignat, E., Calavas, D., Poumarat, F., 2007. Prevalence of mycoplasmas in external ear canal of goats: Influence of the sanitary status of the herd. *Small Ruminant Research* 73, 296–299.
- Mourot, D., Loussouarn, S., 1981. Sensibilité des ferments lactiques aux antibiotiques utilisés en médecine vétérinaire. *Recueil de Médecine Vétérinaire* 157, 175–177.
- Nicholas, R., Ayling, R., McAuliffe, L., 2008. *Mycoplasma Diseases of Ruminants*. CAB International, Oxfordshire, UK, 239 pp.
- Nouvel, L.X., Sirand-Pugnet, P., Marena, M., Sagné, E., Barbe, V., Mangelot, S., Schenowitz, C., Jacob, D., Barré, A., Claverol, S., Blanchard, A., Citti, C., 2010. Comparative genomic and proteomic analyses of two *Mycoplasma agalactiae* strains: Clues to the macro- and micro-events that are shaping mycoplasma diversity. *BMC Genomics* 11, 86.
- Nouvel, L.X., Marena, M.S., Glew, M.D., Sagné, E., Giammarinaro, P., Tardy, F., Poumarat, F., Rosengarten, R., Citti, C., 2012. Molecular typing of *Mycoplasma agalactiae*: Tracing European-wide genetic diversity and an endemic clonal population. *Comparative Immunology, Microbiology and Infectious Diseases* 35, 487–496.
- OIE, 2012. WAHID Interface. World Animal Health Database. <http://www.oie.int> (accessed 14 September 2012).
- Oravcová, K., López-Enríquez, L., Rodríguez-Lázaro, D., Hernández, M., 2009. *Mycoplasma agalactiae* p40 Gene, a novel marker for diagnosis of contagious agalactia in sheep by real-time PCR: Assessment of analytical performance and in-house validation using naturally contaminated milk samples. *Journal of Clinical Microbiology* 47, 445–450.
- Ostrowski, S., Thiaucourt, F., Amirbekov, M., Mahmadsheev, A., Manso-Silván, L., Dupuy, V., Vahobov, D., Ziyoev, O., Michel, S., 2011. Fatal outbreak of *Mycoplasma capricolum* pneumonia in endangered markhors. *Emerging Infectious Diseases* 17, 2338–2341.
- Otero, N.J., Jaramillo Meza, L., Miranda Morales, R.E., Navarro Hernández, J.A., Quintero Martínez, M.T., 2009. Association of *Raillietia caprae* with the presence of *Mycoplasmas* in the external ear canal of goats. *Preventive Veterinary Medicine* 92, 150–153.
- Paterna, A., Sánchez, A., Amores, J., Gómez-Martín, A., Corrales, J.C., Contreras, A., De la Fe, C., 2012. Survival of *Mycoplasma agalactiae* and *Mycoplasma mycoides* subspecies *capri* in heat treated goat colostrum. *The Veterinary Journal* 196, 263–265.
- Poumarat, F., Le Grand, D., Gaurivaud, P., Gay, E., Chazel, M., Game, Y., Bergonier, D., 2012. Comparative assessment of two commonly used commercial ELISA tests for the serological diagnosis of contagious agalactia of small ruminants caused by *Mycoplasma agalactiae*. *BMC Veterinary Research* 8, 109.
- Schubert, E., Sachse, K., Jores, J., Heller, M., 2011. Serological testing of cattle experimentally infected with *Mycoplasma mycoides* subsp. *mycoides*. Small colony using four different tests reveals a variety of seroconversion patterns. *BMC Veterinary Research* 7, 72.
- Shahram, M., Nicholas, R.A., Wood, A.P., Kelly, D.P., 2010. Further evidence to justify reassignment of *Mycoplasma mycoides* subspecies *mycoides* large colony type to *Mycoplasma mycoides* subspecies *capri*. *Systematic and Applied Microbiology* 33, 20–24.
- Sierra, D., Contreras, A., Sánchez, A., Luengo, C., Corrales, J.C., Morales, C.T., De la Fe, C., Guirao, I., Gonzalo, C., 2009. Detection limits of non-beta-lactam antibiotics in goat's milk by microbiological residues screening tests. *Journal of Dairy Science* 92, 4200–4206.
- Sirand-Pugnet, P., Lartigue, C., Marena, M., Jacob, D., Barré, A., Barbe, V., Schenowitz, C., Mangelot, S., Couloux, A., Segurens, B., Daruvar, A., Blanchard, A., Citti, C., 2007. Being pathogenic, plastic, and sexual while living with a nearly minimal bacterial genome. *PLOS Genetics* 3, 744–758.
- Skapski, A., Hygonenq, M.C., Sagné, E., Guirao, I., Citti, C., Baranowski, E., 2011. Genome-scale analysis of *Mycoplasma agalactiae* loci involved in interaction with host cells. *PLoS One* 6, e25291.
- Sperser, J., Aurich, C., Aurich, J.E., Rosengarten, R., 2002. High prevalence of mycoplasmas in the genital tract of asymptomatic stallions in Austria. *Veterinary Microbiology* 87, 119–129.
- Szeredi, L., Tenk, M., Dan, A., 2003. Infection of two goatherds with *Mycoplasma mycoides* subsp. *capri* in Hungary, evidence of a possible faecal excretion. *Journal of Veterinary Medicine B, Infectious Diseases and Veterinary Public Health* 50, 172–177.
- Tardy, F., Mercier, P., Solsona, M., Saras, E., Poumarat, F., 2007. *Mycoplasma mycoides* subsp. *mycoides* biotype large colony isolates from healthy and diseased goats: Prevalence and typing. *Veterinary Microbiology* 121, 268–277.
- Tardy, F., Maigre, L., Tricot, A., Poumarat, F., Nguyen, L., Le Grand, D., 2011. Comparison of isolates of *Mycoplasma mycoides* subspecies *capri* from

- asymptomatic and septicaemic goats. *Journal of Comparative Pathology* 144, 70–77.
- Tardy, F., Baranowski, E., Nouvell, L.X., Mick, V., Manso-Silván, L., Thiaucourt, F., Thébault, P., Breton, M., Sirand-Pugnet, P., Blanchard, A., Garnier, A., Gibert, P., Game, Y., Poumarat, F., Citti, C., 2012. Emergence of atypical *Mycoplasma agalactiae* strains harboring a new prophage and associated with an alpine wild ungulate mortality episode. *Applied and Environmental Microbiology* 78, 4659–4668.
- Thiaucourt, F., Bölske, G., 1996. Contagious caprine pleuropneumonia and other pulmonary mycoplasmoses of sheep and goats. *Revue Scientifique et Technique (Office International des Épizooties)* 15, 1397–1414.
- Thompson, C.C., Vieira, N.M., Vicente, A.C.P., Thompson, F.L., 2011. Towards a genome based taxonomy of mycoplasmas. *Infection, Genetics and Evolution* 11, 1798–1804.
- Tola, S., Manunta, D., Rocca, S., Rocchigiani, A.M., Idini, G., Angioi, P.P., Leori, G., 1999. Experimental vaccination against *Mycoplasma agalactiae* using different inactivated vaccines. *Vaccine* 16, 2764–2768.
- Trichard, C.J., Jordaan, P., Prozesky, L., Jacobsz, E.P., Henton, M.M., 1993. The identification of *Mycoplasma mycoides mycoides* LC as the aetiological agent of balanoposthitis and vulvovaginitis in sheep in South Africa. *Onderstepoort Journal of Veterinary Research* 60, 29–37.
- Verbisck, G., González Candela, M., Galián, J., Cubero Pablo, M.J., Martín Atance, P., León Vizcaíno, L., 2008. Epidemiology of *Mycoplasma agalactiae* infection in free-ranging Spanish ibex (*Capra pyrenaica*) in Andalusia, Southern Spain. *Journal of Wildlife Disease* 44, 369–380.
- Verbisck, G., González-Candela, M., Cubero, M.J., León, L., Serrano, E., Perales, A., 2010. *Mycoplasma agalactiae* in Iberian ibex (*Capra pyrenaica*) in Spain. *Veterinary Record* 167, 425–426.
- Walz, P.H., Mullaney, T.P., Render, J.A., Walker, R.D., Mosser, T., Baker, J.C., 1997. Otitis media in preweaned Holstein dairy calves in Michigan due to *Mycoplasma bovis*. *Journal of Veterinary Diagnostic Investigation* 9, 250–254.
- Wise, K.S., Foecking, M.F., Röske, K., Lee, Y.J., Lee, Y.M., Madan, A., Calcutt, M.J., 2006. Distinctive repertoire of contingency genes conferring mutation-based phase variation and combinatorial expression of surface lipoproteins in *Mycoplasma capricolum* subsp. *capricolum* of the *Mycoplasma mycoides* phylogenetic cluster. *Journal of Bacteriology* 188, 4926–4941, Erratum in: *Journal of Bacteriology* 2006, 188, 6716–6717.