

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7888236>

# Biosynthesis of Covalently Bound Flavin: Isolation and in Vitro Flavinylation of the Monomeric Sarcosine Oxidase Apoprotein †

ARTICLE *in* BIOCHEMISTRY · JUNE 2005

Impact Factor: 3.02 · DOI: 10.1021/bi047271x · Source: PubMed

---

CITATIONS

33

---

READS

29

4 AUTHORS, INCLUDING:



Marilyn Schuman Jorns

Drexel University College of Medicine

101 PUBLICATIONS 3,051 CITATIONS

SEE PROFILE

Published in final edited form as:

Biochemistry. 2005 May 3; 44(17): 6452–6462.

## Biosynthesis of Covalently Bound Flavin: Isolation and *In Vitro* Flavinylation of Monomeric Sarcosine Oxidase Apoprotein

Alshaimaa Hassan-Abdallah, Robert C. Bruckner, Guohua Zhao, and Marilyn Schuman Jorns\*

Department of Biochemistry and Molecular Biology, Drexel University, College of Medicine, Philadelphia, PA 19102

### Abstract

The covalently bound FAD in native monomeric sarcosine oxidase (MSOX) is attached to the protein by a thioether bond between the 8 $\alpha$ -methyl group of the flavin and Cys315. Large amounts of soluble apoenzyme are produced by controlled expression in a riboflavin-dependent *E. coli* strain. A time-dependent increase in catalytic activity is observed upon incubation of apoMSOX with FAD, accompanied by the covalent incorporation of FAD to about 80% of the level observed with native enzyme. The spectral and catalytic properties of the reconstituted enzyme are otherwise indistinguishable from native MSOX. The reconstitution reaction exhibits apparent second order kinetics ( $k = 139 \text{ M}^{-1} \text{ min}^{-1}$ , 23 °C) and is accompanied by the formation of a stoichiometric amount of hydrogen peroxide. A time-dependent reduction of FAD is observed when the reconstitution reaction is conducted under anaerobic conditions. The results provide definitive evidence for autoflavinylation in a reaction that proceeds via a reduced flavin intermediate and requires *only* apoMSOX and FAD. Flavinylation of apoMSOX is not observed with 5-deazaFAD or 1-deazaFAD, an outcome attributed to a decrease in the acidity of the 8 $\alpha$ -methyl group protons. Covalent flavin attachment is observed with 8-nor-8-chloroFAD in an aromatic nucleophilic displacement reaction that proceeds via a quinonoid intermediate but not a reduced flavin intermediate. The reconstituted enzyme contains a modified cysteine-flavin linkage (8-nor-8-S-cysteinyl) as compared with native MSOX (8 $\alpha$ -S-cysteinyl), a difference that may account for its ~10-fold lower catalytic activity.

Although flavins often function as noncovalently bound prosthetic groups, a growing number of flavoenzymes have been found to contain covalently bound flavin. This diverse group of enzymes perform a remarkable range of reactions, including ion pumping, antibiotic synthesis and metabolism of biogenic amines (1–3). Monomeric sarcosine oxidase (MSOX<sup>1</sup>) is a prototypical member of a recently discovered family of amine-oxidizing enzymes that all contain covalently bound flavin (4–7). MSOX is an inducible bacterial enzyme that plays an important role in the catabolism of sarcosine (N-methylglycine), a common soil metabolite (8). The enzyme is widely used in the clinical evaluation of renal function (9). MSOX catalyzes the oxygen-dependent demethylation of sarcosine to yield glycine, formaldehyde and hydrogen peroxide. High resolution crystal structures are available for free MSOX and complexes of the enzyme with competitive inhibitors (10, 11). MSOX contains 1 mol of FAD, attached to the protein via a thioether linkage between the 8 $\alpha$ -methyl group of the isoalloxazine ring and Cys315 (8 $\alpha$ -S-cysteinyl-FAD) (12) (see Table 1 for structure). The same covalent linkage is found for FAD in monoamine oxidase A and B, mammalian enzymes that exhibit 20% sequence identity with MSOX. In other enzymes, tyrosine or histidine may replace cysteine as

\*To whom requests for reprints should be addressed. Phone: (215) 762-7495 FAX: (215) 762-4452, E-mail: marilyn.jorns@drexelmed.edu

†This work was supported in part by Grant GM 31704 (M. S. J.) from the National Institutes of Health.

<sup>1</sup>Abbreviations: MSOX, monomeric sarcosine oxidase; FAD, flavin adenine dinucleotide; IPTG, isopropyl  $\beta$ -D-thiogalactopyranoside; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis

the site of flavin attachment but the covalent linkage nearly always involves the 8 $\alpha$ -methyl group of the flavin (1).

Covalent flavin attachment is thought to occur in a reaction catalyzed by the apoprotein itself (1, 13–15). A molecular mechanism involving iminoquinone methide and reduced flavin intermediates (Scheme 1, intermediates I and II, respectively) has been proposed for MSOX and other enzymes (1, 10) but *in vitro* studies to characterize the reaction have been limited by difficulties in isolating reconstitutable apoenzymes. A notable exception is provided by *p*-cresol methylhydroxylase, a hemoflavoenzyme ( $\alpha_2\beta_2$ ) that contains separate flavin- and heme-binding subunits. In this perhaps unusual example, both subunits are required for covalent flavinylation (8 $\alpha$ -O-tyrosyl-FAD). As a consequence, the reaction can be blocked by expressing the flavin-binding subunit by itself and then triggered by mixing the separately isolated subunits in the presence of FAD (13). Expression of apo-monoamine oxidase A and B was achieved by using a riboflavin-dependent strain of *Saccharomyces cerevisiae*. Isolation of the mitochondrial membrane-bound proteins was, however, thwarted by the extreme instability of the solubilized apoenzymes (16, 17). An unstable apoenzyme form of 6-hydroxy-D-nicotine oxidase was isolated in low yield by expressing the recombinant enzyme in the presence of diphenylene iodonium (14). Analytical amounts of apo-succinate dehydrogenase have been prepared by using a reticulocyte lysate transcription-translation system (15). Flavinylation of both of these apoenzymes (8 $\alpha$ -N(3)-histidyl-FAD) was markedly accelerated by small molecule effectors.

MSOX is a relatively small (44 kDa), stable, cytoplasmic protein containing a single prosthetic group. The recombinant enzyme is highly overexpressed in *E. coli* and readily isolated in gram quantities. These features suggested that MSOX might provide a viable system for *in vitro* flavinylation studies. In this paper we describe a system for expression and preparative scale (40 mg/liter of culture) isolation of a reasonably stable apoprotein that can be reconstituted with FAD and certain FAD derivatives.

## EXPERIMENTAL PROCEDURES

### Materials

Amplex Red was obtained from Molecular Probes. NADH peroxidase, horseradish peroxidase and *o*-dianisidine were purchased from Sigma. 8-Nor-8-chlororiboflavin and 1-deazariboflavin were generous gifts from Drs. John Lambooy and Wallace Ashton, respectively. 5-Deazariboflavin was previously synthesized in this laboratory according to the method of O'Brien et al. (18). Recombinant FAD synthetase from *Corynebacterium ammoniagenes* was kindly provided by Dr. Bill McIntire. *Pfu* DNA polymerase was obtained from Stratagene. *Nde*I and *Xho*I restriction enzymes were purchased from New England Biolabs. QIAquick PCR Purification and QIAquick Gel Extraction Kits were obtained from Qiagen. Plasmid pET24a was supplied by Novagen. T4 DNA ligase was purchased from New England Biolabs. The Co<sup>2++</sup> affinity matrix was obtained from BD Bioscience. Sep-Pak C18 Cartridges were purchased from Waters.

### Expression and Purification of MSOX Holoenzyme

MSOX was expressed in *E. coli* and purified as previously described (12) except that the cells (*E. coli* DH1/pMAW) were grown in Terrific Broth (19) instead of LB. Enzyme activity was measured using the Nash assay (12) or a horseradish peroxidase assay with *o*-dianisidine as the chromogenic substrate (20). Protein was determined using the Bio-Rad micro protein assay (12). Enzyme concentration was also estimated based on flavin absorbance at 454 nm ( $\epsilon_{454} = 12,200 \text{ M}^{-1} \text{ cm}^{-1}$ ) (12). The isolated enzyme exhibited catalytic and spectral properties similar

to that reported for enzyme isolated from cells grown on LB (12) (see Table 2). However, the yield of purified enzyme (3.97 g from 10 L of cell culture) was 10-fold higher.

### Generation of a Riboflavin-dependent Derivative of *E. coli* BL21 (DE3)

Chemical mutagenesis of *E. coli* BL21 (DE3) was performed by exposing the cells to ethyl methane sulfonate for 15 min, according to the procedure described by Miller (21). The mutagenized cells were enriched for riboflavin auxotrophs using the ampicillin enrichment method described by Molholt (22). Five riboflavin-dependent colonies [*E. coli* BL21 (DE3) M1a-M1e] were isolated based on their ability to grow on LB only when the medium was supplemented with riboflavin (100 µg/mL). *E. coli* BL21 (DE3) M1b cells were selected for expression of the MSOX apoenzyme, as described below.

### Construction of Plasmid pGZ26

Plasmid pMAW contains the cloned gene for MSOX from *Bacillus sp. B-0618* (12). PCR was used to introduce an *NdeI* restriction site at the 5' end of the gene, delete the stop codon and introduce a *XhoI* restriction site at the 3' end of the gene. Reactions were performed using plasmid pMAW as template, 5'-AGGAATCATATGAGCACACATTTTGATG-3' as forward primer (*NdeI* site is underlined) and 5'-

AGACTACTCGAGGATAGTTGTTTTTGTAAACG-3' as backward primer (*XhoI* site is underlined). PCR was conducted using *Pfu* DNA polymerase and a Hybaid Touchdown Thermocycler with the following cycle settings: 30 cycles of 94°C for 40 s, 52 °C for 40 s, and 72°C for 2 min, followed by a single cycle of 72°C for 30 min. The PCR product was purified using the QIAquick PCR Purification Kit, cut with *NdeI* and *XhoI*, purified by agarose gel (1.5%) electrophoresis and recovered using the QIAquick Gel Extraction Kit. The desired *NdeI-XhoI* fragment (containing the MSOX gene minus its stop codon) was then subcloned between the *NdeI* and *XhoI* sites of plasmid pET24a using T4 DNA ligase. The resulting construct (plasmid pGZ26) places MSOX expression under the control of the T7 promoter and adds 8 amino acids [LeuGlu(His)<sub>6</sub>] to the 3'-end of the recombinant protein. Plasmid pGZ26 was sequenced across the entire MSOX insert. Sequencing was conducted by Nucleic Acid/Protein Research Core Facility at the Children's Hospital of Philadelphia. Plasmid pGZ26 was used to transform *E. coli* BL21 (DE3) M1b cells to kanamycin resistance.

### Expression and Purification of MSOX Apoenzyme

*E. coli* BL21 (DE3) M1b/pGZ26 cells were grown at 37 °C in LB containing riboflavin (4 µg/mL) and kanamycin (100 µg/mL) until A<sub>595</sub> = 0.9. The cells then were collected by centrifugation and washed twice with riboflavin-free LB. The washed cells were resuspended in riboflavin-free LB at 15 °C. MSOX expression was induced with 1 mM IPTG. Cells were harvested after 3 days at 15 °C.

All steps of the purification were conducted at 4 °C. Cells were disrupted by sonication in the presence of protease inhibitors, as described for the isolation of MSOX holoenzyme (12). The pellet obtained after centrifugation of the cell lysate was discarded. The supernatant was dialyzed against 50 mM potassium phosphate buffer, pH 7.5, containing 20% glycerol and 500 mM NaCl and then mixed with 15 ml of Co<sup>2++</sup> affinity matrix. After gentle rocking for 45 min the mixture was poured into a column. The column was washed with dialysis buffer until the eluate exhibited no detectable absorbance at 280 nm. ApoMSOX was then eluted using a 1-liter linear gradient from 0 to 150 mM imidazole in dialysis buffer. The enzyme was dialyzed against 50 mM potassium phosphate buffer, pH 7.5, containing 50 mM KCl, concentrated using a Centricon YM-30 ultrafiltration cell and stored in aliquots at -80 °C. The yield from 3 liters of culture was 123 mg. Activity and protein were measured as described for the holoenzyme. The concentration of apoenzyme was also estimated based on its absorbance at 280 nm ( $\epsilon_{280} = 36,270 \text{ M}^{-1} \text{ cm}^{-1}$ ) (12).

### Preparation of FAD Analogs

The amount of FAD synthetase required to achieved complete conversion of each riboflavin derivative to the FAD level (3.6 to 7.1  $\mu\text{g/mL}$ ) was determined in preliminary analytical trials under the conditions described below. Reaction progress was monitored by thin-layer chromatography on silica gel plates using *n*-butanol/acetic acid/water (12:3:5). Riboflavin derivatives (200  $\mu\text{M}$ ) were incubated for 18 h at 37 °C with FAD synthetase in 50 mM potassium phosphate buffer, pH 7.6, containing 5 mM ATP and 20 mM magnesium chloride. Preparative scale (200 mL) reaction mixtures were then filtered through an Amicon PM-10 membrane to remove protein. Except as noted below, the filtrate was applied to a 35-mL Sep-Pak C18 Cartridge, previously activated by washing with methanol and then equilibrated with water. The cartridge was washed with water. Elution conditions varied somewhat, depending on the FAD analog. 5-DeazaFAD was eluted with 10% acetonitrile in water. 1-DeazaFAD was not eluted by a 1-liter linear gradient from 0 to 5% acetonitrile in water. The compound began to slowly leach off the cartridge upon further washing with 5% acetonitrile. 1-DeazaFAD was eluted by switching to 10% acetonitrile in water. The Sep-Pak eluates were concentrated by lyophilization and stored at -20 °C. In the case of 8-nor-8-chloroFAD, the filtrate was applied to a 1-liter DEAE-cellulose column previously equilibrated with 5 mM potassium phosphate buffer, pH 6.8. The column was washed successively with 5 mM and 10 mM potassium phosphate buffers, pH 6.8. 8-Nor-8-chloroFAD was eluted with a 12-liter linear gradient from 20 mM to 75 mM potassium phosphate buffer, pH 6.8. The eluate was applied to a 35-mL Sep-Pak C18 Cartridge, previously treated as described above. The cartridge was washed with water. 8-Nor-8-chloroFAD was eluted with methanol, rotoevaporated to dryness, dissolved in water and stored at -20 °C

The concentration of stock solutions of the flavin analogs was estimated using the following extinction coefficients: 8-nor-8-chloroFAD,  $\epsilon_{448} = 10,600 \text{ M}^{-1} \text{ cm}^{-1}$  (23); 5-deazaFAD,  $\epsilon_{399} = 11,500 \text{ M}^{-1} \text{ cm}^{-1}$  (24); 1-deazaFAD,  $\epsilon_{540} = 6,800 \text{ M}^{-1} \text{ cm}^{-1}$  (25).

### Reconstitution of ApoMSOX with FAD or FAD Analogs

Reactions were conducted at 23 °C in 100 mM potassium phosphate buffer, pH 8.0, containing apoMSOX and excess flavin, as indicated in the text. Reactions were initiated by addition of apoMSOX. The kinetics of the reconstitution reaction with FAD were monitored by withdrawing small aliquots at various times to measure sarcosine oxidase activity using a horseradish peroxidase-coupled assay with *o*-dianisidine as the chromogenic substrate (20). Reaction progress was also monitored by measuring hydrogen peroxide formation during reconstitution of apoMSOX with FAD or 8-nor-8-chloroFAD. These reactions were conducted as described above except the reconstitution mixtures also contained horseradish peroxidase (18 U/ml), Amplex Red (80  $\mu\text{M}$ ) and dimethylsulfoxide (0.4 %). Hydrogen peroxide formation was monitored at 563 nm ( $\Delta\epsilon_{563} = 52,200 \text{ M}^{-1} \text{ cm}^{-1}$ ). The value for  $\Delta\epsilon_{563}$  was determined in studies with known amounts of hydrogen peroxide. Hydrogen peroxide solutions for these studies were standardized using an NADH peroxidase-coupled assay (26). Data obtained for reconstitution reactions were corrected for a small linear reagent blank ( $\Delta A_{563}/\text{min} = 3.2 \times 10^{-7}$ ), observed in the presence of all components except apoMSOX. Owing to the large spectral change observed upon reaction of apoMSOX with 8-nor-8-chloroFAD, the kinetics of covalent attachment could also be directly monitored at 499 nm, as described in the text.

### Isolation and Characterization of Reconstituted Enzyme

Unbound flavin was removed by ultrafiltration using a Centricon YM-30. The absorption spectrum of reconstituted enzyme was recorded after centrifugation. Extinction coefficients for covalently bound flavins were determined based on absorbance changes observed after denaturation of the reconstituted enzyme with 3 M guanidine hydrochloride. Calculations with FAD-reconstituted enzyme were made using the extinction coefficient reported for free FAD

in the presence of guanidine hydrochloride ( $\epsilon_{450} = 11,900 \text{ M}^{-1} \text{ cm}^{-1}$ ) (12). The extinction coefficient reported for free 8-nor-8-S-(N-acetylcysteinyl)-FAD ( $\epsilon_{480} = 25,200 \text{ M}^{-1} \text{ cm}^{-1}$ ) (23) was used to calculate the extinction coefficient of 8-nor-8-chloroFAD-reconstituted enzyme. To determine the stoichiometry of flavin incorporation, protein concentration was estimated based on the absorbance of the denatured protein at 280 nm ( $\epsilon_{280} = 36,270 \text{ M}^{-1} \text{ cm}^{-1}$ ) (12). The absorbance at 280 nm was corrected for the contribution from free FAD ( $\epsilon_{280} = 22,900 \text{ M}^{-1} \text{ cm}^{-1}$ ) (12) or free 8-nor-8-S-(N-acetylcysteinyl)-FAD ( $\epsilon_{280} = 18,900 \text{ M}^{-1} \text{ cm}^{-1}$ ) (27).

### Spectroscopy and Data Analysis

Absorption spectra were recorded using an Agilent Technologies 8453 diode array spectrophotometer or a Perkin-Elmer Lambda 2S spectrometer. Horseradish peroxidase assays using Amplex Red were conducted using only the Perkin-Elmer Lambda 2S spectrometer because the bright light source in the diode array spectrophotometer caused significant photodecomposition of the chromogenic substrate. For anaerobic reconstitution experiments, a solution of apoMSOX in 100 mM potassium phosphate buffer, pH 8.0, containing 50 mM glucose, was placed in the main compartment of a special cuvette with two side arms. The first and second side arms contained small aliquots of glucose oxidase and FAD or 8-nor-8-chloroFAD, respectively. The cuvette was made anaerobic by multiple cycles of evacuation, followed by filling with oxygen-free argon gas. Identical spectra were observed before and after tipping glucose oxidase (5 U/mL). Reconstitution was initiated by mixing apoMSOX (276  $\mu\text{M}$ ) with FAD or 8-nor-8-chloroFAD (120  $\mu\text{M}$ ). All spectra are corrected for dilution. Kinetic data were fitted using the curve fit function in Sigma Plot (SPSS Inc.).

## RESULTS

### Expression of ApoMSOX

A riboflavin-dependent derivative of *E. coli* BL21 (DE3) was generated by chemical mutagenesis using ethyl methane sulfonate (21), followed by enrichment for riboflavin auxotrophs as described by Molholt (22). This strain [BL21 (DE3) M1b], does not grow in LB unless the medium is supplemented with riboflavin. We reasoned that apoMSOX might be produced if riboflavin was removed from the growth medium prior to induction of enzyme expression. Native MSOX is typically expressed at 37 °C under the control of the leaky *lac* promoter and purified by a procedure involving several chromatography steps. Huge amounts of pure enzyme are obtained, especially when the cell density is increased by using Terrific broth in place of LB (400 versus 40 mg MSOX/liter of culture). Two changes were, however, introduced in order to facilitate apoenzyme expression and isolation: 1) MSOX expression was placed under the control of the T7 promoter to achieve tightly regulated expression; 2) a carboxyl-terminal affinity tag, (His)<sub>6</sub>, was incorporated to streamline protein purification.

The riboflavin-dependent strain was transformed with the new expression plasmid (pGZ26) and grown at 37 °C in riboflavin-supplemented LB until a desired cell density was attained ( $A_{595} = 0.9$ ). The cells were then transferred to unsupplemented LB and MSOX expression was induced at 15 °C. Extracts from cells grown under these conditions exhibit a prominent 44 kDa protein band that co-migrates with native MSOX on SDS-PAGE gels (Figure 1, lanes 2 and 1, respectively). Nearly all of the protein in the whole cell extract was present in the soluble fraction, as judged by SDS-PAGE analysis of supernatant and pellet fractions (data not shown). In contrast, apoenzyme expression at 37 °C generated inclusion bodies that resisted various renaturation attempts.



### Isolation of a Reconstitutable Apoenzyme

The soluble 44 kDa protein produced at 15 °C was isolated by using a  $\text{Co}^{++}$  affinity matrix. This single step procedure yields pure protein (Figure 1, lanes 3 and 4) in amounts (40 mg/liter of culture) similar to that observed for the isolation of native enzyme from cells grown in LB. The preparation was predominantly apoMSOX, as judged by the low catalytic activity and minimal absorbance in the visible region, as compared with native enzyme (Figure 2, Table 2). The residual FAD in the apoenzyme preparation may be due to a small amount of flavin in unsupplemented LB or the cytosol of washed cells. Attempts to eliminate the latter source by incubation of washed cells in unsupplemented LB for 1 or 4 h at 15 °C prior to enzyme induction reduced the yield but did not affect the properties of the isolated apoprotein

To test for reconstitutability, the apoprotein was incubated with excess FAD (500  $\mu\text{M}$ ) at 23 °C, aliquots were removed at various times and assayed for catalytic activity. A time-dependent increase in activity was observed in a reaction that exhibits apparent first order kinetics ( $k_{\text{obs}} = 0.078 \pm 0.004 \text{ min}^{-1}$ ) (Figure 3). The final specific activity attained in this experiment ( $\text{SA} = 15 \text{ U/mg}$ ) was about 45% of that observed with native enzyme. Somewhat lower values ( $\sim 12 \text{ U/mL}$ ) were obtained after long term storage. The results show that apoMSOX could be reconstituted with FAD.

### Isolation and Properties of FAD-reconstituted MSOX

A larger scale reconstitution reaction was conducted, unbound flavin was removed by ultrafiltration and the preparation was then centrifuged to remove any insoluble protein. The reconstituted enzyme was isolated in about 50% yield and exhibited a specific activity (27.0 U/ml) that was 78% of that observed for native enzyme (Table 2). The higher specific activity of the isolated reconstituted enzyme suggests that inactive protein is removed by centrifugation. The absorption spectrum of reconstituted MSOX exhibits maxima at 454, 373 and 275 nm with a pronounced shoulder at 475 (Figure 4, curve 1), similar to that observed for native enzyme.

The reconstituted enzyme was denatured with guanidine hydrochloride to determine whether the flavin was covalently attached to the protein. Denaturation results in a hypsochromic shift of both visible absorption maxima and the loss of the shoulder at 475 nm (Figure 4, curve 2). A colorless filtrate was obtained upon ultrafiltration of the denatured reconstituted enzyme. Both protein and flavin were found in the yellow-colored retentate which exhibited a characteristic flavin absorption spectrum (data not shown). The results show that the FAD in reconstituted MSOX is covalently attached to the protein. The data provide definitive evidence for autoflavinylation in a reaction that requires only apoenzyme and FAD.

The spectral changes observed after denaturation of reconstituted MSOX were used to estimate the extinction coefficient of the intact reconstituted enzyme at 454 nm. The calculated value ( $\epsilon_{454} = 12,300 \text{ M}^{-1} \text{ cm}^{-1}$ ) is virtually identical to that previously reported for native MSOX (Table 2). The flavin content of the reconstituted enzyme is about 80% of that observed for native MSOX, as judged by values obtained for the ratio,  $A_{280}/A_{454}$ , or the estimated molar ratio of flavin to protein (Table 2). The somewhat lower flavin content in reconstituted enzyme is consistent with the observed difference in its specific activity as compared with native MSOX (Table 2).

Apparent steady state kinetic parameters for reconstituted MSOX and native enzyme were determined by monitoring the rate of hydrogen peroxide formation at various sarcosine concentrations in air-saturated buffer. In these studies, enzyme concentration was estimated based on absorption at 454 nm in order to compensate for differences in flavin content. Importantly, reconstituted MSOX exhibits apparent  $k_{\text{cat}}$  and  $K_{\text{m}}$  values that are virtually identical to those observed with native enzyme (Table 2).

### Effect of FAD Concentration of the Rate of Covalent Flavinylation

In addition to the covalent flavin linkage, there are numerous noncovalent interactions between FAD and the protein, as judged by the crystal structure of native MSOX (10). The observed noncovalent flavin contacts suggested that apoMSOX might form an initial noncovalent complex with FAD prior to covalent attachment. In this case, the rate of covalent flavinylation should exhibit saturation kinetics with respect to FAD. To evaluate this hypothesis, apoMSOX was incubated with various concentrations of FAD in the range from 50 to 500  $\mu\text{M}$ . The rate of reconstitution was measured by monitoring the time-dependent increase in catalytic activity. Contrary to our expectation, the observed apparent first order rate constants exhibited a linear ( $r^2 = 0.97$ ) dependence on the FAD concentration (Figure 5). However, the plot does not go through the origin, suggesting that covalent flavin attachment may involve initial formation of an unstable apoMSOX·FAD complex. An apparent second order rate constant of  $139 \text{ M}^{-1} \text{ min}^{-1}$  was estimated from the slope of the plot in Figure 5.

Catalytic activity, albeit at reduced rates, is observed with mutant forms of other enzymes that bind flavin noncovalently when the normal flavin attachment site is eliminated by mutagenesis (28–32). These studies suggested that addition of FAD to the catalytic assay mixture might stimulate sarcosine oxidation by apoMSOX. FAD concentrations in the range from 25 to 1000  $\mu\text{M}$  were tested. No effect on the initial rate of sarcosine oxidation was detected, consistent with very weak noncovalent binding of FAD to apoMSOX.

### Is a Reduced Flavin Intermediate Formed during Reconstitution of ApoMSOX with FAD?

Covalent flavin attachment is postulated to involve the formation of a reduced flavin intermediate (Scheme 1, intermediate II). We reasoned that rapid oxidation of the putative reduced flavin intermediate should occur under aerobic conditions, accompanied by the reduction of oxygen to hydrogen peroxide. Control studies showed that hydrogen peroxide could be measured in the presence of excess FAD by using a horseradish peroxidase-coupled assay with Amplex Red as the chromogenic substrate. Oxidation of Amplex Red to resorufin results in a large increase in absorbance at 563 nm ( $\Delta\epsilon_{563} = 52,200 \text{ M}^{-1} \text{ cm}^{-1}$ ), a wavelength where FAD exhibits minimal absorbance.

Significantly, hydrogen peroxide formation was initiated by the addition of apoMSOX to reaction mixtures containing excess FAD, horseradish peroxidase and Amplex Red, as judged by the time-dependent increase in absorbance at 563 nm. The reaction exhibited biphasic kinetics with 85% of the observed change occurring in the initial fast phase ( $k_{\text{fast}} = 0.142 \pm 0.002 \text{ min}^{-1}$ ), followed by a 20-fold slower phase ( $k_{\text{slow}} = 7.3 \pm 0.1 \times 10^{-3} \text{ min}^{-1}$ ). The slow phase is largely attributable to a blank reaction observed when apoMSOX is added to hydrogen peroxide assay mixtures in the absence of FAD ( $k = 5.8 \pm 0.2 \times 10^{-3} \text{ min}^{-1}$ ) (Figure 6). The fast phase of hydrogen peroxide formation was 1.8-fold faster than the rate of reconstitution, as estimated based on the increase in enzyme activity observed under nearly the same conditions ( $k = 0.078 \pm 0.004 \text{ min}^{-1}$  at 500  $\mu\text{M}$  FAD). However, hydrogen peroxide assay mixtures included a small amount of dimethylsulfoxide (0.4%), the solvent used to prepare stock solutions of Amplex Red. The rate of reconstitution was, therefore, re-measured in presence of 0.4% dimethylsulfoxide by monitoring the increase in enzyme activity. The rate observed under these conditions ( $k = 0.120 \pm 0.006 \text{ min}^{-1}$ ) was in good agreement with the value obtained for the fast phase of hydrogen peroxide formation. The amount of hydrogen peroxide produced in the fast phase (3.6 nmol/mL) was stoichiometric with the amount of reconstituted MSOX formed under the same conditions (3.5 nmol/mL), as estimated based on the increase in catalytic activity.

The spectral course of the reconstitution of apoMSOX with FAD was monitored under anaerobic conditions in an attempt to directly detect the postulated reduced flavin intermediate.



A very high concentration of apoMSOX (276  $\mu\text{M}$ ) was used in this experiment in order to achieve an appreciable reaction rate at a moderate FAD concentration (120  $\mu\text{M}$ ) that would be compatible with spectral measurements using a cuvette with a 1-cm light path. Indeed, the anaerobic reaction of apoMSOX with FAD resulted a time-dependent reduction of the flavin (Figure 7). The amount of reduced flavin formed (55.8  $\mu\text{M}$ ) was estimated based on the observed decrease in absorbance at 450 nm and the difference in the extinction coefficient of free oxidized FAD *versus* MSOX-bound reduced FAD ( $\Delta\epsilon_{450} = 10,300 \text{ M}^{-1} \text{ cm}^{-1}$ ). This value is about 50% of the value calculated for the amount of reconstituted MSOX (124.6  $\mu\text{M}$ ), as estimated based on the increase in catalytic activity. This discrepancy may be due to difficulties in removing oxygen from a very concentrated protein solution that could not be overcome by including glucose/glucose oxidase as an oxygen scavenger. Immediate oxidation of the reduced flavin was observed upon aeration of the sample. Comparison of the initial spectrum (Figure 7, curve 1) with that observed after aeration (Figure 7, curve 7) shows that covalent flavinylation is accompanied by a small bathochromic shift and a small increase in the intensity of the 450 nm absorption band of free FAD. The observed change is consistent with differences in the spectral properties of free *versus* MSOX-bound FAD ( $\lambda_{\text{max}} = 450$  *versus* 454 nm;  $\epsilon_{450} = 11,300$  *versus*  $\epsilon_{454} = 12,200 \text{ M}^{-1} \text{ cm}^{-1}$ ). The magnitude of the observed spectral change ( $A_{454} \text{ reoxidized} - A_{450} \text{ initial} = 0.082$ ) is 73% of the value calculated based on the estimated concentration of reconstituted MSOX.

### Attempted Reconstitution of ApoMSOX with 5-deazaFAD or 1-deazaFAD

5-DeazaFAD and 1-deazaFAD contain carbon in place of nitrogen at positions 5 and 1 of the flavin ring, respectively (see Table 1 for structures). No evidence for incorporation of 5-deazaFAD or 1-deazaFAD into apoMSOX was obtained even after prolonged incubation (24 h) under conditions (500  $\mu\text{M}$  flavin in 100 mM potassium phosphate, pH 8.0 at 23 °C) where flavinylation is readily observed with FAD. Failure to observe flavinylation with 5-deazaFAD or 1-deazaFAD might reflect the inability of these derivatives to bind noncovalently to apoMSOX. This possibility was investigated by determining whether the analogs could inhibit covalent attachment of FAD. In these experiments, the rate of apoMSOX reconstitution with 100  $\mu\text{M}$  FAD was monitored by measuring the increase in catalytic activity in absence or presence of 500  $\mu\text{M}$  5-deazaFAD or 1-deazaFAD. The rate of reconstitution in the presence of 5-deazaFAD and 1-deazaFAD was 45% and 62%, respectively, of that observed with FAD alone. The observed competition clearly indicates that the analogs can bind noncovalently to apoMSOX.

### Reconstitution of ApoMSOX with 8-Nor-8-chloroFAD

The 8 $\alpha$ -methyl group of FAD is replaced by a chlorine atom in 8-nor-8-chloroFAD (see Table 1 for structure). This FAD analog exhibits a typical flavin absorption spectrum with two maxima in the visible region ( $\lambda_{\text{max}} = 448, 365 \text{ nm}$ ) (Figure 8, curve 1). Dramatically different spectral properties are, however, observed after reconstitution of apoMSOX with 8-nor-8-chloroFAD. The reconstituted enzyme exhibited a single maximum in the visible region at 475 nm (Figure 8, curve 2) with a shoulder around 460 nm. Denaturation with guanidine hydrochloride shifted the absorption maximum to 480 nm, resulting in a more pronounced shoulder at 460 nm (Figure 8, curve 3). The absorption spectrum of the denatured enzyme is virtually identical to that reported for 8-nor-8-S-(N-acetylcysteinyl)-FAD (23) (see Table 1 for structure). The results suggested that flavinylation of the apoprotein had occurred in reaction involving nucleophilic displacement of the 8-chloro substituent by a cysteine residue (Scheme 2). Consistent with this hypothesis, a colorless filtrate was obtained after ultrafiltration of the denatured enzyme. The 480 nm-absorbing chromophore was found in the retentate, along with the MSOX protein. Unlike flavinylation with FAD, the postulated nucleophilic displacement reaction with 8-nor-8-chloroFAD is not expected to generate a reduced flavin intermediate. Indeed, little or no hydrogen peroxide was detected during reconstitution of apoMSOX with

8-nor-8-chloroFAD (Figure 6). Furthermore, flavin reduction was not observed when the reaction with 8-nor-8-chloroFAD was conducted under anaerobic conditions. The results provide strong evidence for the formation of a 8-nor-8-S-thioflavin linkage.

The absorbance change observed upon denaturation and the spectral properties reported for free 8-nor-8-S-(N-acetylcysteinyl)-FAD were used to calculate an extinction coefficient for the intact reconstituted enzyme at 475 nm ( $\epsilon_{475} = 25,600 \text{ M}^{-1} \text{ cm}^{-1}$ ). Enzyme reconstituted with 8-nor-8-chloroFAD exhibited a somewhat higher flavin to protein ratio as compared with FAD-reconstituted enzyme but was considerably less active as a catalyst (Table 2). The specific activity and apparent  $k_{\text{cat}}$  for sarcosine oxidation with 8-nor-8-chloroFAD-reconstituted enzyme are about an order of magnitude lower and the apparent  $K_m$  for sarcosine is more than 2-fold larger than observed with native enzyme.

### Kinetics of the Reaction of ApoMSOX with 8-Nor-8-chloroFAD

Formation of the 8-nor-8-S-thioflavin linkage is accompanied by a large spectral change. This suggested that it might be possible to directly monitor the kinetics of the flavinylation reaction at an appropriate wavelength, despite the presence of excess 8-nor-8-chloroFAD. The calculated difference spectrum for the reaction exhibits a maximum at 480 nm and a less intense minimum at 358 nm (Figure 8, inset). A wavelength as close as possible to the difference maximum at 480 nm was selected ( $\lambda = 499 \text{ nm}$ ) such that the initial absorbance due to 8-nor-8-chloroFAD (500  $\mu\text{M}$ ) was about 1. The flavinylation reaction, monitored at 499 nm, exhibited biphasic kinetics with 65% of the absorbance increase occurring in a fast initial phase ( $k_{\text{fast}} = 0.48 \pm 0.02 \text{ min}^{-1}$ ), followed by a 70-fold slower phase ( $k_{\text{slow}} = 6.8 \pm 1.4 \times 10^{-3} \text{ min}^{-1}$ ) (Figure 9). The fast phase is about 6-fold faster than the rate of flavinylation observed with FAD at the same concentration.

MSOX contains three cysteine residues (Cys315, Cys152, Cys262). Cys315 is the site of covalent flavin attachment in native MSOX (12) and the only cysteine residue at the active site (10). Cys315 is likely to be the most reactive cysteine in apoMSOX and responsible for the fast phase of the reaction observed with 8-nor-8-chloroFAD. Unlike FAD, free 8-nor-8-chloroFAD is known to react with thiols (33). This suggested that a non-active site cysteine residue might account for the slow phase of the MSOX reaction with 8-nor-8-chloroFAD. This hypothesis predicts that a portion of the 8-nor-8-S-cysteinyl-FAD in reconstituted enzyme would not be reducible by substrate. To test this hypothesis, an aerobic sample of reconstituted enzyme (0.27 mM oxygen) was mixed with a large excess of sarcosine (500 mM) in a stoppered cuvette. An isosbestic reduction of the covalently bound flavin was observed in a reaction that was complete within 20 min. However, further reduction occurred immediately after addition of solid dithionite (Figure 10). Reaction of FAD-reconstituted enzyme (Figure 10, inset) or native MSOX (data not shown) with sarcosine under the same conditions resulted in an immediate 89% or 90% bleaching, respectively, of the absorbance due to the oxidized enzyme at 454 nm. No further decrease at 454 nm was observed upon incubation or after addition of solid dithionite. The results show that all of the flavin in FAD-reconstituted and native MSOX is reducible by substrate. In contrast, about 15% of the flavin in 8-nor-8-chloroFAD-reconstituted enzyme cannot be reduced by sarcosine, as judged by the difference in absorbance at 475 nm observed with substrate- versus dithionite-reduced enzyme.

## DISCUSSION

Biosynthesis of a reconstitutible apoenzyme form of MSOX with a C-terminal affinity tag has been achieved by tightly regulated expression using a riboflavin-dependent strain of *E. coli* as host cell. The transformed cells are grown to a desired density at 37 °C in LB medium supplemented with riboflavin, harvested, washed and then transferred to unsupplemented LB at 15 °C for induction of MSOX expression. Large amounts of soluble apoprotein are produced

under these conditions (40 mg/liter of culture) whereas inclusion bodies are formed at higher induction temperatures. The pure protein, isolated by a single affinity chromatography step, is predominantly apoMSOX, as judged by its low residual catalytic activity and visible absorbance.

A time-dependent increase in catalytic activity is observed upon incubation of the apoenzyme with FAD, accompanied by the covalent incorporation of FAD to about 80% of the level observed with native enzyme. Except for a somewhat lower flavin content, the spectral and catalytic properties of the reconstituted enzyme are virtually indistinguishable from those observed with native MSOX. The results provide definitive evidence for autoflavinylation in a reaction that requires *only* apoMSOX and FAD. Surprisingly, the observed rate of flavinylation exhibits a linear, rather than an anticipated hyperbolic, dependence on the concentration of FAD ( $k = 139 \text{ M}^{-1} \text{ min}^{-1}$  at  $23^\circ\text{C}$ ). However, the plot does not intersect the origin, as would be expected for a second order reaction, suggesting that covalent attachment may involve prior formation of an unstable noncovalent E-FAD complex. In contrast, the flavin-binding subunit from *p*-cresol methylhydroxylase exhibits only noncovalent binding with FAD (34). Covalent attachment of FAD is triggered by a chaperone-like action of the heme-binding subunit that induces a conformational change in the flavin-binding subunit (13).

Strong evidence for a reduced flavin intermediate in the flavinylation reaction (Scheme 1, intermediate II) is provided by the similar reconstitution rates observed by measuring hydrogen peroxide formation or the increase in sarcosine oxidase activity, by the observed stoichiometric production of hydrogen peroxide and covalently bound FAD and by the time-dependent reduction of FAD observed when reconstitution is conducted under anaerobic conditions. Since the reaction is facilitated by the apoprotein, flavinylation of apoMSOX can be regarded as an autocatalytic process. Of course, the apoprotein is not a true catalyst since it is modified during the reaction, analogous to single turnover proteins like  $\text{O}^6$ -methylguanine-DNA alkyltransferases that repair alkylated DNA by transferring the alkyl group to one of their own cysteine residues (35). It is nevertheless interesting that apoMSOX flavinylation and MSOX-catalyzed amine oxidation both involve formation of a reduced flavin intermediate that reacts rapidly with oxygen to generate oxidized flavin and hydrogen peroxide.

Further insight into the mechanism of covalent flavin attachment was sought in studies with FAD analogs. Flavinylation is not observed with 5-deazaFAD or 1-deazaFAD, even after prolonged incubation with apoMSOX, although the analogs do bind noncovalently to apoMSOX as judged by their ability to inhibit flavinylation with FAD. The same outcome was reported for 5-deazaFAD in *in vitro* studies with apo-*p*-cresol methylhydroxylase (36) and apo-6-hydroxy-D-nicotine oxidase (14). Similarly, no flavin incorporation was observed when monoamine oxidase (A or B) was expressed in a riboflavin-dependent yeast strain grown in the presence of 5-deazariboflavin (16). Reduced 5-deazaflavins exhibit greatly diminished reactivity with oxygen as compared with unmodified flavins. It has been suggested that the failure to observe covalent attachment of 5-deazaFAD to monoamine oxidase can be attributed to the inability of the reduced flavin intermediate to be re-oxidized (16). This hypothesis cannot account for the results obtained with apoMSOX and 1-deazaFAD since reduced 1-deazaflavins exhibit high reactivity with oxygen, similar to that observed with unmodified flavin. 5-Deazaflavins and 1-deazaflavins do, however, share one property in common. Both substitutions result in a large decrease in redox potential as compared with unmodified flavin ( $\Delta E_{m,7} \sim -80 \text{ mV}$ , see Table 1). Proton abstraction from the  $8\alpha$ -methyl group is required for formation of the iminoquinone methide intermediate (Scheme 1, intermediate I) in the mechanism postulated for covalent flavin attachment. The acidity of the  $8\alpha$ -methyl group protons is enhanced by electron delocalization from the  $8\alpha\text{-CH}_2^-$  carbanion into the flavin ring. This property can account for the *selective* exchange of the  $8\alpha$ -methyl group protons with

solvent, as reported for free FMN in D<sub>2</sub>O (37). Since 5-deazaflavins and 1-deazaflavins are much poorer electron acceptors, these substitutions are likely to decrease the acidity of the 8 $\alpha$ -methyl group protons and therefore interfere with covalent flavin attachment. Consistent with this hypothesis, an enhanced rate of flavinylation of apoMSOX is observed when FAD is replaced by an analog with a higher redox potential<sup>2</sup>.

The 8 $\alpha$ -methyl group in FAD is replaced by a chlorine atom in 8-nor-8-chloroFAD, a substitution that blocks the normal mode of flavin attachment. Covalent flavin attachment is, however, observed in a reaction thought to involve displacement of the 8-chloro substituent by cysteine thiolate (Scheme 2), similar to the reaction observed with free 8-nor-8-chloroflavin and various thiolates (33). The aromatic nucleophilic displacement reaction proceeds via a quinoid intermediate but does not involve a reduced flavin intermediate, unlike the flavinylation reaction with FAD. Formation of the postulated 8-nor-8-S-thioflavin linkage in the apoMSOX reaction with 8-nor-8-chloroFAD is strongly supported by the dramatic change in the absorption spectrum of the flavin analog, the absence of hydrogen peroxide formation and the similar spectral properties observed for reconstituted MSOX and 8-nor-8-S-(N-acetylcysteinyl)-FAD. The covalent flavins in 8-nor-8-chloroFAD-reconstituted enzyme and native MSOX are likely to exhibit similar redox potentials, as judged by data reported for the corresponding riboflavin derivatives (8-nor-8-S-cysteinyl- and 8 $\alpha$ -S-cysteinyl-riboflavin, respectively, see Table 1). The ~10-fold lower catalytic activity observed with 8-nor-8-chloroFAD-reconstituted enzyme may reflect a structural perturbation needed to accommodate the modified cys315-flavin linkage at the active site.

Flavinylation of apoMSOX with 8-nor-8-chloroFAD occurs mainly at Cys315 but also involves a secondary reaction with a 70-fold less reactive, non-active site cysteine (Cys262 or Cys152). Evidence for a secondary reaction is provided by the observed biphasic reconstitution kinetics and the presence of flavin in the reconstituted enzyme that cannot be reduced by sarcosine. The fast phase of the reconstitution reaction is attributed to Cys315. The slow phase is probably due to Cys262 since this residue is partially accessible to solvent whereas Cys152 is completely buried. The reconstituted enzyme contains only slightly more flavin as compared with enzyme reconstituted with FAD (0.68 *versus* 0.61 mol flavin/mol protein). We estimate that 85% of the flavin in enzyme reconstituted with 8-nor-8-chloroFAD can be reduced by substrate, as judged by the extent of bleaching of flavin absorbance at 475 nm observed with sarcosine- or dithionite-reduced enzyme. The fast phase of the reconstitution reaction, however, accounts for 65% of the spectral change observed at 499 nm. A couple of factors might account for the modest discrepancy in these values. Firstly, flavinylation at the active site may generate a more stable protein that is isolated in higher yield as compared to protein flavinylated at a secondary site. Alternatively, covalently bound flavin at the active site and the secondary site may exhibit significantly different extinction coefficients at 475 and 499 nm. Indeed, model studies show that the spectral properties of 8-nor-8-S-alkylthio-flavins are extremely sensitive to solvent polarity (27).

Covalent attachment to the native cysteinyl flavin attachment site in monoamine oxidase A is observed when the protein is expressed in a riboflavin-dependent yeast strain grown in the presence of 8-nor-8-chlororiboflavin (16). In contrast, covalent flavin attachment of 8-nor-8-chloroFAD is not observed with apoenzymes where tyrosyl or histidyl residues serve as the normal flavin attachment site, as judged by results obtained with apo-*p*-cresol methylhydroxylase (36) or apo-6-hydroxy-D-nicotine oxidase (14), respectively. Significantly, apoenzyme flavinylation is observed with 8-nor-8-chloroFAD when the native histidyl attachment site in 6-hydroxy-D-nicotine oxidase is mutated to cysteine (38). The results suggest that a more potent thiolate nucleophile is required for the nucleophilic aromatic

<sup>2</sup>Hassan-Abdallah, A., Bruckner, R.C. Zhao, G. and Jorns, M.S., unpublished results.

substitution reactions with chloride as leaving group as compared with nucleophilic addition to the iminoquinone methide intermediate in the normal flavinylation pathway.

A diverse and expanding group of enzymes, including numerous drug targets, have been found to contain covalently bound flavin. Experimental support for postulated intermediates and a detailed understanding of the role of the protein moiety in covalent flavin biosynthesis have been limited by the lack of a robust system for *in vitro* studies. This obstacle has been overcome by the development of an expression system that allows for the isolation of substantial amounts of a reconstitutable apoenzyme. Further study of the autocatalytic flavinylation reaction observed with apoMSOX may serve as a paradigm for other members of this important category of enzymes.

#### Acknowledgements

We thank Drs. John Lambooy, Wallace Ashton and Bill McIntire for generous gifts of 8-nor-8-chlororiboflavin, 1-deazariboflavin and FAD synthetase, respectively. We thank Drs. Xiao-Ping Zhang and Ayse Hacisalihoglu for their help in producing a riboflavin-dependent strain of *E. coli* BL21(DE3) and the preparation of 8-nor-8-chloroFAD, respectively.

#### References


1. Mewies M, McIntire WS, Scrutton NS. Covalent attachment of flavin adenine dinucleotide (FAD) and flavin mononucleotide (FMN) to enzymes - The current state of affairs. *Protein Sci* 1998;7:7–20. [PubMed: 9514256]
2. Bruckner RC, Zhao GH, Venci D, Jorns MS. Nikkomycin biosynthesis: Formation of a 4-electron oxidation product during turnover of nikD with its physiological substrate. *Biochemistry* 2004;43:9160–9167. [PubMed: 15248773]
3. Barquera B, Hase CC, Gennis RB. Expression and mutagenesis of the NqrC subunit of the NQR respiratory Na<sup>+</sup> pump from *Vibrio cholerae* with covalently attached FMN. *FEBS Letters* 2001;492:45–49. [PubMed: 11248234]
4. Khanna P, Jorns MS. Characterization of the FAD-containing N-methyltryptophan oxidase from *Escherichia coli*. *Biochemistry* 2001;40:1441–1450. [PubMed: 11170472]
5. Dodt G, Kim DG, Reimann SA, Reuber BE, McCabe K, Gould SJ, Mihalik SJ. L-pipecolic acid oxidase, a human enzyme essential for the degradation of L-pipecolic acid, is most similar to the monomeric sarcosine oxidases. *Biochem J* 2000;345:487–494. [PubMed: 10642506]
6. Venci D, Zhao G, Jorns MS. Molecular characterization of nikD, a new flavoenzyme important in the biosynthesis of nikkomycin antibiotics. *Biochemistry* 2002;41:15795–15802. [PubMed: 12501208]
7. Eschenbrenner M, Chlumsky LJ, Khanna P, Strasser F, Jorns MS. Organization of the multiple coenzymes and subunits and role of the covalent flavin link in the complex heterotetrameric sarcosine oxidase. *Biochemistry* 2001;40:5352–5367. [PubMed: 11330998]
8. Kvalnes-Krick, K. and Jorns, M. S. (1991) Role of the covalent and noncovalent flavins in sarcosine oxidase, in *Chemistry and Biochemistry of Flavoenzymes* (Muller, F., Ed.) pp 425–435, CRC Press, Inc., Boca Raton.
9. Fossati P, Prencipe L, Berti G. Enzymic creatinine assay: A new colorimetric method based on hydrogen peroxide measurement. *Clin Chem* 1983;29:1494–1496. [PubMed: 6872208]
10. Trickey P, Wagner MA, Jorns MS, Mathews FS. Monomeric sarcosine oxidase: Structure of a covalently-flavinylated secondary amine oxidizing enzyme. *Structure* 1999;7:331–345. [PubMed: 10368302]
11. Wagner MA, Trickey P, Chen Z, Mathews FS, Jorns MS. Monomeric sarcosine oxidase: 1. Flavin reactivity and active site binding determinants. *Biochemistry* 2000;39:8813–8824. [PubMed: 10913292]
12. Wagner MA, Khanna P, Jorns MS. Structure of the flavocoenzyme of two homologous amine oxidases: Monomeric sarcosine oxidase and N-methyltryptophan oxidase. *Biochemistry* 1999;38:5588–5595. [PubMed: 10220347]




13. Kim J, Fuller JH, Kuusk V, Cunane L, Chen ZW, Mathews FS, McIntire WS. The cytochrome subunit is necessary for covalent FAD attachment to the flavoprotein subunit of p-cresol methylhydroxylase. *J Biol Chem* 1995;270:31202–31209. [PubMed: 8537385]
14. Brandsch R, Bichler V. Autoflavinylation of apo-6-hydroxy-D-nicotine oxidase. *J Biol Chem* 1991;266:19056–19062. [PubMed: 1918024]
15. Robinson KM, Lemire BD. Covalent attachment of FAD to the yeast succinate dehydrogenase flavoprotein requires import into mitochondria, presequence removal, and folding. *J Biol Chem* 1996;271:4055–4060. [PubMed: 8626739]
16. Miller JR, Edmondson DE. Influence of flavin analogue structure on the catalytic activities and flavinylation reactions of recombinant human liver monoamine oxidases A and B. *J Biol Chem* 1999;274:23515–23525. [PubMed: 10438531]
17. Nandigama RK, Edmondson DE. Influence of FAD structure on its binding and activity with the C406A mutant of recombinant human liver monoamine oxidase A. *J Biol Chem* 2000;275:20527–20532. [PubMed: 10877844]
18. O'Brien DE, Weinstock LT, Cheng CC. Synthesis of 10-deazariboflavin and related 2,4-dioxypyrimido[4,5-b]quinolines. *J Heterocycl Chem* 1970;7:99–105.
19. Tartof KD, Hobbs CA. Improved media for growing plasmid and cosmid clones. *Focus* 1987;9:12–16.
20. Wagner MA, Jorns MS. Monomeric sarcosine oxidase: 2. Kinetic studies with sarcosine, alternate substrates and substrate analogs. *Biochemistry* 2000;39:8825–8829. [PubMed: 10913293]
21. Miller, J. H. (1972) *Experiments in Molecular Genetics* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
22. Molholt B. Isolation of amino acid auxotrophs of *E. coli* using ampicillin enrichment. *Microb Genet Bull* 1967;27:8–9.
23. Moore EG, Cardemil E, Massey V. Production of a covalent flavin linkage in lipoamide dehydrogenase. *J Biol Chem* 1978;253:6413–6422. [PubMed: 681358]
24. Spencer R, Fisher J, Walsh C. Preparation, characterization, and chemical properties of the flavin coenzyme analogues of 5-deazariboflavin, 5-deazariboflavin 5'-phosphate, and 5-deazariboflavin 5'-diphosphate, 5'-5'-adenosine ester. *Biochemistry* 1976;15:1043–1053. [PubMed: 3206]
25. Entsch B, Husain M, Ballou DP, Massey V, Walsh C. Oxygen reactivity of p-hydroxybenzoate hydroxylase containing 1-deaza-FAD. *J Biol Chem* 1980;255:1420–1429. [PubMed: 6766449]
26. Wagner MA, Jorns MS. Folate utilization by monomeric *versus* heterotetrameric sarcosine oxidases. *Arch Biochem Biophys* 1997;342:176–181. [PubMed: 9185627]
27. Raibekas AA, Jorns MS. Affinity probing of flavin binding sites: I. Covalent attachment of 8-(methylsulfonyl)FAD to pig heart lipoamide dehydrogenase. *Biochemistry* 1994;33:12649–12655. [PubMed: 7918491]
28. Robinson KM, Rothery RA, Weiner JH, Lemire BD. The covalent attachment of FAD to the flavoprotein of *Saccharomyces cerevisiae* succinate dehydrogenase is not necessary for import and assembly into mitochondria. *Eur J Biochem* 1994;222:983–990. [PubMed: 8026509]
29. Blaut M, Whittaker K, Valdovinos A, Ackrell BAC, Gonsalus RP, Cecchini G. Fumarate reductase mutants of *Escherichia coli* that lack covalently bound flavin. *J Biol Chem* 1989;264:13599–13604. [PubMed: 2668268]
30. Fraaije MW, van den Heuvel RHH, Van Berkel WJH, Mattevi A. Structural analysis of flavinylation in vanillyl-alcohol oxidase. *J Biol Chem* 2000;275:38654–38658. [PubMed: 10984479]
31. Scrutton NS, Packman LC, Mathews FS, Rohlfs RJ, Hille R. Assembly of redox centers in the trimethylamine dehydrogenase of bacterium W(3)A(1) -Properties of the wild-type enzyme and a C30A mutant expressed from a cloned gene in *Escherichia coli*. *J Biol Chem* 1994;269:13942–13950. [PubMed: 8188674]
32. Efimov I, Cronin CN, McIntire WS. Effects of noncovalent and covalent FAD binding on the redox and catalytic properties of p-cresol methylhydroxylase. *Biochemistry* 2001;40:2155–2166. [PubMed: 11329284]
33. Moore EG, Ghisla S, Massey V. Properties of flavins where the 8-methyl group is replaced by mercapto-residues. *J Biol Chem* 1979;254:8173–8178. [PubMed: 475888]



34. Engst S, Kuusk V, Efimov I, Cronin CN, McIntire WS. Properties of p-cresol methylhydroxylase flavoprotein overproduced by *Escherichia coli*. *Biochemistry* 1999;38:16620–16628. [PubMed: 10600124]
35. Pegg AE. Repair of O<sup>6</sup>-alkylguanine by alkyltransferases. *Rev Mutat Res* 2000;462:83–100.
36. Efimov I, McIntire WS. A study of the spectral and redox properties and covalent flavinylation of the flavoprotein component of p-cresol methylhydroxylase reconstituted with FAD analogues. *Biochemistry* 2004;43:10532–10546. [PubMed: 15301551]
37. Bullock FJ, Jardetsky O. A experimental demonstration of the nuclear magnetic resonance assignments in the 6,7-dimethylisoalloxazine nucleus. *J Org Chem* 1965;30:2056–2057.
38. Stoltz M, Henninger HP, Brandsch R. The design of an alternative, covalently flavinylated 6-hydroxy-D-nicotine oxidase by replacing the FAD-binding histidine by cysteine and reconstitution of the holoenzyme with 8-(methylsulfonyl)FAD. *FEBS Letters* 1996;386(2–3):194–196. [PubMed: 8647280]
39. Walsh C, Fisher J, Spencer R, Graham DW, Ashton WT, Brown JE, Brown RD, Rogers EF. Chemical and enzymatic properties of riboflavin analogues. *Biochemistry* 1978;17:1942–1951. [PubMed: 207304]
40. Edmondson DE, Singer TP. Oxidation-reduction properties of 8- $\alpha$ -substituted flavins. *J Biol Chem* 1973;248:8144–8149. [PubMed: 4752947]
41. Draper RD, Ingraham LL. A potentiometric study of the flavin semiquinone equilibrium. *Arch Biochem Biophys* 1968;125:802–808. [PubMed: 5671043]
42. Stankovich MT, Massey V. Determination of the redox potential of deazariboflavin by equilibration with flavins. *Biochim Biophys Acta* 1976;452:335–344. [PubMed: 12806]
43. Spencer R, Fisher J, Walsh C. One- and two-electron redox chemistry of 1-carba-1-deazariboflavin. *Biochemistry* 1977;16:3586–3594. [PubMed: 19057]




**Figure 1.** Expression of apoMSOX. The SDS-15% polyacrylamide gel was stained for protein with Coomassie blue. Lane 1, native MSOX; Lane 2, whole cell extract of *E. coli* BL21 (DE3) *M1b*/pGZ26 cells, prepared as described in the text; Lanes 3 and 4, 5 and 10  $\mu$ g, respectively, of purified apoMSOX.




**Figure 2.**

Comparison of the visible absorption spectrum of apoMSOX (curve 1) with that observed for native enzyme (curve 2). Spectra were recorded in 100 mM potassium phosphate buffer, pH 8.0 at 23 °C and are normalized to the same absorbance at 280 nm.




**Figure 3.**

Reconstitution of apoMSOX with FAD. The reaction was initiated by adding apoMSOX (11.3  $\mu$ M) to 100 mM potassium phosphate buffer, pH 8.0, containing 500  $\mu$ M FAD at 23  $^{\circ}$ C. Aliquots were withdrawn at the indicated times and assayed for MSOX activity using a horseradish peroxidase assay (20). The solid line is a fit of the data (circles) to a single exponential equation  $[A = A_0 + \Delta A(1 - e^{-kt})]$  where  $A$  is the observed activity,  $A_0$  is the activity at  $t = 0$ ,  $\Delta A$  is the total increase in activity and  $k$  is the apparent first order rate constant. Each data point is the average of three duplicate runs.



**Figure 4.**


Spectral properties of MSOX reconstituted with FAD. Curve 1 is the absorption spectrum of reconstituted enzyme in 100 mM potassium phosphate buffer, pH 8.0 at 23 °C. Curve 2 was recorded after denaturation with 3 M guanidine hydrochloride.



**Figure 5.**


Effect of FAD concentration on the rate of reconstitution. The observed rate of reconstitution was estimated based on the increase in catalytic activity, as described in the legend to Figure 3, except that the FAD concentration was varied in the range from 50 to 500  $\mu\text{M}$ .






**Figure 6.**

Formation of hydrogen peroxide during reconstitution of apoMSOX with FAD or 8-nor-8-chloroFAD. Reactions were initiated by adding apoMSOX (11.3  $\mu$ M) to 100 mM potassium phosphate buffer, pH 8.0, containing 0 or 500  $\mu$ M FAD or 500  $\mu$ M 8-nor-8-chloroFAD, horseradish peroxidase (18 U/ml), Amplex Red (80  $\mu$ M) and 0.4% dimethylsulfoxide at 23  $^{\circ}$ C. Hydrogen peroxide formation was monitored by following the oxidation of Amplex Red at 563 nm. The spectral change observed in the presence of FAD (red line) was fitted to a biphasic exponential equation (black line). The spectral change observed in the absence of FAD (green line) was fitted to a single exponential equation (black line). The blue line shows the spectral change observed with 8-nor-8-chloroFAD.




**Figure 7.**

Anaerobic reconstitution of apoMSOX with FAD. Curve 1 is the absorption spectrum recorded immediately after mixing 276  $\mu\text{M}$  apoMSOX with 120  $\mu\text{M}$  FAD in 100 mM potassium phosphate buffer, pH 8.0, containing 50 mM glucose and glucose oxidase (0.5 U/mL) at 23  $^{\circ}\text{C}$ . Curves 2–6 were recorded 2.5, 5.0, 10.0, 20.0 and 144 min, respectively, after mixing. Curve 7 was recorded after opening the cuvette to air. All spectra are corrected for the residual absorbance of apoMSOX (see Figure 2, curve 1).




**Figure 8.**

Comparison of the spectral properties of free 8-nor-8-chloroFAD (curve 1) with 8-nor-8-chloroFAD-reconstituted MSOX (curve 2). Curve 3 was obtained after denaturation of the reconstituted enzyme with 3 M guanidine hydrochloride. Spectra were recorded in 100 mM potassium phosphate buffer, pH 8.0, at 23 °C and are normalized to the same flavin concentration (21.8  $\mu\text{M}$ ). The inset shows the difference spectrum obtained by subtracting the spectrum of free 8-nor-8-chloroFAD from that observed for the intact, reconstituted enzyme.




**Figure 9.**

Kinetics of the reaction of apoMSOX with 8-nor-8-chloroFAD. The reaction was initiated by adding apoMSOX (11.3  $\mu$ M) to 100 mM potassium phosphate buffer, pH 8.0, containing 500  $\mu$ M 8-nor-8-chloroFAD at 23  $^{\circ}$ C and monitored at 499 nm. The solid line is a fit of the data (circles) to a biphasic exponential equation.




**Figure 10.**

Reduction of reconstituted enzymes with sarcosine. Curve 1 is the absorption spectrum of enzyme reconstituted with 8-nor-8-chloroFAD in 100 mM potassium phosphate buffer, pH 8.0, at 23 °C. Curves 2–7 were recorded 1, 2, 3, 4, 5 and 20 min, respectively after addition of 500 mM sarcosine to the aerobic sample in a tightly stoppered cuvette. No additional change was observed upon further incubation. (The arrow indicates the direction of absorbance change at wavelengths below 384 nm, the isosbestic point of the reaction.) Curve 8 was recorded immediately after adding solid dithionite. The inset shows absorption spectra obtained with FAD-reconstituted enzyme under the same conditions. Curves 1 and 2 were recorded before and immediately after, respectively, adding 500 mM sarcosine. No further decrease of the 454 nm band was observed upon addition of solid dithionite (data not shown).

**Scheme 1.**







Postulated mechanism for covalent flavinylation of MSOX with FAD.



**Scheme 2.**

Covalent attachment of 8-nor-8-chloroFAD to MSOX via a mechanism involving nucleophilic aromatic substitution.

**Table 1**  
Structure and redox potential of selected flavin derivatives<sup>a</sup>

Flavin	$E_{m,7}$ (mV)	Ref.
 8-nor-8-chloroflavin	-144	(39)
 8α-S-cysteinyflavin	-169	(40)
 8-nor-8-S-(N-acetylcysteiny)-O-flavin	-180	(33)
 unmodified flavin	-199	(41)
 5-deazaflavin	-273	(42)
 1-deazaflavin	-280	(43)

<sup>a</sup>Redox potentials are for the corresponding riboflavin derivatives. The carboxyl and amino groups in structure shown for 8α-S-cysteinyflavin are replaced by peptide bonds in the 8α-S-cysteinyflavin-FAD covalently attached to Cys315 in MSOX.

**Table 2**  
Spectral and catalytic properties of native MSOX, apoMSOX and reconstituted enzyme<sup>a</sup>

Preparation	$\epsilon_{4XY}$ (M <sup>-1</sup> cm <sup>-1</sup> )	A <sub>280</sub> /A <sub>4XY</sub>	mol flavin/mol protein	Specific Activity (U/mg)	k <sub>cat</sub> (app) (min <sup>-1</sup> )	K <sub>m</sub> (app) (mM)
native MSOX	12,200 <sup>b</sup>	5.7 (5.5) <sup>b</sup>	0.80 (0.87) <sup>b</sup>	34.7	2510 ± 10 (2730 ± 32) <sup>c</sup>	5.6 ± 0.1 (4.5 ± 0.1) <sup>c</sup>
apoMSOX	nd	73	nd	1.1	nd	nd
FAD-reconstituted	12,300	7.3	0.61	27.0	2330 ± 20	5.2 ± 0.1
8-nor-8-chloroFAD-reconstituted	25,600	3.1	0.68	3.0	392 ± 7	13 ± 0.6

<sup>a</sup> Specific activity and apparent steady state kinetic parameters were determined using a horseradish peroxidase assay with *o*-dianisidine as the chromogenic substrate (20). 4XY = 454 nm for FAD-reconstituted, native MSOX and apoMSOX; 4XY = 475 nm for 8-nor-8-chloroFAD-reconstituted MSOX.

<sup>b</sup> Values previously reported by Wagner et al. (12)

<sup>c</sup> Values previously reported by Wagner and Jorns (20)