

Published in final edited form as:

Biochemistry. 2005 August 30; 44(34): 11546–11554. doi:10.1021/bi0508111.

# IMPACT OF THE C-TERMINAL DOMAIN OF TOPOISOMERASE IIα ON THE DNA CLEAVAGE ACTIVITY OF THE HUMAN ENZYME†

Jennifer S. Dickey§ and Neil Osheroff§,¶,\*

Departments of Biochemistry and Medicine (Hematology/Oncology), Vanderbilt University School of Medicine, Nashville, Tennessee 37232-0146

*SDepartment of Biochemistry.* 

¶Department of Medicine (Hematology/Oncology).

#### Abstract

The enzymatic function of the C-terminal domain of eukaryotic topoisomerase II is not well defined. This region of the enzyme is highly variable, hydrophilic, and contains nuclear localization signals and phosphorylation sites. In contrast to eukaryotic topoisomerase II, type II enzymes from chlorella virus completely lack the C-terminal domain. These viral enzymes are characterized by a robust DNA cleavage activity, high coordination between their two active site tyrosyl residues, and reduced sensitivity to anticancer drugs. As a first step towards characterizing the contribution of the Cterminal domain of human topoisomerase IIα to enzyme function, the protein was truncated at amino acid 1175, which corresponds to the C-terminal residue of Paramecium bursaria chlorella virus-1 topoisomerase II as determined by BLAST sequence alignment. Although the overall catalytic activity of the resulting enzyme, hTop $2\alpha\Delta 1175$ , was lower than that of full-length topoisomerase IIα, the mutant protein displayed a double-stranded DNA cleavage activity that was ~2-to 3-fold higher. While the DNA breaks created by hTop2αΔ1175 were primarily double-stranded, cuts generated by topoisomerase IIa were primarily single-stranded. Thus, the enhanced cleavage observed for hTop $2\alpha\Delta 1175$  appears to be due, at least in part, to an increase in active site coordination. Finally, hTop2αΔ1175 displayed a distinctly lower susceptibility to anticancer agents than did topoisomerase  $II\alpha$ , despite the fact that it showed a similar binding affinity for etoposide. Therefore,

PBCV-1

Paramecium bursaria chlorella virus-1

CVM-1

Chlorella virus Marburg-1

**PCR** 

polymerase chain reaction.

<sup>&</sup>lt;sup>†</sup>This work was supported by National Institutes of Health research grant GM33944. JD was a trainee under grant 5 T32 HD07043 from the National Institutes of Health.

<sup>\*</sup>To whom correspondence should be addressed: Neil Osheroff, Department of Biochemistry, 654 Robinson Research Building, Vanderbilt University School of Medicine, Nashville, TN 37232-0146. Tel.: 615-322-4338, Fax: 615-343-1166, E-mail: neil.osheroff@vanderbilt.edu

<sup>&</sup>lt;sup>1</sup>The abbreviations used are

 $<sup>^2</sup>$ Although partially relaxed plasmid molecules can be seen in the reactions catalyzed by topoisomerase II $\alpha$  and hTop2 $\alpha\Delta1175$ , the DNA relaxation patterns generated by the two enzymes differ. In the reaction catalyzed by the full-length enzyme, the negatively supercoiled band is rapidly shifted to an intermediate superhelical density, which gradually shifts to the position of fully relaxed products. This pattern is indicative of a distributive reaction, in which topoisomerase II $\alpha$  dissociates from partially relaxed plasmids during the course of the assay. In contrast, in the reaction catalyzed by hTop2 $\alpha\Delta1175$ , the negatively supercoiled band persists and appears to be shifted primarily to the position of fully relaxed products with minimal accumulation of partially relaxed intermediates. This pattern is indicative of a processive reaction, in which hTop2 $\alpha\Delta1175$  remains largely bound to plasmids throughout the relaxation process.

the C-terminal domain of human topoisomerase  $II\alpha$  appears to play significant roles in modulating the DNA cleavage/ligation reaction of the enzyme and its response to anticancer agents.

Topoisomerase II is an essential enzyme that is responsible for resolving knots and tangles in chromosomal DNA that are generated during normal nuclear processes (1-7). Vertebrate species encode two distinct isoforms of the enzyme, topoisomerase II $\alpha$  and  $\beta$  (8,9). Topoisomerase II $\alpha$  expression is linked to the proliferative status of cells. Enzyme levels rise dramatically during periods of active growth and this isoform is required for replication and mitosis (10,11). In contrast, the expression of topoisomerase II $\beta$  appears to be independent of cell growth and the specific physiological functions of the enzyme have yet to be defined (12,13). Although this isoform is dispensable at the cellular level, deletion of topoisomerase II $\beta$  leads to neurological defects in developing mouse embryos (14).

Topoisomerase II functions by passing an intact double helix through a transient double-stranded break that it generates in a separate segment of DNA (2,3,5,7,15). Although the DNA strand passage activity of the enzyme is essential for cell survival, the double-stranded DNA breaks that it creates have the potential to destabilize the genome (3,5,7,16,17). As a result, topoisomerase II normally cleaves and ligates the genetic material very rapidly. However, physiological conditions that increase levels of topoisomerase II-associated DNA strand breaks convert the enzyme to a potent cellular toxin that fragments the genetic material (3,5,7,16, 18).

Because of its DNA cleavage activity, topoisomerase II plays a central role in cancer. The enzyme is the cytotoxic target for some of the most successful anticancer drugs currently in clinical use (3,5,7,19-21). All of these agents act by increasing cellular levels of topoisomerase II-mediated DNA strand breaks (3,5,7,19). In contrast, emerging evidence suggests that strand breaks generated by the enzyme may initiate chromosomal translocations that trigger specific types of leukemia (17,22-27).

It is difficult to study the DNA cleavage/ligation reaction of eukaryotic topoisomerase II because the enzyme normally maintains low equilibrium levels of strand breaks (2,3,5,7). Recently, however, novel viral type II topoisomerases were discovered that possess a dramatically (30– to 50–fold) higher DNA cleavage activity than their eukaryotic counterparts. These enzymes are encoded by the chlorella viruses, *Paramecium bursaria* chlorella virus-1 (PBCV-1)¹ and *Chlorella* virus Marburg-1 (CVM-1) (28-31). The basis for the high DNA cleavage activity of chlorella virus topoisomerase II is not known. However, both viral enzymes are considerably (~45-60 kDa) shorter than eukaryotic topoisomerase II and lack the entire C-terminal domain typical of the eukaryotic enzyme. The remaining portion of chlorella virus topoisomerase II displays nearly 50% amino acid sequence identity to human topoisomerase IIα. Furthermore, virtually every amino acid residue that is invariant or highly conserved within the eukaryotic type II family also is retained in chlorella virus topoisomerase II.

The enzymatic function of the C-terminal domain of eukaryotic topoisomerase II is not well defined. This region of the enzyme is highly variable, hydrophilic, and contains nuclear localization signals as well as phosphorylation sites (11,32,33). Based on structural studies, it has been proposed that the C-terminal domain of prokaryotic type II topoisomerases interacts with DNA, but comparable studies have not been carried out with eukaryotic topoisomerase II (34-37).

Human topoisomerase II $\alpha$  is 1581 amino acids in length. Truncation of the protein at residue 1296 yields an enzyme that is active, but unable to complement a yeast strain encoding a temperature-sensitive topoisomerase II (33). This failure to complement presumably results from deletion of nuclear localization signals. Truncation of the human enzyme at amino acid

1121 yields an enzyme that is neither able to complement a deficient yeast strain nor able to catalyze DNA strand passage (33). Similar findings have been reported for yeast and *Drosophila* topoisomerase II (33,38-41).

The impact of the C-terminal domain on the DNA cleavage/ligation reaction of human topoisomerase II  $\alpha$  has not been analyzed. However, the enhanced DNA cleavage activity of chlorella virus topoisomerase II suggests that the C-terminal region of the enzyme may affect its ability to generate DNA strand breaks. Therefore, the human enzyme was truncated at amino acid 1175, which corresponds to the terminal amino acid in PBCV-1 topoisomerase II as determined by BLAST sequence alignment. The double-stranded DNA cleavage activity of the truncated human enzyme, hTop2 $\alpha$  $\Delta$ 1175, was 2–3–fold higher than that of full-length topoisomerase II $\alpha$ . Increased cleavage appears to be due, at least in part, to a greater coordination between the two active site tyrosyl residues. Finally, similar to the viral enzyme, hTop2 $\alpha$  $\Delta$ 1175, displayed a decreased sensitivity toward anticancer drugs. On the basis of these results, it is concluded that the loss of the C-terminal domain contributes to the robust DNA scission activity and low drug sensitivity of chlorella virus topoisomerase II.

#### **EXPERIMENTAL PROCEDURES**

#### **Enzymes and Materials**

Negatively supercoiled pBR322 DNA was prepared using a Plasmid Mega Kit (Qiagen) as described by the manufacturer. Kinetoplast DNA (kDNA) was isolated from *Crithidia fasciculata* as described by Englund *et al.* (42). [ $\gamma$ -<sup>32</sup>P]ATP ( $\sim$ 5000 Ci/mmol) was obtained from ICN. Etoposide was from Sigma, amsacrine was from Bristol-Myers Squibb, and CP-115,953 was from Pfizer Global Research. Drugs were stored at 4°C as 10 or 20 mM stock solutions in 100% DMSO. Etoposide labeled with [ $^3$ H] (654 mCi/mmol) was obtained from Moravek Biochemicals and stored at —80 °C in 100% methanol. All other chemicals were analytical reagent grade.

#### Cloning of hTop2αΔ1175

A fragment of the human topoisomerase II $\alpha$  gene in the YepWob6 plasmid was digested with KpnI and SmaI (43). PCR was utilized to introduce a stop codon at 1175 and a SmaI cut size downstream of the stop codon. The following sequences were used for the forward and reverse primers respectively: CATTGGTCAGTTTGGTACCAG-

GCTACATGGTGGCAAGGATTCTGCTAG and CAGTGTATATTGCCCGGGATGATGTTA-

TTAATAAATGTAGCCAAGTCTTCTTTCCACAAATCTGATGG. DNA was denatured at 95 °C for 6 min and subjected to 30 cycles of PCR using the following program: denaturation at 95 °C for 1 min, annealing at 53 °C for 1 min and primer extension at 72 °C for 4 min. Following the last round of PCR, primers were allowed to extend for an additional 8 min at 72 °C and the final product was purified by gel electrophoresis. The new insert was digested with SmaI and KpnI and ligated into the previously digested YepWob6 plasmid, and transformed into *Escherichia coli* DH5 $\alpha$ . The truncated human topoisomerase II $\alpha$  construct was sequenced to confirm the primary structure of the gene. The resultant plasmid carrying  $hTop2\alpha\Delta1175$  was transformed into *Saccharomyces cerevisiae* JEL-1 $\Delta top1$  by heat shock.

#### **Expression and Purification of Recombinant Type II Topoisomerases**

PBCV-1 topoisomerase II was expressed in *S. cerevisiae* JEL-1 $\Delta top1$  and purified by a modification (31) of the procedure of Lavrukhin *et al.* (28). Human topoisomerase II $\alpha$  and hTop2 $\alpha\Delta$ 1175 were expressed in *S. cerevisiae* JEL-1 $\Delta top1$  and purified as described by Kingma *et al.* (44).

#### **Decatenation of Kinetoplast DNA (kDNA)**

DNA decatenation reactions were carried out as described by Miller  $\it et\,al.$  (45). Assay mixtures contained 1 mM ATP, 5 nM kDNA, and 0–20 nM topoisomerase II  $\alpha$  or hTop2 $\alpha\Delta1175$  in a total of 20  $\mu L$  of topoisomerase II activity buffer (10mM Tris-HCl, pH 7.9, 175 mM KCl, 5 mM MgCl $_2$ , 0.1 mM NaEDTA, 2.5% glycerol). Decatenation was initiated by the addition of enzyme, followed by incubation for 30 min at 37 °C. Reactions were terminated by the addition of 3  $\mu L$  of stop solution (0.5% SDS, 77 mM EDTA). Proteinase K was added (2  $\mu L$  of a 0.8 mg/mL solution) and reactions were incubated for 30 min at 45 °C to digest the enzyme. Samples were mixed with agarose gel loading buffer (60% sucrose in 10 mM Tris-HCl, pH 7.9) and subjected to electrophoresis in 1% agarose gels in 100 mM Tris-borate, pH 8.3, and 2 mM EDTA. Gels were stained for 30 min with 0.5  $\mu g/mL$  ethidium bromide and DNA bands were visualized by UV light and quantified using an Alpha Innotech digital imaging system.

#### **DNA Relaxation**

DNA relaxation assays were carried out as described by Fortune and Osheroff (46). For catalytic reactions, assay mixtures contained 1 nM topoisomerase IIa or hTop2a $\Delta$ 1175, 1 mM ATP, and 10 nM negatively supercoiled pBR322 DNA in a total of 20  $\mu$ L of topoisomerase II activity buffer. Samples were incubated at 37 °C for 0–30 min. For non-turnover reactions, assay mixtures contained 110 nM human enzymes, 5 nM negatively supercoiled pBR322 DNA, and 50  $\mu$ M ATP in a total of 20  $\mu$ L of topoisomerase II activity buffer. Samples were incubated for 0–60 s at 37 °C. In all cases, reactions were stopped by the addition of 3  $\mu$ L of stop solution. Samples were processed and products were quantified as in the preceding section.

#### **DNA Cleavage**

Plasmid DNA cleavage reactions were based on the procedure of Fortune and Osheroff (46). Unless stated otherwise, reaction mixtures contained 10 nM negatively supercoiled pBR322 DNA in a total of 20 µL of DNA cleavage buffer (human type II topoisomerases, 10 mM Tris-HCl, pH 7.9, 100 mM KCl, 5 mM MgCl<sub>2</sub>, 0.1 mM NaEDTA, and 2.5% glycerol; PBCV-1 topoisomerase II, 10 mM Tris-HCl, pH 8.5, 62.5 mM NaCl, 62.5 mM KCl, 0.1 mM NaEDTA, 2.5 mM MgCl<sub>2</sub>, and 2.5% glycerol). Where noted, 0-200 µM etoposide, amsacrine, or CP-115,953 was added to cleavage reactions. DNA cleavage was initiated by the addition of topoisomerase II (final enzyme concentrations were hTop2αΔ1175 or topoisomerase IIα, 0.2-1 μM; PBCV-1 topoisomerase II, 0.02 μM). Reaction mixtures were incubated for 6 min at 37 °C (human) or 25 °C (PBCV-1) to establish cleavage/religation equilibria. Cleavage intermediates were trapped by the addition of 2 µL of 5% SDS and 2 µL of 250 mM NaEDTA, pH 8.0, (human) or 2 μL of 1.15% SDS and 2 μL of 115 mM NaEDTA, pH 8.0 (PBCV-1). Proteinase K was added (2  $\mu$ L of 0.8 mg/mL) and mixtures were incubated for 30 min at 45  $^{\circ}$ C to digest the type II enzyme. Samples were mixed with 2 µL of agarose gel loading buffer, heated for 2 min at 45 °C, and subjected to electrophoresis in 1% agarose gels in 40 mM Trisacetate, pH 8.3, and 2 mM EDTA containing 0.5 µg/mL ethidium bromide. Double- and singlestranded DNA cleavage was monitored by the conversion of negatively supercoiled plasmid to linear and nicked circular molecules, respectively. DNA bands were quantified by digital imaging as described above.

DNA sites cleaved by hTop2 $\alpha\Delta$ 1175 or human topoisomerase II $\alpha$  were mapped as described by a modification (47) of the procedure of O'Reilly and Kreuzer (48). A linear 4330 bp fragment (HindIII/EcoRI) of pBR322 plasmid DNA singly labeled with [ $^{32}$ P]phosphate on the 5'-terminus of the HindIII site was used as the cleavage substrate. Reaction mixtures contained 0.35 nM DNA molecules and 60 nM enzyme in a total of 50  $\mu$ L of DNA cleavage buffer. Assays were carried out in the absence or presence of 50  $\mu$ M etoposide. Reactions were started by the addition of enzyme and mixtures were incubated for 10 min at 37 °C. Cleavage intermediates were trapped and processed as described above. Products were precipitated twice

in 100% ethanol, dried, resuspended in 40% formamide, 8.4 mM EDTA, 0.02% bromophenol blue, and 0.02% xylene cyanole FF, and subjected to electrophoresis in 6% polyacrylamide gels. Gels were fixed in 10% methanol/10% acetic acid for 5 min and dried. DNA cleavage products were visualized using a Bio-Rad Molecular Imager FX.

It should be noted that DNA cleavage reactions were performed in the absence of ATP. The nucleotide cofactor does not influence the mechanism of topoisomerase II-mediated DNA scission (5). Furthermore, the omission of ATP from cleavage assays is advantageous in that the topological state of the DNA is not altered during the reaction. Thus, negatively supercoiled substrates are neither relaxed nor catenated over the course of the assay (5). Alterations in DNA topology have the potential to become problematic, especially in light of the fact that the efficiency of DNA strand passage differs between full-length human topoisomerase II $\alpha$  and hTop2 $\alpha$  $\Delta$ 1175 (see Figures 2 and 3).

#### **Topoisomerase II-DNA Binding**

The ability of topoisomerase II $\alpha$  or hTop2 $\alpha\Delta$ 1175 to bind DNA was assessed using a nitrocellulose filter-binding assay (31). Linear pBR322 DNA radiolabeled with [\$^{32}P]phosphate was prepared as described in the preceding section. Binding mixtures contained 5 nM DNA and 10–400 nM enzyme in a total of 20  $\mu$ L of DNA binding buffer (10 mM Tris-HCl, pH 7.9, 5 mM KCl, 0.1 mM NaEDTA, and 2.5% glycerol) and were incubated for 6 min at 37 °C. Nitrocellulose membranes (0.45  $\mu$ m HA, Millipore) were prepared by incubation in DNA binding buffer for 10 min. Samples were applied to the membranes and filtered in vacuo. Membranes were washed 3 times with 1 mL of DNA binding buffer, dried, and submerged in 8 mL of scintillation fluid (Econo-Safe, Research Products International). Radioactivity remaining on the membranes was quantified using a Beckman LS 5000 TD Scintillation Counter. The percent DNA bound to topoisomerase II was determined based on the ratio of the radioactivity on the membranes vs. that of the input DNA.

#### **DNA Religation**

DNA religation mediated by topoisomerase II $\alpha$  or hTop2 $\alpha\Delta$ 1175 was monitored according to the procedure of Robinson and Osheroff (49). Enzyme—DNA cleavage/religation equilibria were established as above for plasmid DNA cleavage. Religation was initiated by shifting reactions from 37 °C to 0 °C. Reactions were stopped 0–30 s after the temperature shift by the addition of 2  $\mu$ L of 5% SDS followed by 2  $\mu$ L of 250 mM NaEDTA, pH 8.0. Samples were processed and analyzed as described for topoisomerase II plasmid DNA cleavage reactions. Religation was monitored by quantifying the loss of linear DNA.

#### **Topoisomerase II-Etoposide Binding**

The ability of human topoisomerase IIa, hTop2a $\Delta$ 1175, or PBCV-1 topoisomerase II to bind etoposide was monitored as described by Kingma and Osheroff (50). Nitrocellulose membranes (0.45  $\mu$ m HA) were presoaked in DNA cleavage buffer for 10 min. Binding mixtures contained 3  $\mu$ M enzyme and 25-100  $\mu$ M [ $^3$ H]etoposide in a total of 60  $\mu$ L of cleavage buffer. Samples were incubated at 25 °C for 6 min and applied to the nitrocellulose membranes in vacuo. Filters were immediately washed 3 times with 1 mL of ice-cold wash DNA cleavage buffer, dried, and submerged in 8 mL of scintillation fluid. Radioactivity remaining on the membranes was quantified as described above. The amount of radioactive etoposide remaining on the filter in the absence of enzyme was subtracted for binding calculations.

#### **RESULTS**

Because of the deleterious effects of DNA strand breaks in proliferating cells, eukaryotic type II topoisomerases normally maintain very low levels of DNA scission intermediates. Typically,

less than 0.2% of a population of human topoisomerase II $\alpha$  is in a DNA cleavage complex at any given time (2,3,5,7). In marked contrast, topoisomerase II from chlorella virus PBCV-1 or CVM-1 maintains a dramatically higher level of cleavage intermediates (typically >25%) (29-31). The major structural difference between the chlorella and human type II topoisomerases is the lack of a C-terminal domain in the viral enzyme. Therefore, to determine the impact of the C-terminal domain on the DNA cleavage activity of topoisomerase II $\alpha$ , we truncated the human enzyme at amino acid 1175. This position corresponds to the terminal residue in PBCV-1 topoisomerase II, as determined by BLAST sequence alignment. The resulting enzyme, hTop2 $\alpha$ Δ1175, was expressed in yeast and purified along with full-length topoisomerase II $\alpha$  and PBCV-1 topoisomerase II.

#### Catalytic Activity of hTop2αΔ1175

Before addressing the effects of the C-terminal domain on the DNA cleavage activity of human topoisomerase  $II\alpha$ , we determined whether the truncated enzyme retained overall catalytic activity. Two independent assays were used to assess the ability of hTop2 $\alpha\Delta$ 1175 to catalyze the DNA strand passage reaction. First, the ability of the truncated enzyme to decatenate kDNA was characterized (Figure 1). hTop2 $\alpha\Delta$ 1175 was able to unlink catenated DNA rings, albeit with an activity that was ~2-fold lower than that of full-length topoisomerase  $II\alpha$ .

To further characterize the influence of the C-terminal region on catalytic activity, the ability of hTop2 $\alpha\Delta$ 1175 to relax negatively supercoiled plasmid DNA was determined. Under optimal reaction conditions for human topoisomerase II $\alpha$ , hTop2 $\alpha\Delta$ 1175 was able to remove negative supercoils from plasmid molecules (Figure 2). However, reaction rates were lower than observed for the full-length enzyme. In addition, as determined by the loss of the negatively supercoiled band relative to the production of partially relaxed plasmid species, the reaction catalyzed by the truncated enzyme appeared to be more processive. This latter result was unexpected. If the C-terminal domain of eukaryotic topoisomerase II interacts with DNA, its presence might enhance enzyme-nucleic acid binding, and hence increase processivity. However, as discussed below, the assumption that the C-terminal domain of human topoisomerase II $\alpha$  increases the affinity of the enzyme for DNA is not correct.

The decreased catalytic activity of hTop2 $\alpha\Delta$ 1175 could be due to deficiencies in DNA strand passage or enzyme turnover. To distinguish between these two possibilities, a non-turnover DNA relaxation assay was utilized. Reaction mixtures contained a 22-fold stoichiometric excess of either full-length topoisomerase II $\alpha$  or hTop2 $\alpha\Delta$ 1175 over plasmid molecules. This excess eliminates the need for enzyme turnover and dissociation during the reaction, since complete relaxation of the plasmid can be achieved if each topoisomerase II molecule catalyzes only a single DNA strand passage event. Human topoisomerase II $\alpha$  relaxed the DNA substrate within 60 s (Figure 3). In contrast, hTop2 $\alpha\Delta$ 1175 displayed little DNA relaxation over the course of the assay. This finding suggests that hTop2 $\alpha\Delta$ 1175 catalyzes the DNA strand passage event more slowly than the full-length enzyme. Taken together, the above results confirm that the C-terminal domain of the human enzyme is dispensable for catalytic activity. However, this portion of the protein contributes to the catalytic efficiency of topoisomerase II $\alpha$ .

#### DNA Cleavage Activity of hTop2αΔ1175

The most dramatic enzymatic difference between human topoisomerase II $\alpha$  and chlorella virus topoisomerase II is the enhanced double-stranded DNA scission activity of the viral enzyme (29-31). Therefore, the ability of hTop2 $\alpha$  $\Delta$ 1175 to cleave DNA was compared to that of the full-length human enzyme. As seen in Figure 4, loss of the C-terminal domain of topoisomerase II $\alpha$  increased levels of double-stranded DNA cleavage. Over a range of enzyme concentrations, hTop2 $\alpha$  $\Delta$ 1175 generated 2 to 3 times more double-stranded DNA breaks than did full-length topoisomerase II $\alpha$ . While this difference does not approach the enhancement observed with

the viral enzyme (Figure 4, *insert*), the loss of the C-terminal domain clearly enhances the ability of topoisomerase  $II\alpha$  to cleave DNA.

#### **DNA Binding**

Several factors may contribute to the increased DNA cleavage activity of hTop $2\alpha\Delta1175$ . One possibility is that the truncated enzyme binds more tightly to its DNA substrate. To test this hypothesis, the ability of hTop $2\alpha\Delta1175$  to bind DNA was compared to that of full-length human topoisomerase II $\alpha$ . As seen in Figure 5, the affinity of the two enzymes for DNA was similar. Thus, the increase in DNA scission that accompanies the loss of the C-terminal domain is not due to an enhanced affinity for nucleic acids.

#### **DNA Religation**

A number of topoisomerase II-targeted anticancer drugs raise levels of enzyme-generated DNA strand breaks by inhibiting the religation of cleaved DNA molecules (3,5,7,51). Thus, the rate of DNA religation mediated by hTop2 $\alpha\Delta$ 1175 was determined (Figure 6). The truncated enzyme resealed DNA breaks at a rate that was slightly faster than that of human topoisomerase II $\alpha$ . This result indicates that the loss of the C-terminal domain does not increase levels of double-stranded DNA breaks by impairing the religation reaction of the enzyme.

#### **Coordination Between Enzyme Subunits**

PBCV-1 topoisomerase II displays significantly greater coordination between its two active site tyrosyl residues than does human topoisomerase II $\alpha$ . In this regard, the overwhelming majority of DNA breaks created by the chlorella virus enzyme are double-stranded in nature. Under optimal reaction conditions, PBCV-1 topoisomerase II generates nearly five double-stranded DNA breaks for every nick (i.e., single-stranded break) that it creates (Figure 7). In contrast, topoisomerase II $\alpha$  displays markedly lower coordination between its active site tyrosyl residues (52). For every double-stranded break that the human enzyme creates, it generates 2–5 nicks (i.e., cleavage intermediates in which one or the other active site tyrosyl residue, but not both, has cleaved the DNA) (Figure 7). It is not obvious how this coordination contributes to the catalytic activity of topoisomerase II $\alpha$ . However, the ability of the two active site tyrosyl residues to work in concert clearly should enhance the generation of double-stranded DNA breaks at the expense of single-stranded cuts.

To determine whether the C-terminal domain of topoisomerase II $\alpha$  affects active site coordination, the ratio of single-to double-stranded DNA breaks generated by hTop2 $\alpha\Delta$ 1175 was calculated. Results indicate that active site coordination of the truncated human enzyme was intermediate between those of full-length topoisomerase II $\alpha$  and PBCV-1 topoisomerase II. Based on the averages of three independent DNA cleavage titrations, hTop2 $\alpha\Delta$ 1175 produced approximately one nick for every double-stranded DNA break that it generated (Figure 7). This marked rise in active site coordination can account, at least in part, for the increase in double- stranded DNA breaks that accompany the loss of the C-terminal domain of topoisomerase II $\alpha$ .

#### Sensitivity of hTop2αΔ1175 to Anticancer Drugs

Compared to human topoisomerase II $\alpha$ , PBCV-1 topoisomerase II displays low sensitivity to DNA cleavage-enhancing anticancer drugs. When global levels of DNA cleavage are monitored using plasmid substrates, the viral enzyme is nearly refractory to a number of topoisomerase II-targeted agents (29).

It is not apparent why PBCV-1 topoisomerase II is resistant to anticancer agents. One possibility is that the C-terminal domain of the enzyme contributes to drug binding.

Consequently, the ability of topoisomerase II $\alpha$ , hTop2 $\alpha\Delta$ 1175, and PBCV-1 topoisomerase II to bind [ $^3$ H]etoposide was assessed using a nitrocellulose filter technique (Figure 8). Results indicate that the three enzymes bind similar amounts of the drug. Therefore, the insensitivity of the viral enzyme toward anticancer agents, at least etoposide, is not due to a lack of drug binding.

Etoposide, amsacrine, and CP-115,953 increase double-stranded DNA cleavage mediated by full-length topoisomerase II $\alpha$  5– to 10–fold (Figure 9). Since the C-terminal domain of the human enzyme does not appear to contribute to drug binding, the effects of these anticancer agents on the DNA cleavage activity of hTop2 $\alpha$ Δ1175 were determined. As seen in Figure 9, the truncated enzyme was less sensitive to the drugs. In most cases, the relative DNA cleavage enhancement observed was approximately 50% that seen with full-length topoisomerase II $\alpha$ .

Despite the diminished susceptibility of hTop2 $\alpha\Delta1175$  for anticancer drugs, the loss of the C-terminal domain does not appear to alter the DNA cleavage specificity of the truncated enzyme, either in the absence or presence of etoposide (Figure 10). It should be noted that the DNA samples shown in Figure 10 were subjected to electrophoresis under denaturing conditions. Therefore, the overall levels of DNA cleavage seen in the autoradiogram reflect both single-and double-stranded DNA breaks. Although hTop2 $\alpha\Delta1175$  generates higher levels of double-stranded DNA breaks than full-length human topoisomerase II $\alpha$ , the truncated enzyme produces considerably lower levels of single-stranded breaks. Consequently, the intensity of the DNA cleavage pattern observed with the full-length enzyme in the absence of etoposide is somewhat greater than that seen for hTop2 $\alpha\Delta1175$ .

Finally, in most cases, the intensity of etoposide-induced DNA cleavage bands generated by the truncated enzyme was lower than that of the corresponding bands produced by topoisomerase II $\alpha$ . However, in a few instances, the intensity of bands generated by hTop2 $\alpha\Delta$ 1175 in the presence of etoposide was comparable to or slightly greater than those produced by the full-length enzyme. Whether this result reflects differences in levels of single-stranded DNA cleavage or subtle effects of the C-terminal domain on DNA cleavage site utilization in the presence of anticancer drugs is not known at the present time.

Taken together, the above findings indicate that the C-terminal domain of topoisomerase II substantially influences the degree to which the enzyme responds to anticancer agents, but not the sites at which it forms DNA cleavage complexes.

#### DISCUSSION

The C-terminal domain of eukaryotic type II topoisomerases varies considerably from species to species (38,41,53,54). It is believed that this portion of the protein contributes to the physiology of topoisomerase II, as it contains phosphorylation sites and nuclear localization signals (11,32,33,55). However, the contribution of the C-terminal domain to the enzymatic activity of topoisomerase II has yet to be defined. Structural studies indicate that the C-terminal domain of prokaryotic type II topoisomerases interacts with nucleic acids (34-37). Unfortunately, comparable studies have not been carried out with eukaryotic enzymes.

In contrast to eukaryotic topoisomerase II, type II enzymes from chlorella virus completely lack the C-terminal domain. These viral enzymes are characterized by a robust DNA cleavage activity and a high coordination between their two active site tyrosyl residues (28-31). Therefore, as a first step toward characterizing the contribution of the C-terminal domain of human topoisomerase II $\alpha$  to enzyme function, the human protein was truncated at amino acid 1175, which corresponds to the C-terminal residue of PBCV-1 topoisomerase II.

The overall catalytic activity of the resulting enzyme, hTop2 $\alpha\Delta1175$ , was lower than that of the full-length protein, however, it displayed a double-stranded DNA cleavage activity that was ~2— to 3—fold higher than that of topoisomerase II $\alpha$ . While the truncated enzyme did not necessarily produce more total DNA breaks than its full-length counterpart, the cuts that it generated were primarily double-stranded in nature. In contrast, the DNA breaks created by topoisomerase II $\alpha$  were primarily single-stranded. Coupled with the findings that the full-length and truncated enzymes cleaved DNA at a similar spectrum of sites and displayed comparable rates of DNA ligation, we conclude that the enhanced double-stranded DNA cleavage observed for hTop2 $\alpha\Delta1175$  is due largely to an increase in active site coordination.

It is not obvious why the two active site tyrosyl residues of human topoisomerase  $II\alpha$  normally do not act in a concerted fashion or why the enzyme would evolve a C-terminal domain that contributed to this lack of coordination. One possibility is that this absence of coordination contributes to the preservation of genomic stability by decreasing the likelihood that topoisomerase  $II\alpha$  would generate double-stranded DNA breaks in the genetic material in the absence of strand passage.

Despite the high DNA scission activity of chlorella virus type II topoisomerases, these enzymes are relatively insensitive to the DNA cleavage-enhancing effects of anticancer drugs (29,31). Remarkably, hTop $2\alpha\Delta1175$  also displayed a distinctly lower susceptibility to anticancer agents. This loss of sensitivity does not result from a decreased affinity for drugs (at least for etoposide). Therefore, the C-terminal domain of topoisomerase II appears to play a role in the response of the enzyme to anticancer agents.

The C-terminal domain of human topoisomerase II $\alpha$  has long been known to play an important role in the physiological regulation of the enzyme. Results of the present study indicate that this region of the protein also plays a part in modulating topoisomerase II-mediated catalysis. The mechanistic basis for the contributions of the C-terminal domain to enzyme function is not known. It is clear from the properties of hTop2 $\alpha$  $\Delta$ 1175 that the C-terminal domain does not contain any essential "active site" amino acid residues. Thus, it is possible that this portion of the enzyme may act by altering the conformation of topoisomerase II $\alpha$ . Alternatively, based on the finding that the C-terminal domain of bacterial DNA gyrase plays a role in mediating enzyme-DNA binding, this region of the human enzyme may subtly alter interactions between the active site of topoisomerase II $\alpha$  and its nucleic acid substrate. Ultimately, structural studies will be required to address this important issue.

#### **ACKNOWLEDGEMENTS**

We would like to thank Ryan P. Bender for helpful discussions regarding the drug binding experiments, and Dr. Erin L. Baldwin and A. Kathleen McClendon for critical reading of the manuscript.

#### **REFERENCES**

- 1. Wang JC. DNA topoisomerases. Annu. Rev. Biochem 1996;65:635-692. [PubMed: 8811192]
- 2. Wang JC. Moving one DNA double helix through another by a type II DNA topoisomerase: the story of a simple molecular machine. Quart. Rev. Biophys 1998;31:107–144.
- 3. Burden DA, Osheroff N. Mechanism of action of eukaryotic topoisomerase II and drugs targeted to the enzyme. Biochim. Biophys. Acta 1998;1400:139–154. [PubMed: 9748545]
- 4. Nitiss JL. Investigating the biological functions of DNA topoisomerases in eukaryotic cells. Biochim. Biophys. Acta 1998;1400:63–81. [PubMed: 9748506]
- 5. Fortune JM, Osheroff N. Topoisomerase II as a target for anticancer drugs: when enzymes stop being nice. Prog. Nucleic Acid Res. Mol. Biol 2000;64:221–253. [PubMed: 10697411]
- Champoux JJ. DNA TOPOISOMERASES: Structure, Function, and Mechanism. Annu. Rev. Biochem 2001;70:369–413. [PubMed: 11395412]

7. Wilstermann AM, Osheroff N. Stabilization of eukaryotic topoisomerase II-DNA cleavage complexes. Curr. Top. Med. Chem 2003;3:321–338. [PubMed: 12570766]

- 8. Drake FH, Zimmerman JP, McCabe FL, Bartus HF, Per SR, Sullivan DM, Ross WE, Mattern MR, Johnson RK, Crooke ST. Purification of topoisomerase II from amsacrine-resistant P388 leukemia cells. Evidence for two forms of the enzyme. J. Biol. Chem 1987;262:16739–16747. [PubMed: 2824504]
- Drake FH, Hofmann GA, Bartus HF, Mattern MR, Crooke ST, Mirabelli CK. Biochemical and pharmacological properties of p170 and p180 forms of topoisomerase II. Biochemistry 1989;28:8154– 8160. [PubMed: 2557897]
- 10. Hsiang YH, Wu HY, Liu LF. Proliferation-dependent regulation of DNA topoisomerase II in cultured human cells. Cancer Res 1988;48:3230–3235. [PubMed: 2835157]
- 11. Isaacs RJ, Davies SL, Sandri MI, Redwood C, Wells NJ, Hickson ID. Physiological regulation of eukaryotic topoisomerase II. Biochim. Biophys. Acta 1998;1400:121–137. [PubMed: 9748535]
- 12. Austin CA, Marsh KL. Eukaryotic DNA topoisomerase IIβ. Bioessays 1998;20:215–226. [PubMed: 9631649]
- 13. Woessner RD, Mattern MR, Mirabelli CK, Johnson RK, Drake FH. Proliferation- and cell cycle-dependent differences in expression of the 170 kilodalton and 180 kilodalton forms of topoisomerase II in NIH-3T3 cells. Cell Growth Differ 1991;2:209–214. [PubMed: 1651102]
- Yang X, Li W, Prescott ED, Burden SJ, Wang JC. DNA topoisomerase IIb and neural development. Science 2000;287:131–134. [PubMed: 10615047]
- 15. Berger JM, Gamblin SJ, Harrison SC, Wang JC. Structure and mechanism of DNA topoisomerase II. Nature 1996;379:225–232. [PubMed: 8538787]
- Baguley BC, Ferguson LR. Mutagenic properties of topoisomerase-targeted drugs. Biochim. Biophys. Acta 1998;1400:213–222. [PubMed: 9748584]
- Felix CA. Secondary leukemias induced by topoisomerase-targeted drugs. Biochim. Biophys. Acta 1400 1998:233–255.
- 18. Kaufmann SH. Cell death induced by topoisomerase-targeted drugs: more questions than answers. Biochim. Biophys. Acta 1998;1400:195–211. [PubMed: 9748575]
- Pommier, Y.; Fesen, MR.; Goldwasser, F. Topoisomerase II inhibitors: the epipodophyllotoxins, m-AMSA, and the ellipticine derivatives. In: Chabner, BA.; Longo, DL., editors. Cancer Chemotherapy and Biotherapy: Principles and Practice. Lippincott-Raven Publishers; Philadelphia: 1996. p. 435-461.
- 20. Hande KR. Clinical applications of anticancer drugs targeted to topoisomerase II. Biochim. Biophys. Acta 1998;1400:173–184. [PubMed: 9748560]
- 21. Hande KR. Etoposide: four decades of development of a topoisomerase II inhibitor. Eur. J. Cancer 1998;34:1514–1521. [PubMed: 9893622]
- 22. Rowley JD. The critical role of chromosome translocations in human leukemias. Annu. Rev. Genet 1998;32:495–519. [PubMed: 9928489]
- 23. Felix CA. Leukemias related to treatment with DNA topoisomerase II inhibitors. Med. Pediatr. Oncol 2001;36:525–535. [PubMed: 11340607]
- 24. Slater DJ, Hilgenfeld E, Rappaport EF, Shah N, Meek RG, Williams WR, Lovett BD, Osheroff N, Autar RS, Ried T, Felix CA. MLL-SEPTIN6 fusion recurs in novel translocation of chromosomes 3, X, and 11 in infant acute myelomonocytic leukaemia and in t(X;11) in infant acute myeloid leukaemia, and MLL genomic breakpoint in complex MLL-SEPTIN6 rearrangement is a DNA topoisomerase II cleavage site. Oncogene 2002;21:4706–4714. [PubMed: 12096348]
- 25. Lindsey RH Jr. Bromberg KD, Felix CA, Osheroff N. 1,4- Benzoquinone is a topoisomerase II poison. Biochemistry 2004;43:7563–7574. [PubMed: 15182198]
- 26. Mistry AR, Felix CA, Whitmarsh RJ, Mason A, Reiter A, Cassinat B, Parry A, Walz C, Wiemels JL, Segal MR, Ades L, Blair IA, Osheroff N, Peniket AJ, Lafage-Pochitaloff M, Cross NC, Chomienne C, Solomon E, Fenaux P, Grimwade D. DNA topoisomerase II in therapy-related acute promyelocytic leukemia. N. Engl. J. Med 2005;352:1529–1538. [PubMed: 15829534]
- 27. Pedersen-Bjergaard J. Insights into leukemogenesis from therapy-related leukemia. N. Engl. J. Med 2005;352:1591–1594. [PubMed: 15829541]

28. Lavrukhin OV, Fortune JM, Wood TG, Burbank DE, Van Etten JL, Osheroff N, Lloyd RS. Topoisomerase II from Chlorella virus PBCV-1. Characterization of the smallest known type II topoisomerase. J. Biol. Chem 2000;275:6915–6921. [PubMed: 10702252]

- Fortune JM, Lavrukhin OV, Gurnon JR, Van Etten JL, Lloyd RS, Osheroff N. Topoisomerase II from Chlorella virus PBCV-1 has an exceptionally high DNA cleavage activity. J. Biol. Chem 2001;276:24401–24408. [PubMed: 11323425]
- 30. Fortune JM, Dickey JS, Lavrukhin OV, Van Etten JL, Lloyd RS, Osheroff N. Site-specific DNA cleavage by Chlorella virus topoisomerase II. Biochemistry 2002;41:11761–11769. [PubMed: 12269818]
- 31. Dickey JS, Choi TJ, Van Etten JL, Osheroff N. Chlorella virus Marburg topoisomerase II: high DNA cleavage activity as a characteristic of Chlorella virus type II enzymes. Biochemistry 2005;44:3899–3908. [PubMed: 15751965]
- 32. Mirski SE, Gerlach JH, Cummings HJ, Zirngibl R, Greer PA, Cole SP. Bipartite nuclear localization signals in the C terminus of human topoisomerase IIα. Exp. Cell Res 1997;237:452–455. [PubMed: 9434641]
- 33. Jensen S, Andersen AH, Kjeldsen E, Biersack H, Olsen EH, Andersen TB, Westergaard O, Jakobsen BK. Analysis of functional domain organization in DNA topoisomerase II from humans and Saccharomyces cerevisiae. Mol. Cell. Biol 1996;16:3866–3877. [PubMed: 8668204]
- 34. Corbett KD, Shultzaberger RK, Berger JM. The C-terminal domain of DNA gyrase A adopts a DNA-bending beta-pinwheel fold. Proc. Natl. Acad. Sci. U S A 2004;101:7293–7298. [PubMed: 15123801]
- 35. Qi Y, Pei J, Grishin NV. C-terminal domain of gyrase A is predicted to have a beta-propeller structure. Proteins 2002;47:258–264. [PubMed: 11948780]
- 36. Hsieh TJ, Chan NL. Crystallization and preliminary X-ray crystallographic analysis of the C-terminal domain of ParC protein from Bacillus stearothermophilus. Acta Crystallogr. D. Biol. Crystallogr 2004;60:564–566. [PubMed: 14993694]
- 37. Hsieh TJ, Farh L, Huang WM, Chan NL. Structure of the topoisomerase IV C-terminal domain: a broken beta-propeller implies a role as geometry facilitator in catalysis. J. Biol. Chem 2004;279:55587–55593. [PubMed: 15466871]
- 38. Shiozaki K, Yanagida M. A functional 125-kDa core polypeptide of fission yeast DNA topoisomerase II. Mol. Cell. Biol 1991;11:6093–6102. [PubMed: 1658625]
- 39. Crenshaw DG, Hsieh T. Function of the hydrophilic carboxyl terminus of type II DNA topoisomerase from Drosophila melanogaster. I. In vitro studies. J. Biol. Chem 1993;268:21328–21334. [PubMed: 8407971]
- 40. Crenshaw DG, Hsieh T. Function of the hydrophilic carboxyl terminus of type II DNA topoisomerase from Drosophila melanogaster. II. In vivo studies. J. Biol. Chem 1993;268:21335–21343. [PubMed: 8407972]
- 41. Caron PR, Watt P, Wang JC. The C-terminal domain of Saccharomyces cerevisiae DNA topoisomerase II. Mol. Cell. Biol. May 1994:3197–3207.
- 42. Englund PT. The replication of kinetoplast DNA networks in *Crithidia fasciculata*. Cell 1978;14:157–168. [PubMed: 667931]
- 43. Wasserman RA, Austin CA, Fisher LM, Wang JC. Use of yeast in the study of anticancer drugs targeting DNA topoisomerases: expression of a functional recombinant human DNA topoisomerase II alpha in yeast. Cancer Res 1993;53:3591–3596. [PubMed: 8393377]
- 44. Kingma PS, Greider CA, Osheroff N. Spontaneous DNA lesions poison human topoisomerase IIα and stimulate cleavage proximal to leukemic 11q23 chromosomal breakpoints. Biochemistry 1997;36:5934–5939. [PubMed: 9166762]
- 45. Miller KG, Liu LF, Englund PT. A homogeneous type II DNA topoisomerase from HeLa cell nuclei. J. Biol. Chem 1981;256:9334–9339. [PubMed: 6267071]
- 46. Fortune JM, Osheroff N. Merbarone inhibits the catalytic activity of human topoisomerase II $\alpha$  by blocking DNA cleavage. J. Biol. Chem 1998;273:17643–17650. [PubMed: 9651360]
- 47. Baldwin EL, Byl JA, Osheroff N. Cobalt enhances DNA cleavage mediated by human topoisomerase II alpha in vitro and in cultured cells. Biochemistry 2004;43:728–735. [PubMed: 14730977]

48. O'Reilly EK, Kreuzer KN. A unique type II topoisomerase mutant that is hypersensitive to a broad range of cleavage-inducing antitumor agents. Biochemistry 2002;41:7989–7997. [PubMed: 12069589]

- 49. Robinson MJ, Osheroff N. Stabilization of the topoisomerase II-DNA cleavage complex by antineoplastic drugs: inhibition of enzyme-mediated DNA religation by 4'-(9-acridinylamino) methanesulfon-m-anisidide. Biochemistry 1990;29:2511–2515. [PubMed: 2159323]
- Kingma PS, Burden DA, Osheroff N. Binding of etoposide to topoisomerase II in the absence of DNA: decreased affinity as a mechanism of drug resistance. Biochemistry 1999;38:3457–3461.
   [PubMed: 10090731]
- 51. Osheroff N. Effect of antineoplastic agents on the DNA cleavage/religation reaction of eukaryotic topoisomerase II: inhibition of DNA religation by etoposide. Biochemistry 1989;28:6157–6160. [PubMed: 2551366]
- 52. Bromberg KD, Velez-Cruz R, Burgin AB, Osheroff N. DNA ligation catalyzed by human topoisomerase II alpha. Biochemistry 2004;43:13416–13423. [PubMed: 15491148]
- 53. Reece RJ, Maxwell A. Tryptic fragments of the Escherichia coli DNA gyrase A protein. J. Biol. Chem 1989;264:19648–19653. [PubMed: 2555327]
- 54. Sengupta T, Mukherjee M, Mandal C, Das A, Majumder HK. Functional dissection of the C-terminal domain of type II DNA topoisomerase from the kinetoplastid hemoflagellate Leishmania donovani. Nucleic Acids Res 2003;31:5305–5316. [PubMed: 12954766]
- 55. Niimi A, Suka N, Harata M, Kikuchi A, Mizuno S. Co-localization of chicken DNA topoisomerase IIalpha, but not beta, with sites of DNA replication and possible involvement of a C-terminal region of alpha through its binding to PCNA. Chromosoma 2001;110:102–114. [PubMed: 11453553]






Figure 1. Decatenation of kDNA by human topoisomerase IIα and hTop2α $\Delta$ 1175. The percent decatenation was determined by quantifying minicircles that were released from kDNA networks by human topoisomerase IIα (hTop2α, open circles) or hTop2α $\Delta$ 1175 ( $\Delta$ 1175, closed circles) and is shown on the left. Error bars represent the standard deviation of three independent experiments. The image at the right is an ethidium bromide-stained gel showing kDNA decatenation in the absence of enzyme (DNA), or in the presence of human topoisomerase IIα or hTop2α $\Delta$ 1175. Reaction mixtures contained 4 nM enzyme. Positions of the kDNA networks at the origin, as well as those of the released DNA maxicircles and minicircles, are indicated.



## **hTop2αΔ1175**

Figure 2. Relaxation of negatively supercoiled pBR322 plasmid DNA by topoisomerase II $\alpha$  and hTop2 $\alpha\Delta$ 1175. The relaxation of negatively supercoiled pBR322 plasmid DNA by human topoisomerase II $\alpha$  (hTop2 $\alpha$ , top) and hTop2 $\alpha\Delta$ 1175 (bottom) is shown. Relaxation reactions were carried out from 0-30 min using 1 nM enzyme. Ethidium bromide-stained gels that are representative of 3 independent experiments are shown. Negatively supercoiled DNA (form I, FI) and nicked circular DNA (form II, FII) are indicated. A DNA control in the absence of enzyme is shown (DNA).

# hTop2α



## **hTop2αΔ1175**

Figure 3. Non-turnover relaxation of negatively supercoiled pBR322 plasmid DNA by topoisomerase II $\alpha$  and hTop2 $\alpha\Delta$ 1175. The non-turnover relaxation of negatively supercoiled pBR322 plasmid DNA by human topoisomerase II $\alpha$  (hTop2 $\alpha$ , top) and hTop2 $\alpha\Delta$ 1175 (bottom) is shown. Relaxation reactions were carried out from 0-60 s using 110 nM enzyme and 5 nM DNA. Ethidium bromide-stained gels that are representative of 3 independent experiments are shown. Negatively supercoiled DNA (form I, FI) and nicked circular DNA (form II, FII) are indicated. A DNA control in the absence of enzyme is shown (DNA).




Figure 4. Double-stranded DNA cleavage by type II topoisomerases. The ability of human topoisomerase IIα (hTop2α, open circles) and hTop2αΔ1175 (Δ1175, closed circles) to cleave negatively supercoiled pBR322 plasmid DNA is shown. A range of enzyme (0.2-1 μM) was used at a constant concentration of 10 nM plasmid. Double-stranded DNA cleavage is expressed as the percentage of circular plasmid substrate that is converted to linear molecules. Error bars represent the standard deviation of three independent experiments. The inset shows an ethidium bromide-stained agarose gel of a DNA cleavage reaction in the absence of enzyme (DNA), in the presence of 0.6 μM human topoisomerase IIα (hTop2α) or hTop2αΔ1175 (Δ1175), or 0.02 μM PBCV-1 topoisomerase II (PBCV). Double-stranded DNA cleavage

converts negatively supercoiled plasmid (form I, FI) to linear molecules (form III, FIII). The position of nicked circular DNA (form II, FII) also is indicated.




Figure 5. Binding of topoisomerase II $\alpha$  and hTop2 $\alpha\Delta$ 1175 to DNA. The percent binding of radiolabeled linear pBR322 DNA to human topoisomerase II $\alpha$  (hTop2 $\alpha$ , open circles) or hTop2 $\alpha\Delta$ 1175 ( $\Delta$ 1175, closed circles) was determined by the ratio of radioactivity retained on a nitrocellulose filter vs. the input amount of radioactivity. A range of 0-400 nM enzyme was examined at a constant concentration of 5 nM linear pBR322. Error bars represent the standard deviation of three independent experiments.




Figure 6. DNA religation mediated by topoisomerase IIα and hTop2α $\Delta$ 1175. DNA religation mediated by human topoisomerase IIα (hTop2α, open circles) or hTop2α $\Delta$ 1175 ( $\Delta$ 1175, closed circles) is shown. The DNA cleavage/religation equilibrium was established at 37°C and religation was initiated by shifting samples to 0°C. The DNA cleavage observed at equilibrium for each enzyme was set to 100% at time zero. Error bars represent the standard deviation of three independent experiments.




Figure 7. The ratio of single-stranded to double-stranded DNA breaks generated by type II topoisomerases. The ratio of single-stranded DNA breaks to double-stranded DNA breaks generated by human topoisomerase IIa (hTop2a, open bars), hTop2a $\Delta$ 1175 ( $\Delta$ 1175, closed bars), or PBCV-1 topoisomerase II (PBCV-1, shaded bar) is shown. A range of 0.2-1  $\mu$ M enzyme was examined for the human enzymes, and the ratio of breaks produced by 0.02  $\mu$ M PBCV-1 topoisomerase II is shown for comparison. Double-stranded and single-stranded DNA cleavage is expressed as the percentage of supercoiled substrate that is converted to linear or nicked circular molecules, respectively. Error bars represent the standard deviation of 3 independent experiments.




Figure 8. Etoposide binding to type II topoisomerases. The ability of 3  $\mu$ M human topoisomerase II $\alpha$  (hTop2 $\alpha$ , left panel), hTop2 $\alpha\Delta$ 1175 ( $\Delta$ 1175, middle panel) or PBCV-1 topoisomerase II (PBCV-1, right panel) to bind [ $^3$ H]etoposide was examined. The free and bound fractions of etoposide were separated by filtration through nitrocellulose membranes, and the binding of etoposide to the filter in the absence of enzyme was subtracted as background. Error bars represent the standard deviation of at least 3 independent experiments.




Figure 9. Double-stranded DNA cleavage by type II enzymes in the presence of topoisomerase II poisons. The ability of human topoisomerase II $\alpha$  (hTop2 $\alpha$ , open circles) and hTop2 $\alpha$ Δ1175 (Δ1175, closed circles) to cleave negatively supercoiled pBR322 plasmid was determined in the presence of three topoisomerase II-targeted anticancer drugs, etoposide, amsacrine, and CP-115,953. Relative DNA cleavage is based on levels of double-stranded DNA breaks generated by the respective enzymes in the absence of drugs. For each individual enzyme, DNA cleavage in the absence of drugs was set to 1. Error bars represent the standard deviation of three independent experiments.




Figure 10. Distribution of topoisomerase II-mediated DNA cleavage sites on a linear DNA substrate. A singly end-labeled 4330 bp fragment of pBR322 was employed. The autoradiogram shows a DNA control (DNA) and cleavage in the presence of human topoisomerase II $\alpha$  (hTop2 $\alpha$ ), topoisomerase II $\alpha$  plus 50  $\mu$ M etoposide (hTop2 $\alpha$ /Etop), hTop2 $\alpha$ Δ1175 (Δ1175), or hTop2 $\alpha$ Δ1175 plus 50  $\mu$ M etoposide (Δ1175/Etop). Data are representative of three independent experiments.